

## Cytokine Receptor Common Gamma Chain Like

### *Priority*

This application claims priority to U.S. Application Serial No. 09/263,626,  
5 filed March 5, 1999, which claims priority to U.S. Provisional Application Serial No.  
60/078,563, filed March 19, 1998, and to U.S. Provisional Application Serial No.  
60/086,505, filed May 22, 1998, and PCT Application No. PCT/US99/05068, filed  
March 5, 1999, all herein incorporated by reference in their entirety.

### *Field of the Invention*

10 The present invention relates to a novel human gene encoding a polypeptide  
which is a member of the Cytokine Receptor family. More specifically, the present  
invention relates to a polynucleotide encoding a novel human polypeptide named  
Cytokine Receptor Common Gamma Chain Like, or "CRCGCL." This invention also  
relates to CRCGCL polypeptides, as well as vectors, host cells, antibodies directed to  
15 CRCGCL polypeptides, and the recombinant methods for producing the same. Also  
provided are diagnostic methods for detecting disorders related to the immune system,  
and therapeutic methods for treating such disorders. The invention further relates to  
screening methods for identifying agonists and antagonists of CRCGCL activity.

### *Background of the Invention*

20 Biochemical and physiological effects often result from the binding of a  
cytokine to a specific receptor molecule. Receptor binding then stimulates certain,  
and often independent, signal transduction pathways. (Kishimoto, T., et al., Cell  
76:253-262 (1994.) The interaction between a cytokine and a receptor is a primary  
25 regulator of a variety of cellular processes, including activation, proliferation, and  
differentiation. (Arai, K. -I, et al., Ann. Rev. Biochem. 59:783-836 (1990); Paul, W.  
E. and Seder, R. A., Cell 76:241-251 (1994)).

Cytokines that bind to the interleukin-2 (IL-2) receptor common gamma chain  
(gamma c), including IL-2, IL-4, IL-7, IL-9, and IL-15, are important for the growth  
30 and differentiation of immune cells, such as T and B lymphocytes, natural killer cells,  
macrophages, and monocytes. These cytokines have overlapping biological effects  
that in part result from the use of the shared receptor subunit gamma c. Recently it

has been shown that these cytokines activate a number of important intracellular signaling molecules, by binding to the interleukin-2 (IL-2) receptor common gamma chain (gamma c), including the Janus kinases JAK1 and JAK3 and members of the transcription factor family of signal transducers and activators of transcription (STATs).

The discovery of these signaling pathways has led to important new insights into their role in lymphocyte maturation, as it has emerged that mutations in the genes encoding both gamma c and JAK3 result in similar forms of severe combined immunodeficiency (SCID). For example, mutations in the human interleukin-2 (IL-2) receptor gamma, mapped to the X chromosome, is associated with X-linked severe combined immunodeficiency. (Human Molecular Genetics, 2(8): 1099 (1993).)

Thus, there is a need for polypeptides that regulate the differentiation and proliferation of cells, since disturbances of such regulation may be involved in disorders relating to immune system. Therefore, there is a need for identification and characterization of such human polypeptides which can play a role in detecting, preventing, ameliorating or correcting such disorders.

### ***Summary of the Invention***

The present invention relates to a novel polynucleotide and the encoded polypeptide of CRCGCL. Moreover, the present invention relates to vectors, host cells, antibodies, and recombinant methods for producing the polypeptides and polynucleotides. Also provided are diagnostic methods for detecting disorders relates to the polypeptides, and therapeutic methods for treating such disorders. The invention further relates to screening methods for identifying binding partners of CRCGCL.

### ***Brief Description of the Drawings***

Figures 1A-1B show the nucleotide sequence (SEQ ID NO:1) and the deduced amino acid sequence (SEQ ID NO:2) of CRCGCL. The predicted leader sequence is located at about amino acids 1-22.

Figure 2 shows the regions of identity between the amino acid sequence of the CRCGCL protein and the translation product of the closest homolog, the Bos Taurus Interleukin-2 receptor gamma (Accession Nos. 1532088) (SEQ ID NO:3), determined by BLAST analysis. Identical amino acids between the two polypeptides are shaded in black, while conservative amino acids are boxed. By examining the regions of amino acids shaded and/or boxed, the skilled artisan can readily identify conserved domains between the two polypeptides. These conserved domains are preferred embodiments of the present invention.

Figure 3 shows an analysis of the CRCGCL amino acid sequence. Alpha, beta, turn and coil regions; hydrophilicity and hydrophobicity; amphipathic regions; flexible regions; antigenic index and surface probability are shown, and all were generated using the default settings. In the "Antigenic Index or Jameson-Wolf" graph, the positive peaks indicate locations of the highly antigenic regions of the CRCGCL protein, i.e., regions from which epitope-bearing peptides of the invention can be

obtained. The domains defined by these graphs are contemplated by the present invention. Tabular representation of the data summarized graphically in Figure 3 can be found in Table 1. The columns are labeled with the headings "Res", "Position", and Roman Numerals I-XIV. The column headings refer to the following features of the amino acid sequence presented in Figure 3, and Table 1: "Res": amino acid residue of SEQ ID NO:2 and Figures 1A and 1B; "Position": position of the corresponding residue within SEQ ID NO:2 and Figures 1A and 1B; I: Alpha, Regions - Garnier-Robson; II: Alpha, Regions - Chou-Fasman; III: Beta, Regions - Garnier-Robson; IV: Beta, Regions - Chou-Fasman; V: Turn, Regions - Garnier-Robson; VI: Turn, Regions - Chou-Fasman; VII: Coil, Regions - Garnier-Robson; VIII: Hydrophilicity Plot - Kyte-Doolittle; IX: Hydrophobicity Plot - Hopp-Woods; X: Alpha, Amphipathic Regions - Eisenberg; XI: Beta, Amphipathic Regions - Eisenberg; XII: Flexible Regions - Karplus-Schulz; XIII: Antigenic Index - Jameson-Wolf; and XIV: Surface Probability Plot - Emini.

0536430.0619.60

## *Detailed Description*

### Definitions

5           The following definitions are provided to facilitate understanding of certain terms used throughout this specification.

          In the present invention, "isolated" refers to material removed from its original environment (e.g., the natural environment if it is naturally occurring), and thus is altered "by the hand of man" from its natural state. For example, an isolated  
10   polynucleotide could be part of a vector or a composition of matter, or could be contained within a cell, and still be "isolated" because that vector, composition of matter, or particular cell is not the original environment of the polynucleotide.

          In the present invention, a "secreted" CRCGCL protein refers to a protein capable of being directed to the ER, secretory vesicles, or the extracellular space as a  
15   result of a signal sequence, as well as a CRCGCL protein released into the extracellular space without necessarily containing a signal sequence. If the CRCGCL secreted protein is released into the extracellular space, the CRCGCL secreted protein can undergo extracellular processing to produce a "mature" CRCGCL protein. Release into the extracellular space can occur by many mechanisms, including  
20   exocytosis and proteolytic cleavage.

          As used herein, a CRCGCL "polynucleotide" refers to a molecule having a nucleic acid sequence contained in SEQ ID NO:1 or the cDNA contained within the clone deposited with the ATCC. For example, the CRCGCL polynucleotide can contain the nucleotide sequence of the full length cDNA sequence, including the 5'  
25   and 3' untranslated sequences, the coding region, with or without the signal sequence, the secreted protein coding region, as well as fragments, epitopes, domains, and variants of the nucleic acid sequence. Moreover, as used herein, a CRCGCL "polypeptide" refers to a molecule having the translated amino acid sequence generated from the polynucleotide as broadly defined. However, one embodiment of  
30   the present invention does not include the polynucleotide sequence of Genbank Accession No. X91553, herein incorporated by reference.

668189-0349260



In specific embodiments, the polynucleotides of the invention are less than 300 kb, 200 kb, 100 kb, 50 kb, 15 kb, 10 kb, or 7.5 kb in length. In a further embodiment, polynucleotides of the invention comprise at least 15 contiguous nucleotides of CRCGCL coding sequence, but do not comprise all or a portion of any  
5 CRCGCL intron. In another embodiment, the nucleic acid comprising CRCGCL coding sequence does not contain coding sequences of a genomic flanking gene (i.e., 5' or 3' to the CRCGCL gene in the genome).

In the present invention, the full length CRCGCL sequence identified as SEQ ID NO:1 was generated by overlapping sequences of the deposited clone (contig  
10 analysis). A representative clone containing all or most of the sequence for SEQ ID NO:1 was deposited with the American Type Culture Collection ("ATCC") on March 23, 1998, and was given the ATCC Deposit Number 209691. A second clone was also deposited with the ATCC on February 25, 1998, and given ATCC Deposit Number 209641. The ATCC is located at 10801 University Boulevard, Manassas,  
15 VA 20110-2209, USA. The ATCC deposit was made pursuant to the terms of the Budapest Treaty on the international recognition of the deposit of microorganisms for purposes of patent procedure.

A CRCGCL "polynucleotide" also includes those polynucleotides capable of hybridizing, under stringent hybridization conditions, to sequences contained in SEQ  
20 ID NO:1, the complement thereof, or the cDNA within the deposited clone. "Stringent hybridization conditions" refers to an overnight incubation at 42 degree C in a solution comprising 50% formamide, 5x SSC (750 mM NaCl, 75 mM sodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA, followed by washing  
25 the filters in 0.1x SSC at about 65 degree C.

Also contemplated are nucleic acid molecules that hybridize to the CRCGCL polynucleotides at moderately high stringency hybridization conditions. Changes in the stringency of hybridization and signal detection are primarily accomplished through the manipulation of formamide concentration (lower percentages of  
30 formamide result in lowered stringency); salt conditions, or temperature. For example, moderately high stringency conditions include an overnight incubation at 37

09376430-081899

degree C in a solution comprising 6X SSPE (20X SSPE = 3M NaCl; 0.2M NaH<sub>2</sub>PO<sub>4</sub>; 0.02M EDTA, pH 7.4), 0.5% SDS, 30% formamide, 100 ug/ml salmon sperm blocking DNA; followed by washes at 50 degree C with 1XSSPE, 0.1% SDS. In addition, to achieve even lower stringency, washes performed following stringent hybridization can be done at higher salt concentrations (e.g. 5X SSC).

Note that variations in the above conditions may be accomplished through the inclusion and/or substitution of alternate blocking reagents used to suppress background in hybridization experiments. Typical blocking reagents include Denhardt's reagent, BLOTTO, heparin, denatured salmon sperm DNA, and commercially available proprietary formulations. The inclusion of specific blocking reagents may require modification of the hybridization conditions described above, due to problems with compatibility.

Of course, a polynucleotide which hybridizes only to polyA<sup>+</sup> sequences (such as any 3' terminal polyA<sup>+</sup> tract of a cDNA shown in the sequence listing), or to a complementary stretch of T (or U) residues, would not be included in the definition of "polynucleotide," since such a polynucleotide would hybridize to any nucleic acid molecule containing a poly (A) stretch or the complement thereof (e.g., practically any double-stranded cDNA clone).

The CRCGCL polynucleotide can be composed of any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. For example, CRCGCL polynucleotides can be composed of single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, the CRCGCL polynucleotides can be composed of triple-stranded regions comprising RNA or DNA or both RNA and DNA. CRCGCL polynucleotides may also contain one or more modified bases or DNA or RNA backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of

modifications can be made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically, or metabolically modified forms.

CRCGCL polypeptides can be composed of amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres, and may contain amino acids other than the 20 gene-encoded amino acids. The CRCGCL polypeptides may be modified by either natural processes, such as posttranslational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications can occur anywhere in the CRCGCL polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given CRCGCL polypeptide. Also, a given CRCGCL polypeptide may contain many types of modifications. CRCGCL polypeptides may be branched, for example, as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched, and branched cyclic CRCGCL polypeptides may result from posttranslation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, pegylation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. (See, for instance, PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York (1993); POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS, B. C. Johnson, Ed., Academic

Press, New York, pgs. 1-12 (1983); Seifter et al., Meth Enzymol 182:626-646 (1990); Rattan et al., Ann NY Acad Sci 663:48-62 (1992).)

"SEQ ID NO:1" refers to a CRCGCL polynucleotide sequence while "SEQ ID NO:2" refers to a CRCGCL polypeptide sequence.

5 A CRCGCL polypeptide "having biological activity" refers to polypeptides exhibiting activity similar, but not necessarily identical to, an activity of a CRCGCL polypeptide, including mature forms, as measured in a particular biological assay, with or without dose dependency. In the case where dose dependency does exist, it need not be identical to that of the CRCGCL polypeptide, but rather substantially  
10 similar to the dose-dependence in a given activity as compared to the CRCGCL polypeptide (i.e., the candidate polypeptide will exhibit greater activity or not more than about 25-fold less and, preferably, not more than about tenfold less activity, and most preferably, not more than about three-fold less activity relative to the CRCGCL polypeptide.)

15

#### **CRCGCL Polynucleotides and Polypeptides**

Clone HTAEK53 was isolated from an activated T-cell cDNA library. Initially, the sequence of clone HTAEK53 was identified as SEQ IDNO:26 and the deduced amino acid sequence was predicted as SEQ ID NO:27, with a recognition  
20 that an apparent frame shift in the sequence existed ~~between nucleotides 256 and 277~~. This frame shift was easily resolved using standard molecular biology techniques, generating the nucleotide sequence of SEQ ID NO:1 and the deduced amino acid sequence shown in SEQ ID NO:2.

The deposited clone contains a cDNA having a total of 1573 nucleotides,  
25 which encodes a predicted open reading frame of 371 amino acid residues. (See Figures 1A-1B.) The open reading frame begins at a N-terminal methionine located at nucleotide position 13, and ends at a stop codon at nucleotide position 1128. The predicted molecular weight of the CRCGCL protein should be about 42 kDa.

Subsequent Northern analysis also showed a 1.6 Kb transcript in a cervical  
30 cancer cell line (HeLa), activated T cells, and a lung carcinoma cell line (A549),

00376430-001899

u

while a shorter variant is also expressed in the lymph node and to a lesser extent in the spleen tissues, a pattern consistent with immune specific expression.

CRCGCL expression was not observed in the following cell lines, HL60, K562, Molt-4, Raji, SW480, G361, as well as the heart, brain, placenta, lung, liver, skeletal muscle, kidney, pancreas, thymus, prostate, testis, ovary, small intestine, colon, or peripheral blood leukocytes, a pattern consistent with immune specific expression.

Using BLAST analysis, SEQ ID NO:2 was found to be homologous to members of the Cytokine Receptor family. Particularly, SEQ ID NO:2 contains domains homologous to the translation product of the Bos Taurus mRNA for Interleukin-2 receptor gamma (Accession Nos. 1532088) (Figure 2) (SEQ ID NO:3), including the following conserved domains: (a) a predicted transmembrane domain located at about amino acids 226-260; (b) a predicted WXWS (SEQ ID NO:30), or [STGL]-x-W-[SG]-x-W-S (SEQ ID NO:18), domain located at about amino acids 198-204 (T-x-P-S-x-W-S) (SEQ ID NO:19), although not a perfect match; and (c) a predicted Jak Box, having the motif W(P,E)X(V,I)P(N,S,D)P (SEQ ID NO:20), domain located at about amino acids 261-268 (I-P-X-V-P-D-P) (SEQ ID NO:21), although not a perfect match. These polypeptide fragments of CRCGCL are specifically contemplated in the present invention. Because Interleukin- 2 receptor gamma (Accession Nos. 1532088) is thought to be important as a cytokine receptor, the homology between Interleukin-2 receptor gamma (Accession Nos. 1532088) and CRCGCL suggests that CRCGCL may also be involved in the differentiation and proliferation of cells. CRCGCL is also homologous to other Interleukin-2 receptor gamma genes isolated from a variety of species, such as human (Accession No. gi/349632), Canis familiaris (Accession No. gi/517412), and mouse (pri/S37582).

Moreover, the encoded polypeptide has a predicted leader sequence located at about amino acids 1-22. (See Figures 1A-1B.) Also shown in Figures 1A-1B, one embodiment of the secreted form of CRCGCL encompasses about amino acids 23-371, amino acids 23-225, or amino acids 1-231. These polypeptide fragments of CRCGCL are specifically contemplated in the present invention.

Other preferred polypeptide fragments comprise the amino acids sequence:  
 QIQIIFYNLETVQVTWNASKYSRTNLTFFHYRFNGDEAYDQCTNYLLQEGHTS  
 GC (SEQ ID NO:22); RRHSLFLHQEWDAPRFHRKSLDGLLPETQF (SEQ ID  
 NO:23);

- 5 LLYEVQYRSPFDTEWQSKQENTCNVTIEGLDAEKCYSFWVRVKAMEDVYGP  
 DTYPSDWSEVTCWQRGEIRDACAETPTPPK (SEQ ID NO:24); and/or  
 MEDVYGPDTYPSDWSEVTCWQRGEIRDACAETPTPPKPKLSKFILISSLAILLM  
 VSLLLLSLWKLWRXKKFLXPSVPDPKSIFPGLFXIHQGNFQEWITDTQNV AHL  
 HKMAGAEQESGPEEPLVVQLAKTEAESPRMLDPQTEEKEASGGSLQLPHQPL  
 10 QGGDVVTIGGFTFVMNDRSYVA (SEQ ID NO:25), as well as fragments thereof.  
 Also preferred are polynucleotide fragments encoding these polypeptide fragments.

Because CRCGCL was isolated from activated T cells, nucleic acids of the  
 invention are useful as reagents for differential identification of the tissue(s) or cell  
 type(s) present in a biological sample and for diagnosis of immune disorders.

- 15 Similarly, polypeptides and antibodies directed to those polypeptides are useful to  
 provide immunological probes for differential identification of the tissue(s) or cell  
 type(s). For a number of disorders of the immune system, expression of this gene at  
 significantly higher or lower levels may be detected in certain tissues (e.g., cancerous  
 and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or  
 20 spinal fluid) taken from an individual having such a disorder, relative to the standard  
 gene expression level, i.e., the expression level in healthy tissue from an individual  
 not having the disorder.

- The tissue distribution in only activated T-cells and homology to the cytokine  
 receptors IL2 and IL13 suggests that this protein is a novel member of the cytokine  
 25 receptor family expressed specifically on T-cells. The tissue distribution of this gene  
 in cells of the immune system suggests that the protein product of this clone would be  
 useful for treatment, prophylaxis and diagnosis of immune and autoimmune diseases,  
 such as lupus, transplant rejection, allergic reactions, arthritis, asthma,  
 immunodeficiency diseases, leukemia, AIDS. In addition its expression in T-cells  
 30 suggests a potential role in the treatment, prophylaxis and detection of thymus  
 disorders such as Graves Disease, lymphocytic thyroiditis, hyperthyroidism and

hypothyroidism. The receptor could also serve as a target for small molecule or monoclonal antibody, blocking its activity, which could be important in the disease states listed herein.

5 The CRCGCL nucleotide sequence identified as SEQ ID NO:1 was assembled from partially homologous ("overlapping") sequences obtained from the deposited clone, and in some cases, from additional related DNA clones. The overlapping sequences were assembled into a single contiguous sequence of high redundancy (usually three to five overlapping sequences at each nucleotide position), resulting in a final sequence identified as SEQ ID NO:1.

10 Therefore, SEQ ID NO:1 and the translated SEQ ID NO:2 are sufficiently accurate and otherwise suitable for a variety of uses well known in the art and described further below. For instance, SEQ ID NO:1 is useful for designing nucleic acid hybridization probes that will detect nucleic acid sequences contained in SEQ ID NO:1 or the cDNA contained in the deposited clone. These probes will also hybridize  
15 to nucleic acid molecules in biological samples, thereby enabling a variety of forensic and diagnostic methods of the invention. Similarly, polypeptides identified from SEQ ID NO:2 may be used to generate antibodies which bind specifically to CRCGCL.

20 Nevertheless, DNA sequences generated by sequencing reactions can contain sequencing errors. The errors exist as misidentified nucleotides, or as insertions or deletions of nucleotides in the generated DNA sequence. The erroneously inserted or deleted nucleotides cause frame shifts in the reading frames of the predicted amino acid sequence. In these cases, the predicted amino acid sequence diverges from the actual amino acid sequence, even though the generated DNA sequence may be greater than 99.9% identical to the actual DNA sequence (for example, one base insertion or  
25 deletion in an open reading frame of over 1000 bases).

Accordingly, for those applications requiring precision in the nucleotide sequence or the amino acid sequence, the present invention provides not only the generated nucleotide sequence identified as SEQ ID NO:1 and the predicted translated amino acid sequence identified as SEQ ID NO:2, but also a sample of plasmid DNA  
30 containing a human cDNA of CRCGCL deposited with the ATCC. The nucleotide sequence of the deposited CRCGCL clone can readily be determined by sequencing

09376430-001800

the deposited clone in accordance with known methods. The predicted CRCGCL amino acid sequence can then be verified from such deposits. Moreover, the amino acid sequence of the protein encoded by the deposited clone can also be directly determined by peptide sequencing or by expressing the protein in a suitable host cell containing the deposited human CRCGCL cDNA, collecting the protein, and determining its sequence.

The present invention also relates to the CRCGCL gene corresponding to SEQ ID NO:1, SEQ ID NO:2, or the deposited clone. The CRCGCL gene can be isolated in accordance with known methods using the sequence information disclosed herein. Such methods include preparing probes or primers from the disclosed sequence and identifying or amplifying the CRCGCL gene from appropriate sources of genomic material.

Also provided in the present invention are species homologs of CRCGCL. Species homologs may be isolated and identified by making suitable probes or primers from the sequences provided herein and screening a suitable nucleic acid source for the desired homologue.

The CRCGCL polypeptides can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

The CRCGCL polypeptides may be in the form of the secreted protein, including the mature form, or may be a part of a larger protein, such as a fusion protein (see below). It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, pro-sequences, sequences which aid in purification, such as multiple histidine residues, or an additional sequence for stability during recombinant production.

CRCGCL polypeptides are preferably provided in an isolated form, and preferably are substantially purified. A recombinantly produced version of a CRCGCL polypeptide, including the secreted polypeptide, can be substantially purified by the one-step method described in Smith and Johnson, Gene 67:31-40



(1988). CRCGCL polypeptides also can be purified from natural or recombinant sources using antibodies of the invention raised against the CRCGCL protein in methods which are well known in the art.

## 5 Polynucleotide and Polypeptide Variants

"Variant" refers to a polynucleotide or polypeptide differing from the CRCGCL polynucleotide or polypeptide, but retaining essential properties thereof. Generally, variants are overall closely similar, and, in many regions, identical to the CRCGCL polynucleotide or polypeptide.

10 By a polynucleotide having a nucleotide sequence at least, for example, 95% "identical" to a reference nucleotide sequence of the present invention, it is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence encoding the  
15 CRCGCL polypeptide. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. The query sequence  
20 may be an entire sequence shown of SEQ ID NO:1, the ORF (open reading frame), or any fragment specified as described herein.

a As a practical matter, whether any particular nucleic acid molecule or polypeptide is at least 90%, 95%, 96%, 97%, 98% or 99% identical to a nucleotide sequence of the <sup>present</sup> ~~presence~~ invention can be determined conventionally using known  
25 computer programs. A preferred method for determining the best overall match between a query sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci. (1990) 6:237-245.) In a sequence alignment the query and subject sequences  
30 are both DNA sequences. An RNA sequence can be compared by converting U's to T's. The result of said global sequence alignment is in percent identity. Preferred

0937430-019260

parameters used in a FASTDB alignment of DNA sequences to calculate percent identity are: Matrix=Unitary, k-tuple=4, Mismatch Penalty=1, Joining Penalty=30, Randomization Group Length=0, Cutoff Score=1, Gap Penalty=5, Gap Size Penalty 0.05, Window Size=500 or the length of the subject nucleotide sequence, whichever is shorter.

If the subject sequence is shorter than the query sequence because of 5' or 3' deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for 5' and 3' truncations of the subject sequence when calculating percent identity. For subject sequences truncated at the 5' or 3' ends, relative to the query sequence, the percent identity is corrected by calculating the number of bases of the query sequence that are 5' and 3' of the subject sequence, which are not matched/aligned, as a percent of the total bases of the query sequence. Whether a nucleotide is matched/aligned is determined by results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This corrected score is what is used for the purposes of the present invention. Only bases outside the 5' and 3' bases of the subject sequence, as displayed by the FASTDB alignment, which are not matched/aligned with the query sequence, are calculated for the purposes of manually adjusting the percent identity score.

For example, a 90 base subject sequence is aligned to a 100 base query sequence to determine percent identity. The deletions occur at the 5' end of the subject sequence and therefore, the FASTDB alignment does not show a matched/alignment of the first 10 bases at 5' end. The 10 unpaired bases represent 10% of the sequence (number of bases at the 5' and 3' ends not matched/total number of bases in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 bases were perfectly matched the final percent identity would be 90%. In another example, a 90 base subject sequence is compared with a 100 base query sequence. This time the deletions are internal deletions so that there are no bases on the 5' or 3' of the subject sequence which are not matched/aligned with the query. In this case the percent

identity calculated by FASTDB is not manually corrected. Once again, only bases 5' and 3' of the subject sequence which are not matched/aligned with the query sequence are manually corrected for. No other manual corrections are to made for the purposes of the present invention.

5 By a polypeptide having an amino acid sequence at least, for example, 95% "identical" to a query amino acid sequence of the present invention, it is intended that the amino acid sequence of the subject polypeptide is identical to the query sequence except that the subject polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the query amino acid sequence. In other  
10 words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a query amino acid sequence, up to 5% of the amino acid residues in the subject sequence may be inserted, deleted, (indels) or substituted with another amino acid. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those  
15 terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

As a practical matter, whether any particular polypeptide is at least 90%, 95%, 96%, 97%, 98% or 99% identical to, for instance, the amino acid sequences shown in SEQ ID NO:2 or to the amino acid sequence encoded by deposited DNA clone can be  
20 determined conventionally using known computer programs. A preferred method for determining the best overall match between a query sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci. (1990) 6:237-245). In a sequence  
25 alignment the query and subject sequences are either both nucleotide sequences or both amino acid sequences. The result of said global sequence alignment is in percent identity. Preferred parameters used in a FASTDB amino acid alignment are: Matrix=PAM 0, k-tuple=2, Mismatch Penalty=1, Joining Penalty=20, Randomization Group Length=0, Cutoff Score=1, Window Size=sequence length, Gap Penalty=5,  
30 Gap Size Penalty=0.05, Window Size=500 or the length of the subject amino acid sequence, whichever is shorter.

For example, a 90 amino acid residue subject sequence is aligned with a 100 residue query sequence to determine percent identity. The deletion occurs at the N-terminus of the subject sequence and therefore, the FASTDB alignment does not show a matching/alignment of the first 10 residues at the N-terminus. The 10 unpaired residues represent 10% of the sequence (number of residues at the N- and C-termini not matched/total number of residues in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 residues were perfectly matched the final percent identity would be 90%. In another example, a 90 residue subject sequence is compared with a 100 residue query sequence. This time the deletions are internal deletions so there are no residues at the N- or C-termini of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only residue positions outside the N- and C-terminal ends of the subject sequence, as displayed in the FASTDB alignment, which are not

For example, a 90 amino acid residue subject sequence is aligned with a 100 residue query sequence to determine percent identity. The deletion occurs at the N-terminus of the subject sequence and therefore, the FASTDB alignment does not show a matching/alignment of the first 10 residues at the N-terminus. The 10 unpaired residues represent 10% of the sequence (number of residues at the N- and C-termini not matched/total number of residues in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 residues were perfectly matched the final percent identity would be 90%. In another example, a 90 residue subject sequence is compared with a 100 residue query sequence. This time the deletions are internal deletions so there are no residues at the N- or C-termini of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only residue positions outside the N- and C-terminal ends of the subject sequence, as displayed in the FASTDB alignment, which are not

matched/aligned with the query sequence are manually corrected for. No other manual corrections are to be made for the purposes of the present invention.

The CRCGCL variants may contain alterations in the coding regions, non-coding regions, or both. Especially preferred are polynucleotide variants containing alterations which produce silent substitutions, additions, or deletions, but do not alter the properties or activities of the encoded polypeptide. Nucleotide variants produced by silent substitutions due to the degeneracy of the genetic code are preferred. Moreover, variants in which 5-10, 1-5, or 1-2 amino acids are substituted, deleted, or added in any combination are also preferred. CRCGCL polynucleotide variants can be produced for a variety of reasons, e.g., to optimize codon expression for a particular host (change codons in the human mRNA to those preferred by a bacterial host such as *E. coli*).

Naturally occurring CRCGCL variants are called "allelic variants," and refer to one of several alternate forms of a gene occupying a given locus on a chromosome of an organism. (Genes II, Lewin, B., ed., John Wiley & Sons, New York (1985).) These allelic variants can vary at either the polynucleotide and/or polypeptide level. Alternatively, non-naturally occurring variants may be produced by mutagenesis techniques or by direct synthesis.

Using known methods of protein engineering and recombinant DNA technology, variants may be generated to improve or alter the characteristics of the CRCGCL polypeptides. For instance, one or more amino acids can be deleted from the N-terminus or C-terminus of the secreted protein without substantial loss of biological function. The authors of Ron et al., *J. Biol. Chem.* 268: 2984-2988 (1993), reported variant KGF proteins having heparin binding activity even after deleting 3, 8, or 27 amino-terminal amino acid residues. Similarly, Interferon gamma exhibited up to ten times higher activity after deleting 8-10 amino acid residues from the carboxy terminus of this protein. (Dobeli et al., *J. Biotechnology* 7:199-216 (1988).)

Moreover, ample evidence demonstrates that variants often retain a biological activity similar to that of the naturally occurring protein. For example, Gayle and coworkers (*J. Biol. Chem.* 268:22105-22111 (1993)) conducted extensive mutational analysis of human cytokine IL-1a. They used random mutagenesis to generate over

3,500 individual IL-1a mutants that averaged 2.5 amino acid changes per variant over the entire length of the molecule. Multiple mutations were examined at every possible amino acid position. The investigators found that "[m]ost of the molecule could be altered with little effect on either [binding or biological activity]." (See, Abstract.) In fact, only 23 unique amino acid sequences, out of more than 3,500 nucleotide sequences examined, produced a protein that significantly differed in activity from wild-type.

Furthermore, even if deleting one or more amino acids from the N-terminus or C-terminus of a polypeptide results in modification or loss of one or more biological functions, other biological activities may still be retained. For example, the ability of a deletion variant to induce and/or to bind antibodies which recognize the secreted form will likely be retained when less than the majority of the residues of the secreted form are removed from the N-terminus or C-terminus. Whether a particular polypeptide lacking N- or C-terminal residues of a protein retains such immunogenic activities can readily be determined by routine methods described herein and otherwise known in the art.

Thus, the invention further includes CRCGCL polypeptide variants which show substantial biological activity. Such variants include deletions, insertions, inversions, repeats, and substitutions selected according to general rules known in the art so as have little effect on activity.

The present application is directed to nucleic acid molecules at least 90%, 95%, 96%, 97%, 98% or 99% identical to the nucleic acid sequences disclosed herein, (e.g., encoding a polypeptide having the amino acid sequence of an N and/or C terminal deletion disclosed below as m-n of SEQ ID NO:2), irrespective of whether they encode a polypeptide having CRCGCL functional activity. This is because even where a particular nucleic acid molecule does not encode a polypeptide having CRCGCL functional activity, one of skill in the art would still know how to use the nucleic acid molecule, for instance, as a hybridization probe or a polymerase chain reaction (PCR) primer. Uses of the nucleic acid molecules of the present invention that do not encode a polypeptide having CRCGCL functional activity include, inter alia, (1) isolating a CRCGCL gene or allelic or splice variants thereof in a cDNA

09376430-001595

library; (2) in situ hybridization (e.g., "FISH") to metaphase chromosomal spreads to provide precise chromosomal location of the CRCGCL gene, as described in Verma et al., Human Chromosomes: A Manual of Basic Techniques, Pergamon Press, New York (1988); and (3) Northern Blot analysis for detecting CRCGCL mRNA expression in specific tissues.

Preferred, however, are nucleic acid molecules having sequences at least 90%, 95%, 96%, 97%, 98% or 99% identical to the nucleic acid sequences disclosed herein, which do, in fact, encode a polypeptide having CRCGCL functional activity. By "a polypeptide having CRCGCL functional activity" is intended polypeptides exhibiting activity similar, but not necessarily identical, to a functional activity of the CRCGCL polypeptides of the present invention (e.g., complete (full-length) CRCGCL, mature CRCGCL and soluble CRCGCL (e.g., having sequences contained in the extracellular domain of CRCGCL) as measured, for example, in a particular immunoassay or biological assay. For example, a CRCGCL functional activity can routinely be measured by determining the ability of a CRCGCL polypeptide to bind a CRCGCL ligand. CRCGCL functional activity may also be measured by determining the ability of a polypeptide, such as cognate ligand which is free or expressed on a cell surface, to induce cells expressing the polypeptide.

Of course, due to the degeneracy of the genetic code, one of ordinary skill in the art will immediately recognize that a large number of the nucleic acid molecules having a sequence at least 90%, 95%, 96%, 97%, 98%, or 99% identical to the nucleic acid sequence of the deposited cDNA, the nucleic acid sequence shown in Figures 1A-1B (SEQ ID NO:1), or fragments thereof, will encode polypeptides "having CRCGCL functional activity." In fact, since degenerate variants of any of these nucleotide sequences all encode the same polypeptide, in many instances, this will be clear to the skilled artisan even without performing the above described comparison assay. It will be further recognized in the art that, for such nucleic acid molecules that are not degenerate variants, a reasonable number will also encode a polypeptide having CRCGCL functional activity. This is because the skilled artisan is fully aware of amino acid substitutions that are either less likely or not likely to significantly effect protein function (e.g., replacing one aliphatic amino acid with a second

aliphatic amino acid), as further described below.

For example, guidance concerning how to make phenotypically silent amino acid substitutions is provided in Bowie et al., "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions," *Science* 247:1306-1310 (1990),  
5 wherein the authors indicate that there are two main strategies for studying the tolerance of an amino acid sequence to change.

The first strategy exploits the tolerance of amino acid substitutions by natural selection during the process of evolution. By comparing amino acid sequences in different species, conserved amino acids can be identified. These conserved amino  
10 acids are likely important for protein function. In contrast, the amino acid positions where substitutions have been tolerated by natural selection indicates that these positions are not critical for protein function. Thus, positions tolerating amino acid substitution could be modified while still maintaining biological activity of the protein.

15 The second strategy uses genetic engineering to introduce amino acid changes at specific positions of a cloned gene to identify regions critical for protein function.

For example, site directed mutagenesis or alanine-scanning mutagenesis (introduction of single alanine mutations at every residue in the molecule) can be used. (Cunningham and Wells, *Science* 244:1081-1085 (1989).) The resulting mutant  
20 molecules can then be tested for biological activity.

As the authors state, these two strategies have revealed that proteins are surprisingly tolerant of amino acid substitutions. The authors further indicate which amino acid changes are likely to be permissive at certain amino acid positions in the protein. For example, most buried (within the tertiary structure of the protein) amino  
25 acid residues require nonpolar side chains, whereas few features of surface side chains are generally conserved. Moreover, tolerated conservative amino acid substitutions involve replacement of the aliphatic or hydrophobic amino acids Ala, Val, Leu and Ile; replacement of the hydroxyl residues Ser and Thr; replacement of the acidic residues Asp and Glu; replacement of the amide residues Asn and Gln, replacement of  
30 the basic residues Lys, Arg, and His; replacement of the aromatic residues Phe, Tyr, and Trp, and replacement of the small-sized amino acids Ala, Ser, Thr, Met, and Gly.



For example, site directed changes at the amino acid level of CRCGCL can be made by replacing a particular amino acid with a conservative amino acid. Preferred conservative mutations include: M1 replaced with A, G, I, L, S, T, or V; G2 replaced with A, I, L, S, T, M, or V; R3 replaced with H, or K; L4 replaced with A, G, I, S, T, M, or V; V5 replaced with A, G, I, L, S, T, or M; L6 replaced with A, G, I, S, T, M, or V; L7 replaced with A, G, I, S, T, M, or V; W8 replaced with F, or Y; G9 replaced with A, I, L, S, T, M, or V; A10 replaced with G, I, L, S, T, M, or V; A11 replaced with G, I, L, S, T, M, or V; V12 replaced with A, G, I, L, S, T, or M; F13 replaced with W, or Y; L14 replaced with A, G, I, S, T, M, or V; L15 replaced with A, G, I, S, T, M, or V; G16 replaced with A, I, L, S, T, M, or V; G17 replaced with A, I, L, S, T, M, or V; W18 replaced with F, or Y; M19 replaced with A, G, I, L, S, T, or V; A20 replaced with G, I, L, S, T, M, or V; L21 replaced with A, G, I, S, T, M, or V; G22 replaced with A, I, L, S, T, M, or V; Q23 replaced with N; G24 replaced with A, I, L, S, T, M, or V; G25 replaced with A, I, L, S, T, M, or V; A26 replaced with G, I, L, S, T, M, or V; A27 replaced with G, I, L, S, T, M, or V; E28 replaced with D; G29 replaced with A, I, L, S, T, M, or V; V30 replaced with A, G, I, L, S, T, or M; Q31 replaced with N; I32 replaced with A, G, L, S, T, M, or V; Q33 replaced with N; I34 replaced with A, G, L, S, T, M, or V; I35 replaced with A, G, L, S, T, M, or V; Y36 replaced with F, or W; F37 replaced with W, or Y; N38 replaced with Q; L39 replaced with A, G, I, S, T, M, or V; E40 replaced with D; T41 replaced with A, G, I, L, S, M, or V; V42 replaced with A, G, I, L, S, T, or M; Q43 replaced with N; V44 replaced with A, G, I, L, S, T, or M; T45 replaced with A, G, I, L, S, M, or V; W46 replaced with F, or Y; N47 replaced with Q; A48 replaced with G, I, L, S, T, M, or V; S49 replaced with A, G, I, L, T, M, or V; K50 replaced with H, or R; Y51 replaced with F, or W; S52 replaced with A, G, I, L, T, M, or V; R53 replaced with H, or K; T54 replaced with A, G, I, L, S, M, or V; N55 replaced with Q; L56 replaced with A, G, I, S, T, M, or V; T57 replaced with A, G, I, L, S, M, or V; F58 replaced with W, or Y; H59 replaced with K, or R; Y60 replaced with F, or W; R61 replaced with H, or K; F62 replaced with W, or Y; N63 replaced with Q; G64 replaced with A, I, L, S, T, M, or V; D65 replaced with E; E66 replaced with D; A67 replaced with G, I, L, S, T, M, or V; Y68 replaced with F, or W; D69 replaced with E; Q70 replaced with N; T72

replaced with A, G, I, L, S, M, or V; N73 replaced with Q; Y74 replaced with F, or W; L75 replaced with A, G, I, S, T, M, or V; L76 replaced with A, G, I, S, T, M, or V; Q77 replaced with N; E78 replaced with D; G79 replaced with A, I, L, S, T, M, or V; H80 replaced with K, or R; T81 replaced with A, G, I, L, S, M, or V; S82 replaced with A, G, I, L, T, M, or V; G83 replaced with A, I, L, S, T, M, or V; L85 replaced with A, G, I, S, T, M, or V; L86 replaced with A, G, I, S, T, M, or V; D87 replaced with E; A88 replaced with G, I, L, S, T, M, or V; E89 replaced with D; Q90 replaced with N; R91 replaced with H, or K; D92 replaced with E; D93 replaced with E; I94 replaced with A, G, L, S, T, M, or V; L95 replaced with A, G, I, S, T, M, or V; Y96 replaced with F, or W; F97 replaced with W, or Y; S98 replaced with A, G, I, L, T, M, or V; I99 replaced with A, G, L, S, T, M, or V; R100 replaced with H, or K; N101 replaced with Q; G102 replaced with A, I, L, S, T, M, or V; T103 replaced with A, G, I, L, S, M, or V; H104 replaced with K, or R; V106 replaced with A, G, I, L, S, T, or M; F107 replaced with W, or Y; T108 replaced with A, G, I, L, S, M, or V; A109 replaced with G, I, L, S, T, M, or V; S110 replaced with A, G, I, L, T, M, or V; R111 replaced with H, or K; W112 replaced with F, or Y; M113 replaced with A, G, I, L, S, T, or V; V114 replaced with A, G, I, L, S, T, or M; Y115 replaced with F, or W; Y116 replaced with F, or W; L117 replaced with A, G, I, S, T, M, or V; K118 replaced with H, or R; S120 replaced with A, G, I, L, T, M, or V; S121 replaced with A, G, I, L, T, M, or V; K123 replaced with H, or R; H124 replaced with K, or R; V125 replaced with A, G, I, L, S, T, or M; R126 replaced with H, or K; F127 replaced with W, or Y; S128 replaced with A, G, I, L, T, M, or V; W129 replaced with F, or Y; H130 replaced with K, or R; Q131 replaced with N; D132 replaced with E; A133 replaced with G, I, L, S, T, M, or V; V134 replaced with A, G, I, L, S, T, or M; T135 replaced with A, G, I, L, S, M, or V; V136 replaced with A, G, I, L, S, T, or M; T137 replaced with A, G, I, L, S, M, or V; S139 replaced with A, G, I, L, T, M, or V; D140 replaced with E; L141 replaced with A, G, I, S, T, M, or V; S142 replaced with A, G, I, L, T, M, or V; Y143 replaced with F, or W; G144 replaced with A, I, L, S, T, M, or V; D145 replaced with E; L146 replaced with A, G, I, S, T, M, or V; L147 replaced with A, G, I, S, T, M, or V; Y148 replaced with F, or W; E149 replaced with D; V150 replaced with A, G, I, L, S, T, or M; Q151 replaced with N; Y152 replaced with F, or





M, or V; E320 replaced with D; A321 replaced with G, I, L, S, T, M, or V; E322 replaced with D; S323 replaced with A, G, I, L, T, M, or V; R325 replaced with H, or K; M326 replaced with A, G, I, L, S, T, or V; L327 replaced with A, G, I, S, T, M, or V; D328 replaced with E; Q330 replaced with N; T331 replaced with A, G, I, L, S, M, or V; E332 replaced with D; E333 replaced with D; K334 replaced with H, or R; E335 replaced with D; A336 replaced with G, I, L, S, T, M, or V; S337 replaced with A, G, I, L, T, M, or V; G338 replaced with A, I, L, S, T, M, or V; G339 replaced with A, I, L, S, T, M, or V; S340 replaced with A, G, I, L, T, M, or V; L341 replaced with A, G, I, S, T, M, or V; Q342 replaced with N; L343 replaced with A, G, I, S, T, M, or V; H345 replaced with K, or R; Q346 replaced with N; L348 replaced with A, G, I, S, T, M, or V; Q349 replaced with N; G350 replaced with A, I, L, S, T, M, or V; G351 replaced with A, I, L, S, T, M, or V; D352 replaced with E; V353 replaced with A, G, I, L, S, T, or M; V354 replaced with A, G, I, L, S, T, or M; T355 replaced with A, G, I, L, S, M, or V; I356 replaced with A, G, L, S, T, M, or V; G357 replaced with A, I, L, S, T, M, or V; G358 replaced with A, I, L, S, T, M, or V; F359 replaced with W, or Y; T360 replaced with A, G, I, L, S, M, or V; F361 replaced with W, or Y; V362 replaced with A, G, I, L, S, T, or M; M363 replaced with A, G, I, L, S, T, or V; N364 replaced with Q; D365 replaced with E; R366 replaced with H, or K; S367 replaced with A, G, I, L, T, M, or V; Y368 replaced with F, or W; V369 replaced with A, G, I, L, S, T, or M; A370 replaced with G, I, L, S, T, M, or V; L371 replaced with A, G, I, S, T, M, or V.

The resulting constructs can be routinely screened for activities or functions described throughout the specification and known in the art. Preferably, the resulting constructs have an increased CRCGCL activity or function, while the remaining CRCGCL activities or functions are maintained. More preferably, the resulting constructs have more than one increased CRCGCL activity or function, while the remaining CRCGCL activities or functions are maintained.

Besides conservative amino acid substitution, variants of CRCGCL include (i) substitutions with one or more of the non-conserved amino acid residues, where the substituted amino acid residues may or may not be one encoded by the genetic code, or (ii) substitution with one or more of amino acid residues having a substituent

05376430-031399

5

10

20

25

30

replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; A26 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; A27 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; E28 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; G29 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; V30 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Q31 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; I32 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Q33 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; I34 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; I35 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Y36 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; F37 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; N38 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; L39 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; E40 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; T41 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; V42 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Q43 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; V44 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; T45 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; W46 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; N47 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; A48 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; S49 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; K50 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; Y51 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; S52 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; R53 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; T54 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; N55 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; L56 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; T57 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; F58 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; H59 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; Y60 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; R61 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; F62 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; N63 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; G64 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; D65 replaced with H, K, R, A, G,

I, L, S, T, M, V, N, Q, F, W, Y, P, or C; E66 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; A67 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Y68 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; D69 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; Q70 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; C71 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or P; T72 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; N73 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; Y74 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; L75 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L76 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Q77 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; E78 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; G79 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; H80 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; T81 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; S82 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; G83 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; C84 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or P; L85 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L86 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; D87 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; A88 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; E89 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; Q90 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; R91 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; D92 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; D93 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; I94 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L95 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Y96 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; F97 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; S98 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; I99 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; R100 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; N101 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; G102 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; T103 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; H104 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F,

668780-0E494E60



W, Y, P, or C; P105 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or C; V106 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; F107 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; T108 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; A109 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; S110 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; R111 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; W112 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; M113 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; V114 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Y115 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; Y116 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; L117 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; K118 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; P119 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or C; S120 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; S121 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; P122 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or C; K123 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; H124 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; V125 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; R126 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; F127 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; S128 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; W129 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; H130 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; Q131 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; D132 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; A133 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; V134 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; T135 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; V136 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; T137 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; C138 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or P; S139 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; D140 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; L141 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; S142 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Y143 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; G144

replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; D145 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; L146 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L147 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Y148 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; E149 replaced with

5 H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; V150 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Q151 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; Y152 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; R153 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; S154 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; P155 replaced with D, E, H, K,

10 R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or C; F156 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; D157 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; T158 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; E159 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; W160 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; Q161 replaced with

15 D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; S162 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; K163 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; Q164 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; E165 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; N166 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; T167

20 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; C168 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or P; N169 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; V170 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; T171 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; I172 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; E173 replaced with H, K, R, A, G, I, L, S,

25 T, M, V, N, Q, F, W, Y, P, or C; G174 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L175 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; D176 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; A177 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; E178 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; K179 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y,

30 P, or C; C180 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or P; Y181 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; S182



G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; T221 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; P222 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or C; T223 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; P224 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or C; P225 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or C; K226 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; P227 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or C; K228 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; L229 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; S230 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; K231 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; F232 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; I233 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L234 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; I235 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; S236 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; S237 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L238 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; A239 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; I240 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L241 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L242 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; M243 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; V244 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; S245 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L246 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L247 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L248 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L249 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; S250 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L251 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; W252 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; K253 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; L254 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; W255 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; R256 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; V257 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; K258 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; K259 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; F260 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; L261 replaced with

668780-02492660



replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; G301 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; A302 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; E303 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; Q304 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; E305 replaced  
5 with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; S306 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; G307 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; P308 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or C; E309 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; E310 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; P311 replaced  
10 with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or C; L312 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; V313 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; V314 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Q315 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; L316 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; A317 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; K318 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; T319 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; E320 replaced with H, K, R, A,  
G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; A321 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; E322 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; S323 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; P324 replaced with  
20 D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or C; R325 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; M326 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L327 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; D328 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; P329 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or C; Q330 replaced with  
25 D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; T331 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; E332 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; E333 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; K334 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; E335 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; A336 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; S337 replaced with D, E, H, K,  
30 R, N, Q, F, W, Y, P, or C; G338 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C;



constructs have more than one loss of CRCGCL activity or function, while the remaining CRCGCL activities or functions are maintained.

Additionally, more than one amino acid (e.g., 2, 3, 4, 5, 6, 7, 8, 9 and 10) can be replaced with the substituted amino acids as described above (either conservative or nonconservative). The substituted amino acids can occur in the full length, mature, or proprotein form of CRCGCL protein, as well as the N- and C- terminal deletion mutants, having the general formula m-n, listed below.

A further embodiment of the invention relates to a polypeptide which comprises the amino acid sequence of a CRCGCL polypeptide having an amino acid sequence which contains at least one amino acid substitution, but not more than 50 amino acid substitutions, even more preferably, not more than 40 amino acid substitutions, still more preferably, not more than 30 amino acid substitutions, and still even more preferably, not more than 20 amino acid substitutions. Of course, in order of ever-increasing preference, it is highly preferable for a peptide or polypeptide to have an amino acid sequence which comprises the amino acid sequence of a CRCGCL polypeptide, which contains at least one, but not more than 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1 amino acid substitutions. In specific embodiments, the number of additions, substitutions, and/or deletions in the amino acid sequence of Figures 1A-1B or fragments thereof (e.g., the mature form and/or other fragments described herein), is 1-5, 5-10, 5-25, 5-50, 10-50 or 50-150, conservative amino acid substitutions are preferable.

#### **Polynucleotide and Polypeptide Fragments**

The present invention is further directed to fragments of the isolated nucleic acid molecules described herein. By a fragment of an isolated nucleic acid molecule having, for example, the nucleotide sequence of the deposited cDNA (clone HTAEK53), a nucleotide sequence encoding the polypeptide sequence encoded by the deposited cDNA, a nucleotide sequence encoding the polypeptide sequence depicted in Figures 1A-1B (SEQ ID NO:2), the nucleotide sequence shown in Figures 1A-1B (SEQ ID NO:1), or the complementary strand thereto, is intended fragments at least 15 nt, and more preferably at least about 20 nt, still more preferably at least 30 nt,



and even more preferably, at least about 40, 50, 100, 150, 200, 250, 300, 325, 350, 375, 400, 450, 500, 550, or 600 nt in length. These fragments have numerous uses that include, but are not limited to, diagnostic probes and primers as discussed herein. Of course, larger fragments, such as those of 501-1500 nt in length are also useful according to the present invention as are fragments corresponding to most, if not all, of the nucleotide sequences of the deposited cDNA (clone HTAEK53) or as shown in Figures 1A-1B (SEQ ID NO:1). By a fragment at least 20 nt in length, for example, is intended fragments which include 20 or more contiguous bases from, for example, the nucleotide sequence of the deposited cDNA, or the nucleotide sequence as shown in Figures 1A-1B (SEQ ID NO:1).

Moreover, representative examples of CRCGCL polynucleotide fragments include, for example, fragments having a sequence from about nucleotide number 1-50, 51-100, 101-150, 151-200, 201-250, 251-300, 301-350, 351-400, 401-450, 451-500, 501-550, 551-600, 651-700, 701-750, 751-800, 800-850, 851-900, 901-950, 951-1000, 1001-1050, 1051-1100, 1101-1150, 1151-1200, 1201-1250, 1251-1300, 1301-1350, 1351-1400, 1401-1450, 1451-1500, 1501-1550, 1551-1600, 1601-1650, 1651-1700, 1701-1750, 1751-1800, 1801-1850, 1851-1900, 1901-1950, 1951-2000, and/or 2001 to the end of SEQ ID NO:1 or the complementary strand thereto, or the cDNA contained in the deposited clone. In this context "about" includes the particularly recited ranges, larger or smaller by several (5, 4, 3, 2, or 1) nucleotides, at either terminus or at both termini.

Preferably, the polynucleotide fragments of the invention encode a polypeptide which demonstrates a CRCGCL functional activity. By a polypeptide demonstrating a CRCGCL "functional activity" is meant, a polypeptide capable of displaying one or more known functional activities associated with a full-length (complete) CRCGCL protein. Such functional activities include, but are not limited to, biological activity, antigenicity [ability to bind (or compete with a CRCGCL polypeptide for binding) to an anti-CRCGCL antibody], immunogenicity (ability to generate antibody which binds to a CRCGCL polypeptide), ability to form multimers with CRCGCL polypeptides of the invention, and ability to bind to a receptor or ligand for a CRCGCL polypeptide.

The functional activity of CRCGCL polypeptides, and fragments, variants derivatives, and analogs thereof, can be assayed by various methods.

For example, in one embodiment where one is assaying for the ability to bind or compete with full-length CRCGCL polypeptide for binding to anti-CRCGCL antibody, various immunoassays known in the art can be used, including but not limited to, competitive and non-competitive assay systems using techniques such as radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoradiometric assays, gel diffusion precipitation reactions, immunodiffusion assays, in situ immunoassays (using colloidal gold, enzyme or radioisotope labels, for example), western blots, precipitation reactions, agglutination assays (e.g., gel agglutination assays, hemagglutination assays), complement fixation assays, immunofluorescence assays, protein A assays, and immunoelectrophoresis assays, etc. In one embodiment, antibody binding is detected by detecting a label on the primary antibody. In another embodiment, the primary antibody is detected by detecting binding of a secondary antibody or reagent to the primary antibody. In a further embodiment, the secondary antibody is labeled. Many means are known in the art for detecting binding in an immunoassay and are within the scope of the present invention.

In another embodiment, where a CRCGCL ligand is identified, or the ability of a polypeptide fragment, variant or derivative of the invention to multimerize is being evaluated, binding can be assayed, e.g., by means well-known in the art, such as, for example, reducing and non-reducing gel chromatography, protein affinity chromatography, and affinity blotting. See generally, Phizicky, E., et al., 1995, Microbiol. Rev. 59:94-123. In another embodiment, physiological correlates of CRCGCL binding to its substrates (signal transduction) can be assayed.

In addition, assays described herein (see Examples) and otherwise known in the art may routinely be applied to measure the ability of CRCGCL polypeptides and fragments, variants derivatives and analogs thereof to elicit CRCGCL related biological activity (either in vitro or in vivo). Other methods will be known to the skilled artisan and are within the scope of the invention.

The present invention is further directed to fragments of the CRCGCL polypeptide described herein. By a fragment of an isolated the CRCGCL polypeptide, for example, encoded by the deposited cDNA (clone HTAEK53), the polypeptide sequence encoded by the deposited cDNA, the polypeptide sequence depicted in Figures 1A-1B (SEQ ID NO:2), is intended to encompass polypeptide fragments contained in SEQ ID NO:2 or encoded by the cDNA contained in the deposited clone. Protein fragments may be "free-standing," or comprised within a larger polypeptide of which the fragment forms a part or region, most preferably as a single continuous region. Representative examples of polypeptide fragments of the invention, include, for example, fragments from about amino acid number 1-20, 21-40, 41-60, 61-80, 81-100, 102-120, 121-140, 141-160, 161-180, 181-200, 201-220, 221-240, 241-260, 261-280, or 281 to the end of the coding region. Moreover, polypeptide fragments can be at least 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, or 150 amino acids in length. In this context "about" includes the particularly recited ranges, larger or smaller by several (5, 4, 3, 2, or 1) amino acids, at either extreme or at both extremes.

Even if deletion of one or more amino acids from the N-terminus of a protein results in modification of loss of one or more biological functions of the protein, other functional activities (e.g., biological activities, ability to multimerize, ability to bind CRCGCL ligand) may still be retained. For example, the ability of shortened CRCGCL muteins to induce and/or bind to antibodies which recognize the complete or mature forms of the polypeptides generally will be retained when less than the majority of the residues of the complete or mature polypeptide are removed from the N-terminus. Whether a particular polypeptide lacking N-terminal residues of a complete polypeptide retains such immunologic activities can readily be determined by routine methods described herein and otherwise known in the art. It is not unlikely that an CRCGCL mutein with a large number of deleted N-terminal amino acid residues may retain some biological or immunogenic activities. In fact, peptides composed of as few as six CRCGCL amino acid residues may often evoke an immune response.

Accordingly, polypeptide fragments include the secreted CRCGCL protein as well as the mature form. Further preferred polypeptide fragments include the secreted CRCGCL protein or the mature form having a continuous series of deleted residues from the amino or the carboxy terminus, or both. For example, any number of amino acids, ranging from 1-60, can be deleted from the amino terminus of either the secreted CRCGCL polypeptide or the mature form. Similarly, any number of amino acids, ranging from 1-30, can be deleted from the carboxy terminus of the secreted CRCGCL protein or mature form. Furthermore, any combination of the above amino and carboxy terminus deletions are preferred. Similarly, polynucleotide fragments encoding these CRCGCL polypeptide fragments are also preferred.

Particularly, N-terminal deletions of the CRCGCL polypeptide can be described by the general formula m-371, where m is an integer from 2 to 370, where m corresponds to the position of the amino acid residue identified in SEQ ID NO:2. More in particular, the invention provides polynucleotides encoding polypeptides comprising, or alternatively consisting of, the amino acid sequence of residues of N-terminal deletions of the CRCGCL polypeptide of the invention shown as SEQ ID NO:2, including polypeptides comprising the amino acid sequence of residues: G-2 to L-371; R-3 to L-371; L-4 to L-371; V-5 to L-371; L-6 to L-371; L-7 to L-371; W-8 to L-371; G-9 to L-371; A-10 to L-371; A-11 to L-371; V-12 to L-371; F-13 to L-371; L-14 to L-371; L-15 to L-371; G-16 to L-371; G-17 to L-371; W-18 to L-371; M-19 to L-371; A-20 to L-371; L-21 to L-371; G-22 to L-371; Q-23 to L-371; G-24 to L-371; G-25 to L-371; A-26 to L-371; A-27 to L-371; E-28 to L-371; G-29 to L-371; V-30 to L-371; Q-31 to L-371; I-32 to L-371; Q-33 to L-371; I-34 to L-371; I-35 to L-371; Y-36 to L-371; F-37 to L-371; N-38 to L-371; L-39 to L-371; E-40 to L-371; T-41 to L-371; V-42 to L-371; Q-43 to L-371; V-44 to L-371; T-45 to L-371; W-46 to L-371; N-47 to L-371; A-48 to L-371; S-49 to L-371; K-50 to L-371; Y-51 to L-371; S-52 to L-371; R-53 to L-371; T-54 to L-371; N-55 to L-371; L-56 to L-371; T-57 to L-371; F-58 to L-371; H-59 to L-371; Y-60 to L-371; R-61 to L-371; F-62 to L-371; N-63 to L-371; G-64 to L-371; D-65 to L-371; E-66 to L-371; A-67 to L-371; Y-68 to L-371; D-69 to L-371; Q-70 to L-371; C-71 to L-371; T-72 to L-371; N-73 to L-371; Y-74 to L-371; L-75 to L-371; L-76 to L-371; Q-77 to L-371; E-78 to L-371; G-79 to

L-371; H-80 to L-371; T-81 to L-371; S-82 to L-371; G-83 to L-371; C-84 to L-371;  
 L-85 to L-371; L-86 to L-371; D-87 to L-371; A-88 to L-371; E-89 to L-371; Q-90 to  
 L-371; R-91 to L-371; D-92 to L-371; D-93 to L-371; I-94 to L-371; L-95 to L-371;  
 Y-96 to L-371; F-97 to L-371; S-98 to L-371; I-99 to L-371; R-100 to L-371; N-101  
 5 to L-371; G-102 to L-371; T-103 to L-371; H-104 to L-371; P-105 to L-371; V-106 to  
 L-371; F-107 to L-371; T-108 to L-371; A-109 to L-371; S-110 to L-371; R-111 to L-  
 371; W-112 to L-371; M-113 to L-371; V-114 to L-371; Y-115 to L-371; Y-116 to L-  
 371; L-117 to L-371; K-118 to L-371; P-119 to L-371; S-120 to L-371; S-121 to L-  
 371; P-122 to L-371; K-123 to L-371; H-124 to L-371; V-125 to L-371; R-126 to L-  
 10 371; F-127 to L-371; S-128 to L-371; W-129 to L-371; H-130 to L-371; Q-131 to L-  
 371; D-132 to L-371; A-133 to L-371; V-134 to L-371; T-135 to L-371; V-136 to L-  
 371; T-137 to L-371; C-138 to L-371; S-139 to L-371; D-140 to L-371; L-141 to L-  
 371; S-142 to L-371; Y-143 to L-371; G-144 to L-371; D-145 to L-371; L-146 to L-  
 371; L-147 to L-371; Y-148 to L-371; E-149 to L-371; V-150 to L-371; Q-151 to L-  
 15 371; Y-152 to L-371; R-153 to L-371; S-154 to L-371; P-155 to L-371; F-156 to L-  
 371; D-157 to L-371; T-158 to L-371; E-159 to L-371; W-160 to L-371; Q-161 to L-  
 371; S-162 to L-371; K-163 to L-371; Q-164 to L-371; E-165 to L-371; N-166 to L-  
 371; T-167 to L-371; C-168 to L-371; N-169 to L-371; V-170 to L-371; T-171 to L-  
 371; I-172 to L-371; E-173 to L-371; G-174 to L-371; L-175 to L-371; D-176 to L-  
 20 371; A-177 to L-371; E-178 to L-371; K-179 to L-371; C-180 to L-371; Y-181 to L-  
 371; S-182 to L-371; F-183 to L-371; W-184 to L-371; V-185 to L-371; R-186 to L-  
 371; V-187 to L-371; K-188 to L-371; A-189 to L-371; M-190 to L-371; E-191 to L-  
 371; D-192 to L-371; V-193 to L-371; Y-194 to L-371; G-195 to L-371; P-196 to L-  
 371; D-197 to L-371; T-198 to L-371; Y-199 to L-371; P-200 to L-371; S-201 to L-  
 25 371; D-202 to L-371; W-203 to L-371; S-204 to L-371; E-205 to L-371; V-206 to L-  
 371; T-207 to L-371; C-208 to L-371; W-209 to L-371; Q-210 to L-371; R-211 to L-  
 371; G-212 to L-371; E-213 to L-371; I-214 to L-371; R-215 to L-371; D-216 to L-  
 371; A-217 to L-371; C-218 to L-371; A-219 to L-371; E-220 to L-371; T-221 to L-  
 371; P-222 to L-371; T-223 to L-371; P-224 to L-371; P-225 to L-371; K-226 to L-  
 30 371; P-227 to L-371; K-228 to L-371; L-229 to L-371; S-230 to L-371; K-231 to L-  
 371; F-232 to L-371; I-233 to L-371; L-234 to L-371; I-235 to L-371; S-236 to L-371;

668780" 06492660

S-237 to L-371; L-238 to L-371; A-239 to L-371; I-240 to L-371; L-241 to L-371; L-242 to L-371; M-243 to L-371; V-244 to L-371; S-245 to L-371; L-246 to L-371; L-247 to L-371; L-248 to L-371; L-249 to L-371; S-250 to L-371; L-251 to L-371; W-252 to L-371; K-253 to L-371; L-254 to L-371; W-255 to L-371; R-256 to L-371; V-257 to L-371; K-258 to L-371; K-259 to L-371; F-260 to L-371; L-261 to L-371; I-262 to L-371; P-263 to L-371; S-264 to L-371; V-265 to L-371; P-266 to L-371; D-267 to L-371; P-268 to L-371; K-269 to L-371; S-270 to L-371; I-271 to L-371; F-272 to L-371; P-273 to L-371; G-274 to L-371; L-275 to L-371; F-276 to L-371; E-277 to L-371; I-278 to L-371; H-279 to L-371; Q-280 to L-371; G-281 to L-371; N-282 to L-371; F-283 to L-371; Q-284 to L-371; E-285 to L-371; W-286 to L-371; I-287 to L-371; T-288 to L-371; D-289 to L-371; T-290 to L-371; Q-291 to L-371; N-292 to L-371; V-293 to L-371; A-294 to L-371; H-295 to L-371; L-296 to L-371; H-297 to L-371; K-298 to L-371; M-299 to L-371; A-300 to L-371; G-301 to L-371; A-302 to L-371; E-303 to L-371; Q-304 to L-371; E-305 to L-371; S-306 to L-371; G-307 to L-371; P-308 to L-371; E-309 to L-371; E-310 to L-371; P-311 to L-371; L-312 to L-371; V-313 to L-371; V-314 to L-371; Q-315 to L-371; L-316 to L-371; A-317 to L-371; K-318 to L-371; T-319 to L-371; E-320 to L-371; A-321 to L-371; E-322 to L-371; S-323 to L-371; P-324 to L-371; R-325 to L-371; M-326 to L-371; L-327 to L-371; D-328 to L-371; P-329 to L-371; Q-330 to L-371; T-331 to L-371; E-332 to L-371; E-333 to L-371; K-334 to L-371; E-335 to L-371; A-336 to L-371; S-337 to L-371; G-338 to L-371; G-339 to L-371; S-340 to L-371; L-341 to L-371; Q-342 to L-371; L-343 to L-371; P-344 to L-371; H-345 to L-371; Q-346 to L-371; P-347 to L-371; L-348 to L-371; Q-349 to L-371; G-350 to L-371; G-351 to L-371; D-352 to L-371; V-353 to L-371; V-354 to L-371; T-355 to L-371; I-356 to L-371; G-357 to L-371; G-358 to L-371; F-359 to L-371; T-360 to L-371; F-361 to L-371; V-362 to L-371; M-363 to L-371; N-364 to L-371; D-365 to L-371; R-366 to L-371; of SEQ ID NO:2. Polynucleotides encoding these polypeptides are also encompassed by the invention.

Also as mentioned above, even if deletion of one or more amino acids from the C-terminus of a protein results in modification of loss of one or more biological functions of the protein, other functional activities (e.g., biological activities, ability to

multimerize, ability to bind CRCGCL ligand) may still be retained. For example the ability of the shortened CRCGCL mutein to induce and/or bind to antibodies which recognize the complete or mature forms of the polypeptide generally will be retained when less than the majority of the residues of the complete or mature polypeptide are removed from the C-terminus. Whether a particular polypeptide lacking C-terminal residues of a complete polypeptide retains such immunologic activities can readily be determined by routine methods described herein and otherwise known in the art. It is not unlikely that an CRCGCL mutein with a large number of deleted C-terminal amino acid residues may retain some biological or immunogenic activities. In fact, peptides composed of as few as six CRCGCL amino acid residues may often evoke an immune response.

Accordingly, the present invention further provides polypeptides having one or more residues deleted from the carboxy terminus of the amino acid sequence of the CRCGCL polypeptide shown in Figures 1A-1B (SEQ ID NO:2), as described by the general formula 1-n, where n is an integer from 2 to 371, where n corresponds to the position of amino acid residue identified in SEQ ID NO:2. More in particular, the invention provides polynucleotides encoding polypeptides comprising, or alternatively consisting of, the amino acid sequence of residues of C-terminal deletions of the CRCGCL polypeptide of the invention shown as SEQ ID NO:2 include polypeptides comprising the amino acid sequence of residues: M-1 to A-370; M-1 to V-369; M-1 to Y-368; M-1 to S-367; M-1 to R-366; M-1 to D-365; M-1 to N-364; M-1 to M-363; M-1 to V-362; M-1 to F-361; M-1 to T-360; M-1 to F-359; M-1 to G-358; M-1 to G-357; M-1 to I-356; M-1 to T-355; M-1 to V-354; M-1 to V-353; M-1 to D-352; M-1 to G-351; M-1 to G-350; M-1 to Q-349; M-1 to L-348; M-1 to P-347; M-1 to Q-346; M-1 to H-345; M-1 to P-344; M-1 to L-343; M-1 to Q-342; M-1 to L-341; M-1 to S-340; M-1 to G-339; M-1 to G-338; M-1 to S-337; M-1 to A-336; M-1 to E-335; M-1 to K-334; M-1 to E-333; M-1 to E-332; M-1 to T-331; M-1 to Q-330; M-1 to P-329; M-1 to D-328; M-1 to L-327; M-1 to M-326; M-1 to R-325; M-1 to P-324; M-1 to S-323; M-1 to E-322; M-1 to A-321; M-1 to E-320; M-1 to T-319; M-1 to K-318; M-1 to A-317; M-1 to L-316; M-1 to Q-315; M-1 to V-314; M-1 to V-313; M-1 to L-312; M-1 to P-311; M-1 to E-310; M-1 to E-309; M-1 to P-308;

M-1 to G-307; M-1 to S-306; M-1 to E-305; M-1 to Q-304; M-1 to E-303; M-1 to A-302; M-1 to G-301; M-1 to A-300; M-1 to M-299; M-1 to K-298; M-1 to H-297; M-1 to L-296; M-1 to H-295; M-1 to A-294; M-1 to V-293; M-1 to N-292; M-1 to Q-291; M-1 to T-290; M-1 to D-289; M-1 to T-288; M-1 to I-287; M-1 to W-286; M-1  
 5 'to E-285; M-1 to Q-284; M-1 to F-283; M-1 to N-282; M-1 to G-281; M-1 to Q-280; M-1 to H-279; M-1 to I-278; M-1 to E-277; M-1 to F-276; M-1 to L-275; M-1 to G-274; M-1 to P-273; M-1 to F-272; M-1 to I-271; M-1 to S-270; M-1 to K-269; M-1 to P-268; M-1 to D-267; M-1 to P-266; M-1 to V-265; M-1 to S-264; M-1 to P-263; M-1 to I-262; M-1 to L-261; M-1 to F-260; M-1 to K-259; M-1 to K-258; M-1 to V-257; M-1 to R-256; M-1 to W-255; M-1 to L-254; M-1 to K-253; M-1 to W-252; M-1 to L-251; M-1 to S-250; M-1 to L-249; M-1 to L-248; M-1 to L-247; M-1 to L-246; M-1 to S-245; M-1 to V-244; M-1 to M-243; M-1 to L-242; M-1 to L-241; M-1 to I-240; M-1 to A-239; M-1 to L-238; M-1 to S-237; M-1 to S-236; M-1 to I-235; M-1 to L-234; M-1 to I-233; M-1 to F-232; M-1 to K-231; M-1 to S-230; M-1 to L-229; M-1 to K-228; M-1 to P-227; M-1 to K-226; M-1 to P-225; M-1 to P-224; M-1  
 15 to T-223; M-1 to P-222; M-1 to T-221; M-1 to E-220; M-1 to A-219; M-1 to C-218; M-1 to A-217; M-1 to D-216; M-1 to R-215; M-1 to I-214; M-1 to E-213; M-1 to G-212; M-1 to R-211; M-1 to Q-210; M-1 to W-209; M-1 to C-208; M-1 to T-207; M-1 to V-206; M-1 to E-205; M-1 to S-204; M-1 to W-203; M-1 to D-202; M-1 to S-201; M-1 to P-200; M-1 to Y-199; M-1 to T-198; M-1 to D-197; M-1 to P-196; M-1 to G-195; M-1 to Y-194; M-1 to V-193; M-1 to D-192; M-1 to E-191; M-1 to M-190; M-1 to A-189; M-1 to K-188; M-1 to V-187; M-1 to R-186; M-1 to V-185; M-1 to W-184; M-1 to F-183; M-1 to S-182; M-1 to Y-181; M-1 to C-180; M-1 to K-179; M-1 to E-178; M-1 to A-177; M-1 to D-176; M-1 to L-175; M-1 to G-174; M-1 to E-173; M-1 to I-172; M-1 to T-171; M-1 to V-170; M-1 to N-169; M-1 to C-168; M-1 to T-167; M-1 to N-166; M-1 to E-165; M-1 to Q-164; M-1 to K-163; M-1 to S-162; M-1 to Q-161; M-1 to W-160; M-1 to E-159; M-1 to T-158; M-1 to D-157; M-1 to F-156; M-1 to P-155; M-1 to S-154; M-1 to R-153; M-1 to Y-152; M-1 to Q-151; M-1 to V-150; M-1 to E-149; M-1 to Y-148; M-1 to L-147; M-1 to L-146; M-1  
 25 to D-145; M-1 to G-144; M-1 to Y-143; M-1 to S-142; M-1 to L-141; M-1 to D-140; M-1 to S-139; M-1 to C-138; M-1 to T-137; M-1 to V-136; M-1 to T-135; M-1 to V-

668780-0619260



134; M-1 to A-133; M-1 to D-132; M-1 to Q-131; M-1 to H-130; M-1 to W-129; M-1 to S-128; M-1 to F-127; M-1 to R-126; M-1 to V-125; M-1 to H-124; M-1 to K-123; M-1 to P-122; M-1 to S-121; M-1 to S-120; M-1 to P-119; M-1 to K-118; M-1 to L-117; M-1 to Y-116; M-1 to Y-115; M-1 to V-114; M-1 to M-113; M-1 to W-112; M-1 to R-111; M-1 to S-110; M-1 to A-109; M-1 to T-108; M-1 to F-107; M-1 to V-106; M-1 to P-105; M-1 to H-104; M-1 to T-103; M-1 to G-102; M-1 to N-101; M-1 to R-100; M-1 to I-99; M-1 to S-98; M-1 to F-97; M-1 to Y-96; M-1 to L-95; M-1 to I-94; M-1 to D-93; M-1 to D-92; M-1 to R-91; M-1 to Q-90; M-1 to E-89; M-1 to A-88; M-1 to D-87; M-1 to L-86; M-1 to L-85; M-1 to C-84; M-1 to G-83; M-1 to S-82; M-1 to T-81; M-1 to H-80; M-1 to G-79; M-1 to E-78; M-1 to Q-77; M-1 to L-76; M-1 to L-75; M-1 to Y-74; M-1 to N-73; M-1 to T-72; M-1 to C-71; M-1 to Q-70; M-1 to D-69; M-1 to Y-68; M-1 to A-67; M-1 to E-66; M-1 to D-65; M-1 to G-64; M-1 to N-63; M-1 to F-62; M-1 to R-61; M-1 to Y-60; M-1 to H-59; M-1 to F-58; M-1 to T-57; M-1 to L-56; M-1 to N-55; M-1 to T-54; M-1 to R-53; M-1 to S-52; M-1 to Y-51; M-1 to K-50; M-1 to S-49; M-1 to A-48; M-1 to N-47; M-1 to W-46; M-1 to T-45; M-1 to V-44; M-1 to Q-43; M-1 to V-42; M-1 to T-41; M-1 to E-40; M-1 to L-39; M-1 to N-38; M-1 to F-37; M-1 to Y-36; M-1 to I-35; M-1 to I-34; M-1 to Q-33; M-1 to I-32; M-1 to Q-31; M-1 to V-30; M-1 to G-29; M-1 to E-28; M-1 to A-27; M-1 to A-26; M-1 to G-25; M-1 to G-24; M-1 to Q-23; M-1 to G-22; M-1 to L-21; M-1 to A-20; M-1 to M-19; M-1 to W-18; M-1 to G-17; M-1 to G-16; M-1 to L-15; M-1 to L-14; M-1 to F-13; M-1 to V-12; M-1 to A-11; M-1 to A-10; M-1 to G-9; M-1 to W-8; M-1 to L-7; of SEQ ID NO:2. Polynucleotides encoding these polypeptides are also encompassed by the invention.

In addition, any of the above listed N- or C-terminal deletions can be combined to produce a N- and C-terminal deleted CRCGCL polypeptide. The invention also provides polypeptides having one or more amino acids deleted from both the amino and the carboxyl termini, which may be described generally as having residues m-n of SEQ ID NO:2, where n and m are integers as described above. Polynucleotides encoding these polypeptides are also encompassed by the invention.

For example, N- and C-terminal deletion mutants comprising the soluble domain of SEQ ID NO:2 are also contemplated. In particular, the invention provides polynucleotides encoding polypeptides comprising, or alternatively consisting of, the amino acid sequence of residues of N-terminal deletions of the CRCGCL polypeptide of the invention shown as SEQ ID NO:2, including polypeptides comprising the amino acid sequence of residues: G-2 to K-231; R-3 to K-231; L-4 to K-231; V-5 to K-231; L-6 to K-231; L-7 to K-231; W-8 to K-231; G-9 to K-231; A-10 to K-231; A-11 to K-231; V-12 to K-231; F-13 to K-231; L-14 to K-231; L-15 to K-231; G-16 to K-231; G-17 to K-231; W-18 to K-231; M-19 to K-231; A-20 to K-231; L-21 to K-231; G-22 to K-231; Q-23 to K-231; G-24 to K-231; G-25 to K-231; A-26 to K-231; A-27 to K-231; E-28 to K-231; G-29 to K-231; V-30 to K-231; Q-31 to K-231; I-32 to K-231; Q-33 to K-231; I-34 to K-231; I-35 to K-231; Y-36 to K-231; F-37 to K-231; N-38 to K-231; L-39 to K-231; E-40 to K-231; T-41 to K-231; V-42 to K-231; Q-43 to K-231; V-44 to K-231; T-45 to K-231; W-46 to K-231; N-47 to K-231; A-48 to K-231; S-49 to K-231; K-50 to K-231; Y-51 to K-231; S-52 to K-231; R-53 to K-231; T-54 to K-231; N-55 to K-231; L-56 to K-231; T-57 to K-231; F-58 to K-231; H-59 to K-231; Y-60 to K-231; R-61 to K-231; F-62 to K-231; N-63 to K-231; G-64 to K-231; D-65 to K-231; E-66 to K-231; A-67 to K-231; Y-68 to K-231; D-69 to K-231; Q-70 to K-231; C-71 to K-231; T-72 to K-231; N-73 to K-231; Y-74 to K-231; L-75 to K-231; L-76 to K-231; Q-77 to K-231; E-78 to K-231; G-79 to K-231; H-80 to K-231; T-81 to K-231; S-82 to K-231; G-83 to K-231; C-84 to K-231; L-85 to K-231; L-86 to K-231; D-87 to K-231; A-88 to K-231; E-89 to K-231; Q-90 to K-231; R-91 to K-231; D-92 to K-231; D-93 to K-231; I-94 to K-231; L-95 to K-231; Y-96 to K-231; F-97 to K-231; S-98 to K-231; I-99 to K-231; R-100 to K-231; N-101 to K-231; G-102 to K-231; T-103 to K-231; H-104 to K-231; P-105 to K-231; V-106 to K-231; F-107 to K-231; T-108 to K-231; A-109 to K-231; S-110 to K-231; R-111 to K-231; W-112 to K-231; M-113 to K-231; V-114 to K-231; Y-115 to K-231; Y-116 to K-231; L-117 to K-231; K-118 to K-231; P-119 to K-231; S-120 to K-231; S-121 to K-231; P-122 to K-231; K-123 to K-231; H-124 to K-231; V-125 to K-231; R-126 to K-231; F-127 to K-231; S-128 to K-231; W-129 to K-231; H-130 to K-231; Q-131 to K-231; D-132 to K-231; A-133 to K-231; V-134 to

K-231; T-135 to K-231; V-136 to K-231; T-137 to K-231; C-138 to K-231; S-139 to K-231; D-140 to K-231; L-141 to K-231; S-142 to K-231; Y-143 to K-231; G-144 to K-231; D-145 to K-231; L-146 to K-231; L-147 to K-231; Y-148 to K-231; E-149 to K-231; V-150 to K-231; Q-151 to K-231; Y-152 to K-231; R-153 to K-231; S-154 to K-231; P-155 to K-231; F-156 to K-231; D-157 to K-231; T-158 to K-231; E-159 to K-231; W-160 to K-231; Q-161 to K-231; S-162 to K-231; K-163 to K-231; Q-164 to K-231; E-165 to K-231; N-166 to K-231; T-167 to K-231; C-168 to K-231; N-169 to K-231; V-170 to K-231; T-171 to K-231; I-172 to K-231; E-173 to K-231; G-174 to K-231; L-175 to K-231; D-176 to K-231; A-177 to K-231; E-178 to K-231; K-179 to K-231; C-180 to K-231; Y-181 to K-231; S-182 to K-231; F-183 to K-231; W-184 to K-231; V-185 to K-231; R-186 to K-231; V-187 to K-231; K-188 to K-231; A-189 to K-231; M-190 to K-231; E-191 to K-231; D-192 to K-231; V-193 to K-231; Y-194 to K-231; G-195 to K-231; P-196 to K-231; D-197 to K-231; T-198 to K-231; Y-199 to K-231; P-200 to K-231; S-201 to K-231; D-202 to K-231; W-203 to K-231; S-204 to K-231; E-205 to K-231; V-206 to K-231; T-207 to K-231; C-208 to K-231; W-209 to K-231; Q-210 to K-231; R-211 to K-231; G-212 to K-231; E-213 to K-231; I-214 to K-231; R-215 to K-231; D-216 to K-231; A-217 to K-231; C-218 to K-231; A-219 to K-231; E-220 to K-231; T-221 to K-231; P-222 to K-231; T-223 to K-231; P-224 to K-231; P-225 to K-231; K-226 to K-231; of SEQ ID NO:2.

Polynucleotides encoding these polypeptides are also encompassed by the invention.

Additionally, the invention provides polynucleotides encoding polypeptides comprising, or alternatively consisting of, the amino acid sequence of residues of C-terminal deletions of the CRCGCL polypeptide of the invention shown as SEQ ID NO:2 include polypeptides comprising the amino acid sequence of residues: Q-23 to K-231; Q-23 to S-230; Q-23 to L-229; Q-23 to K-228; Q-23 to P-227; Q-23 to K-226; Q-23 to P-225; Q-23 to P-224; Q-23 to T-223; Q-23 to P-222; Q-23 to T-221; Q-23 to E-220; Q-23 to A-219; Q-23 to C-218; Q-23 to A-217; Q-23 to D-216; Q-23 to R-215; Q-23 to I-214; Q-23 to E-213; Q-23 to G-212; Q-23 to R-211; Q-23 to Q-210; Q-23 to W-209; Q-23 to C-208; Q-23 to T-207; Q-23 to V-206; Q-23 to E-205; Q-23 to S-204; Q-23 to W-203; Q-23 to D-202; Q-23 to S-201; Q-23 to P-200; Q-23 to Y-199; Q-23 to T-198; Q-23 to D-197; Q-23 to P-196; Q-23 to G-195; Q-23 to Y-194;

Q-23 to V-193; Q-23 to D-192; Q-23 to E-191; Q-23 to M-190; Q-23 to A-189; Q-23 to K-188; Q-23 to V-187; Q-23 to R-186; Q-23 to V-185; Q-23 to W-184; Q-23 to F-183; Q-23 to S-182; Q-23 to Y-181; Q-23 to C-180; Q-23 to K-179; Q-23 to E-178; Q-23 to A-177; Q-23 to D-176; Q-23 to L-175; Q-23 to G-174; Q-23 to E-173; Q-23 to I-172; Q-23 to T-171; Q-23 to V-170; Q-23 to N-169; Q-23 to C-168; Q-23 to T-167; Q-23 to N-166; Q-23 to E-165; Q-23 to Q-164; Q-23 to K-163; Q-23 to S-162; Q-23 to Q-161; Q-23 to W-160; Q-23 to E-159; Q-23 to T-158; Q-23 to D-157; Q-23 to F-156; Q-23 to P-155; Q-23 to S-154; Q-23 to R-153; Q-23 to Y-152; Q-23 to Q-151; Q-23 to V-150; Q-23 to E-149; Q-23 to Y-148; Q-23 to L-147; Q-23 to L-146; Q-23 to D-145; Q-23 to G-144; Q-23 to Y-143; Q-23 to S-142; Q-23 to L-141; Q-23 to D-140; Q-23 to S-139; Q-23 to C-138; Q-23 to T-137; Q-23 to V-136; Q-23 to T-135; Q-23 to V-134; Q-23 to A-133; Q-23 to D-132; Q-23 to Q-131; Q-23 to H-130; Q-23 to W-129; Q-23 to S-128; Q-23 to F-127; Q-23 to R-126; Q-23 to V-125; Q-23 to H-124; Q-23 to K-123; Q-23 to P-122; Q-23 to S-121; Q-23 to S-120; Q-23 to P-119; Q-23 to K-118; Q-23 to L-117; Q-23 to Y-116; Q-23 to Y-115; Q-23 to V-114; Q-23 to M-113; Q-23 to W-112; Q-23 to R-111; Q-23 to S-110; Q-23 to A-109; Q-23 to T-108; Q-23 to F-107; Q-23 to V-106; Q-23 to P-105; Q-23 to H-104; Q-23 to T-103; Q-23 to G-102; Q-23 to N-101; Q-23 to R-100; Q-23 to I-99; Q-23 to S-98; Q-23 to F-97; Q-23 to Y-96; Q-23 to L-95; Q-23 to I-94; Q-23 to D-93; Q-23 to D-92; Q-23 to R-91; Q-23 to Q-90; Q-23 to E-89; Q-23 to A-88; Q-23 to D-87; Q-23 to L-86; Q-23 to L-85; Q-23 to C-84; Q-23 to G-83; Q-23 to S-82; Q-23 to T-81; Q-23 to H-80; Q-23 to G-79; Q-23 to E-78; Q-23 to Q-77; Q-23 to L-76; Q-23 to L-75; Q-23 to Y-74; Q-23 to N-73; Q-23 to T-72; Q-23 to C-71; Q-23 to Q-70; Q-23 to D-69; Q-23 to Y-68; Q-23 to A-67; Q-23 to E-66; Q-23 to D-65; Q-23 to G-64; Q-23 to N-63; Q-23 to F-62; Q-23 to R-61; Q-23 to Y-60; Q-23 to H-59; Q-23 to F-58; Q-23 to T-57; Q-23 to L-56; Q-23 to N-55; Q-23 to T-54; Q-23 to R-53; Q-23 to S-52; Q-23 to Y-51; Q-23 to K-50; Q-23 to S-49; Q-23 to A-48; Q-23 to N-47; Q-23 to W-46; Q-23 to T-45; Q-23 to V-44; Q-23 to Q-43; Q-23 to V-42; Q-23 to T-41; Q-23 to E-40; Q-23 to L-39; Q-23 to N-38; Q-23 to F-37; Q-23 to Y-36; Q-23 to I-35; Q-23 to I-34; Q-23 to Q-33; Q-23 to I-32; Q-23 to Q-31; Q-23 to V-30; Q-23 to G-29; of SEQ ID NO:2. Polynucleotides encoding these polypeptides are also encompassed by the invention.

052430-02192650

Moreover, a signal sequence may be added to these C-terminal constructs. For example, amino acids 1-22 of SEQ ID NO:2, amino acids 2-22 of SEQ ID NO:2, amino acids 3-22 of SEQ ID NO:2, amino acids 4-22 of SEQ ID NO:2, amino acids 5-22 of SEQ ID NO:2, amino acids 6-22 of SEQ ID NO:2, amino acids 7-22 of SEQ ID NO:2, amino acids 8-22 of SEQ ID NO:2, amino acids 9-22 of SEQ ID NO:2, amino acids 10-22 of SEQ ID NO:2, amino acids 11-22 of SEQ ID NO:2, amino acids 12-22 of SEQ ID NO:2, amino acids 13-22 of SEQ ID NO:2, amino acids 14-22 of SEQ ID NO:2, amino acids 15-22 of SEQ ID NO:2, amino acids 16-22 of SEQ ID NO:2, amino acids 17-22 of SEQ ID NO:2, amino acids 18-22 of SEQ ID NO:2, amino acids 19-22 of SEQ ID NO:2, amino acids 20-22 of SEQ ID NO:2, or amino acids 21-22 of SEQ ID NO:2 can be added to the N-terminus of each C-terminal constructs listed above.

Also included are a nucleotide sequence encoding a polypeptide consisting of a portion of the complete CRCGCL amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 209641 or 209691, where this portion excludes any integer of amino acid residues from 1 to about 361 amino acids from the amino terminus of the complete amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 209641 or 209691, or any integer of amino acid residues from 1 to about 361 amino acids from the carboxy terminus, or any combination of the above amino terminal and carboxy terminal deletions, of the complete amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 209641 or 209691. Polynucleotides encoding all of the above deletion mutant polypeptide forms also are provided.

The present application is also directed to proteins containing polypeptides at least 90%, 95%, 96%, 97%, 98% or 99% identical to the CRCGCL polypeptide sequence set forth herein m-n. In preferred embodiments, the application is directed to proteins containing polypeptides at least 90%, 95%, 96%, 97%, 98% or 99% identical to polypeptides having the amino acid sequence of the specific CRCGCL N- and C-terminal deletions recited herein. Polynucleotides encoding these polypeptides are also encompassed by the invention.

0976430-03139

Among the especially preferred fragments of the invention are fragments characterized by structural or functional attributes of CRCGCL. Such fragments include amino acid residues that comprise alpha-helix and alpha-helix forming regions ("alpha-regions"), beta-sheet and beta-sheet-forming regions ("beta-regions"), turn and turn-forming regions ("turn-regions"), coil and coil-forming regions ("coil-regions"), hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, surface forming regions, and high antigenic index regions (i.e., containing four or more contiguous amino acids having an antigenic index of greater than or equal to 1.5, as identified using the default parameters of the Jameson-Wolf program) of complete (i.e., full-length) CRCGCL (SEQ ID NO:2). Certain preferred regions are those set out in Figure 3 and include, but are not limited to, regions of the aforementioned types identified by analysis of the amino acid sequence depicted in Figures 1A-1B (SEQ ID NO:2), such preferred regions include; Garnier-Robson predicted alpha-regions, beta-regions, turn-regions, and coil-regions; Chou-Fasman predicted alpha-regions, beta-regions, turn-regions, and coil-regions; Kyte-Doolittle predicted hydrophilic and hydrophobic regions; Eisenberg alpha and beta amphipathic regions; Emini surface-forming regions; and Jameson-Wolf high antigenic index regions, as predicted using the default parameters of these computer programs. Polynucleotides encoding these polypeptides are also encompassed by the invention.

In additional embodiments, the polynucleotides of the invention encode functional attributes of CRCGCL. Preferred embodiments of the invention in this regard include fragments that comprise alpha-helix and alpha-helix forming regions ("alpha-regions"), beta-sheet and beta-sheet forming regions ("beta-regions"), turn and turn-forming regions ("turn-regions"), coil and coil-forming regions ("coil-regions"), hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions and high antigenic index regions of CRCGCL.

The data representing the structural or functional attributes of CRCGCL set forth in Figures 1A-1B and/or Table 1, as described above, was generated using the various modules and algorithms of the DNA\*STAR set on default parameters. In a preferred embodiment, the data presented in columns VIII, IX, XIII, and XIV of

Table 1 can be used to determine regions of CRCGCL which exhibit a high degree of potential for antigenicity. Regions of high antigenicity are determined from the data presented in columns VIII, IX, XIII, and/or IV by choosing values which represent regions of the polypeptide which are likely to be exposed on the surface of the polypeptide in an environment in which antigen recognition may occur in the process of initiation of an immune response.

Certain preferred regions in these regards are set out in Figure 3, but may, as shown in Table 1, be represented or identified by using tabular representations of the data presented in Figure 3. The DNA\*STAR computer algorithm used to generate Figure 3 (set on the original default parameters) was used to present the data in Figure 3 in a tabular format (See Table 1). The tabular format of the data in Figure 3 may be used to easily determine specific boundaries of a preferred region.

The above-mentioned preferred regions set out in Figure 3 and in Table 1 include, but are not limited to, regions of the aforementioned types identified by analysis of the amino acid sequence set out in Figures 1A-1B. As set out in Figure 3 and in Table 1, such preferred regions include Garnier-Robson alpha-regions, beta-regions, turn-regions, and coil-regions, Chou-Fasman alpha-regions, beta-regions, and coil-regions, Kyte-Doolittle hydrophilic regions and hydrophobic regions, Eisenberg alpha- and beta-amphipathic regions, Karplus-Schulz flexible regions, Emini surface-forming regions and Jameson-Wolf regions of high antigenic index.

Table 1

|    | Res | Position | I | II | III | IV | V | VI | VII | VIII  | IX    | X | XI | XII | XIII  | XIV  |
|----|-----|----------|---|----|-----|----|---|----|-----|-------|-------|---|----|-----|-------|------|
| 5  | Met | 1        | . | .  | B   | .  | . | .  | .   | -0.56 | 0.01  | * | *  | .   | -0.10 | 0.59 |
|    | Gly | 2        | . | .  | B   | .  | . | .  | .   | -0.98 | 0.23  | * | .  | .   | -0.10 | 0.34 |
|    | Arg | 3        | . | A  | B   | .  | . | .  | .   | -1.40 | 0.49  | * | *  | .   | -0.60 | 0.22 |
|    | Leu | 4        | . | A  | B   | .  | . | .  | .   | -1.30 | 0.74  | * | *  | .   | -0.60 | 0.18 |
|    | Val | 5        | . | A  | B   | .  | . | .  | .   | -1.26 | 1.04  | * | *  | .   | -0.60 | 0.20 |
| 10 | Leu | 6        | . | A  | B   | .  | . | .  | .   | -1.24 | 1.04  | * | *  | .   | -0.60 | 0.10 |
|    | Leu | 7        | A | A  | .   | .  | . | .  | .   | -1.49 | 1.54  | * | *  | .   | -0.60 | 0.12 |
|    | Trp | 8        | A | A  | .   | .  | . | .  | .   | -2.46 | 1.36  | * | *  | .   | -0.60 | 0.17 |
|    | Gly | 9        | A | A  | .   | .  | . | .  | .   | -2.34 | 1.36  | . | .  | .   | -0.60 | 0.15 |
|    | Ala | 10       | A | A  | .   | .  | . | .  | .   | -2.30 | 1.46  | . | .  | .   | -0.60 | 0.16 |
| 15 | Ala | 11       | A | A  | .   | .  | . | .  | .   | -2.30 | 1.46  | . | .  | .   | -0.60 | 0.12 |
|    | Val | 12       | . | A  | B   | .  | . | .  | .   | -1.83 | 1.23  | . | .  | .   | -0.60 | 0.10 |
|    | Phe | 13       | . | A  | B   | .  | . | .  | .   | -1.89 | 1.23  | . | .  | .   | -0.60 | 0.10 |
|    | Leu | 14       | . | A  | B   | .  | . | .  | .   | -1.83 | 1.16  | . | .  | .   | -0.60 | 0.10 |
|    | Leu | 15       | A | .  | .   | .  | . | T  | .   | -1.84 | 1.57  | . | .  | .   | -0.20 | 0.14 |
| 20 | Gly | 16       | . | .  | .   | .  | . | T  | C   | -1.84 | 1.54  | . | .  | .   | 0.00  | 0.16 |
|    | Gly | 17       | . | .  | .   | .  | T | T  | .   | -1.80 | 1.26  | . | .  | .   | 0.20  | 0.19 |
|    | Trp | 18       | A | .  | .   | .  | . | T  | .   | -1.44 | 1.26  | . | .  | .   | -0.20 | 0.19 |
|    | Met | 19       | . | .  | B   | .  | . | .  | .   | -0.63 | 1.00  | * | .  | .   | -0.40 | 0.19 |
|    | Ala | 20       | . | .  | B   | .  | . | .  | .   | -0.17 | 0.97  | . | .  | .   | -0.40 | 0.34 |
| 25 | Leu | 21       | . | .  | B   | .  | . | .  | .   | -0.17 | 0.97  | . | .  | .   | -0.40 | 0.32 |
|    | Gly | 22       | . | .  | .   | .  | . | T  | C   | -0.41 | 0.49  | . | .  | F   | 0.32  | 0.32 |
|    | Gln | 23       | . | .  | .   | .  | . | T  | C   | -0.71 | 0.37  | . | .  | F   | 0.79  | 0.32 |
|    | Gly | 24       | . | .  | .   | .  | . | T  | C   | -0.11 | 0.37  | . | .  | F   | 0.96  | 0.39 |
|    | Gly | 25       | . | .  | .   | .  | . | T  | C   | 0.13  | -0.31 | . | .  | F   | 1.73  | 0.69 |
| 30 | Ala | 26       | . | .  | .   | .  | . | .  | C   | 0.09  | -0.31 | . | .  | F   | 1.70  | 0.39 |
|    | Ala | 27       | A | .  | .   | .  | . | .  | .   | 0.43  | -0.07 | * | .  | F   | 1.33  | 0.29 |
|    | Glu | 28       | A | .  | .   | B  | . | .  | .   | -0.46 | -0.10 | . | *  | .   | 0.81  | 0.52 |
|    | Gly | 29       | A | .  | .   | B  | . | .  | .   | -0.11 | 0.16  | . | *  | .   | 0.04  | 0.36 |
|    | Val | 30       | A | .  | .   | B  | . | .  | .   | -0.66 | 0.06  | . | *  | .   | -0.13 | 0.61 |
|    | Gln | 31       | . | .  | B   | B  | . | .  | .   | -0.96 | 0.24  | . | *  | .   | -0.30 | 0.25 |
| 35 | Ile | 32       | . | .  | B   | B  | . | .  | .   | -0.61 | 0.93  | . | .  | .   | -0.60 | 0.18 |
|    | Gln | 33       | . | .  | B   | B  | . | .  | .   | -1.31 | 1.26  | . | .  | .   | -0.60 | 0.37 |
|    | Ile | 34       | . | .  | B   | B  | . | .  | .   | -0.97 | 1.40  | . | *  | .   | -0.60 | 0.19 |
|    | Ile | 35       | . | .  | B   | B  | . | .  | .   | -0.92 | 1.40  | . | *  | .   | -0.60 | 0.43 |
|    | Tyr | 36       | . | .  | B   | B  | . | .  | .   | -0.92 | 1.40  | . | *  | .   | -0.60 | 0.20 |
| 40 | Phe | 37       | . | .  | B   | B  | . | .  | .   | -0.34 | 1.00  | * | .  | .   | -0.60 | 0.50 |
|    | Asn | 38       | . | .  | .   | B  | . | .  | C   | -1.20 | 0.80  | * | .  | .   | -0.25 | 1.03 |
|    | Leu | 39       | . | .  | .   | B  | . | .  | C   | -0.31 | 0.76  | . | .  | .   | -0.40 | 0.49 |
|    | Glu | 40       | . | .  | B   | B  | . | .  | .   | -0.28 | 0.40  | . | *  | .   | -0.60 | 0.98 |
|    | Thr | 41       | . | .  | B   | B  | . | .  | .   | -0.34 | 0.26  | . | *  | .   | -0.30 | 0.45 |
| 45 | Val | 42       | . | .  | B   | B  | . | .  | .   | 0.07  | 0.34  | . | *  | .   | -0.30 | 0.79 |
|    | Gln | 43       | . | .  | B   | B  | . | .  | .   | 0.07  | 0.57  | . | *  | .   | -0.60 | 0.48 |
|    | Val | 44       | . | .  | B   | B  | . | .  | .   | 0.29  | 0.97  | . | *  | .   | -0.60 | 0.53 |
|    | Thr | 45       | . | .  | B   | B  | . | .  | .   | -0.01 | 0.99  | * | *  | .   | -0.60 | 0.73 |
|    | Trp | 46       | . | .  | B   | B  | . | .  | .   | 0.34  | 0.73  | * | *  | .   | -0.60 | 0.56 |
| 50 | Asn | 47       | A | .  | .   | B  | . | .  | .   | 0.96  | 0.33  | * | *  | .   | -0.15 | 1.51 |
|    | Ala | 48       | . | .  | .   | B  | T | .  | C   | 0.66  | 0.44  | * | *  | F   | 0.44  | 1.64 |
|    | Ser | 49       | . | .  | .   | .  | . | T  | C   | 1.62  | 0.34  | * | *  | F   | 1.28  | 2.10 |
|    | Lys | 50       | . | .  | .   | .  | T | T  | .   | 1.62  | -0.57 | * | *  | F   | 2.72  | 2.55 |
|    | Tyr | 51       | . | .  | .   | .  | T | T  | .   | 1.91  | -0.49 | * | *  | F   | 2.76  | 3.65 |
| 55 | Ser | 52       | . | .  | .   | .  | T | T  | .   | 1.10  | -0.59 | * | *  | F   | 3.40  | 4.38 |
|    | Arg | 53       | . | .  | .   | B  | T | .  | .   | 1.38  | -0.29 | * | *  | F   | 2.36  | 1.80 |
|    | Thr | 54       | . | .  | B   | B  | . | .  | .   | 0.98  | 0.20  | * | *  | F   | 1.02  | 1.66 |
|    | Asn | 55       | . | .  | B   | B  | . | .  | .   | 0.90  | 0.23  | * | *  | F   | 0.68  | 1.07 |
|    | Leu | 56       | . | .  | B   | B  | . | .  | .   | 0.90  | 0.34  | * | *  | .   | 0.04  | 0.75 |
| 60 | Thr | 57       | . | .  | B   | B  | . | .  | .   | 1.31  | 1.10  | * | *  | .   | -0.60 | 0.81 |
|    | Phe | 58       | . | .  | B   | B  | . | .  | .   | 0.50  | 0.61  | * | *  | .   | -0.60 | 0.99 |
|    | His | 59       | . | .  | B   | B  | . | .  | .   | 0.81  | 1.00  | . | *  | .   | -0.45 | 1.04 |
|    | Tyr | 60       | . | .  | B   | B  | . | .  | .   | 0.47  | 0.71  | . | *  | .   | -0.45 | 1.15 |
|    | Arg | 61       | . | .  | B   | B  | . | .  | .   | 1.28  | 0.66  | . | *  | .   | -0.45 | 1.32 |

668100-0219260



Table 1 (continued)

|    | Res | Position | I | II | III | IV | V | VI | VII | VIII  | IX    | X | XI | XII | XIII  | XIV  |
|----|-----|----------|---|----|-----|----|---|----|-----|-------|-------|---|----|-----|-------|------|
| 5  | Phe | 62       | . | .  | .   | B  | T | .  | .   | 1.59  | -0.13 | . | *  | .   | 1.19  | 1.62 |
|    | Asn | 63       | . | .  | .   | .  | T | T  | .   | 1.70  | -0.63 | . | *  | F   | 2.38  | 1.79 |
|    | Gly | 64       | . | .  | .   | .  | T | T  | .   | 1.49  | -0.89 | * | *  | F   | 2.57  | 0.92 |
|    | Asp | 65       | . | .  | .   | .  | T | T  | .   | 1.73  | -0.13 | . | *  | F   | 2.76  | 1.67 |
|    | Glu | 66       | . | .  | .   | .  | T | T  | .   | 1.62  | -0.91 | . | *  | F   | 3.40  | 1.73 |
| 10 | Ala | 67       | . | .  | .   | .  | T | .  | .   | 1.66  | -0.91 | . | .  | F   | 2.86  | 3.04 |
|    | Tyr | 68       | . | .  | .   | .  | T | .  | .   | 1.34  | -0.77 | * | .  | .   | 2.22  | 0.97 |
|    | Asp | 69       | . | .  | .   | .  | T | .  | .   | 1.69  | -0.29 | * | .  | F   | 1.73  | 0.81 |
|    | Gln | 70       | . | .  | .   | .  | T | .  | .   | 1.44  | 0.11  | * | .  | F   | 0.94  | 1.29 |
|    | Cys | 71       | . | .  | B   | .  | . | T  | .   | 0.63  | 0.37  | * | .  | F   | 0.40  | 1.29 |
| 15 | Thr | 72       | . | .  | B   | .  | . | T  | .   | 0.41  | 0.30  | * | .  | F   | 0.25  | 0.64 |
|    | Asn | 73       | . | .  | B   | .  | . | T  | .   | 0.66  | 0.99  | * | .  | .   | -0.20 | 0.30 |
|    | Tyr | 74       | . | .  | B   | .  | . | T  | .   | 0.66  | 0.99  | * | .  | .   | -0.20 | 0.98 |
|    | Leu | 75       | . | .  | B   | .  | . | .  | .   | 0.31  | 0.41  | * | .  | .   | -0.25 | 1.18 |
|    | Leu | 76       | . | .  | B   | .  | . | .  | .   | 0.94  | 0.36  | * | .  | .   | -0.10 | 0.73 |
| 20 | Gln | 77       | . | .  | B   | .  | . | .  | .   | 0.94  | 0.46  | * | .  | F   | 0.00  | 0.63 |
|    | Glu | 78       | . | .  | B   | .  | . | .  | .   | 0.64  | 0.19  | * | .  | F   | 0.70  | 1.10 |
|    | Gly | 79       | . | .  | .   | .  | T | .  | .   | 0.54  | -0.11 | * | .  | F   | 1.95  | 1.79 |
|    | His | 80       | . | .  | .   | .  | T | .  | .   | 0.69  | -0.37 | . | .  | F   | 2.20  | 1.02 |
|    | Thr | 81       | . | .  | .   | .  | T | T  | .   | 0.69  | -0.20 | * | .  | F   | 2.50  | 0.32 |
| 25 | Ser | 82       | . | .  | .   | .  | T | T  | .   | -0.12 | 0.49  | . | .  | F   | 1.35  | 0.26 |
|    | Gly | 83       | . | .  | .   | .  | T | T  | .   | -0.12 | 0.74  | . | .  | F   | 1.10  | 0.16 |
|    | Cys | 84       | . | .  | B   | .  | . | T  | .   | -0.37 | 0.24  | . | .  | .   | 0.60  | 0.19 |
|    | Leu | 85       | . | A  | B   | .  | . | .  | .   | -0.33 | 0.26  | . | .  | .   | -0.05 | 0.14 |
|    | Leu | 86       | A | A  | .   | .  | . | .  | .   | -0.02 | -0.13 | * | *  | .   | 0.30  | 0.24 |
| 30 | Asp | 87       | A | A  | .   | .  | . | .  | .   | 0.39  | -0.16 | * | *  | .   | 0.30  | 0.79 |
|    | Ala | 88       | A | A  | .   | .  | . | .  | .   | 0.73  | -0.73 | . | *  | F   | 0.90  | 1.87 |
|    | Glu | 89       | A | A  | .   | .  | . | .  | .   | 1.40  | -1.41 | . | *  | F   | 0.90  | 3.79 |
|    | Gln | 90       | A | .  | .   | .  | . | T  | .   | 1.32  | -2.10 | . | *  | F   | 1.30  | 3.79 |
|    | Arg | 91       | A | .  | .   | .  | . | T  | .   | 1.32  | -1.41 | . | *  | F   | 1.30  | 2.63 |
| 35 | Asp | 92       | A | .  | .   | .  | . | T  | .   | 1.08  | -1.23 | . | *  | F   | 1.30  | 1.25 |
|    | Asp | 93       | A | .  | .   | .  | . | T  | .   | 0.97  | -0.47 | . | *  | F   | 1.00  | 1.13 |
|    | Ile | 94       | . | .  | B   | B  | . | .  | .   | 0.67  | -0.09 | . | *  | .   | 0.30  | 0.50 |
|    | Leu | 95       | . | .  | B   | B  | . | .  | .   | -0.22 | 0.30  | . | *  | .   | -0.30 | 0.40 |
|    | Tyr | 96       | . | .  | B   | B  | . | .  | .   | -0.22 | 0.99  | . | *  | .   | -0.60 | 0.17 |
| 40 | Phe | 97       | . | .  | B   | B  | . | .  | .   | -0.22 | 0.99  | * | *  | .   | -0.60 | 0.47 |
|    | Ser | 98       | . | .  | B   | B  | . | .  | .   | -0.57 | 0.70  | * | *  | .   | -0.32 | 0.92 |
|    | Ile | 99       | . | .  | B   | .  | . | T  | .   | 0.01  | 0.44  | * | *  | .   | 0.36  | 0.58 |
|    | Arg | 100      | . | .  | .   | .  | T | T  | .   | 0.79  | 0.17  | * | *  | F   | 1.49  | 0.97 |
|    | Asn | 101      | . | .  | .   | .  | T | T  | .   | 0.82  | -0.11 | * | *  | F   | 2.37  | 0.98 |
| 45 | Gly | 102      | . | .  | .   | .  | T | T  | .   | 0.67  | -0.07 | * | *  | F   | 2.80  | 2.17 |
|    | Thr | 103      | . | .  | .   | B  | . | .  | C   | 0.27  | -0.11 | * | *  | F   | 1.77  | 0.82 |
|    | His | 104      | . | .  | .   | B  | . | .  | C   | 0.84  | 0.67  | * | *  | F   | 0.59  | 0.44 |
|    | Pro | 105      | . | .  | B   | B  | . | .  | .   | 0.14  | 0.76  | * | .  | .   | -0.04 | 0.65 |
|    | Val | 106      | . | .  | B   | B  | . | .  | .   | -0.16 | 0.83  | * | .  | .   | -0.32 | 0.45 |
| 50 | Phe | 107      | . | .  | B   | B  | . | .  | .   | 0.30  | 0.73  | * | .  | .   | -0.60 | 0.45 |
|    | Thr | 108      | . | .  | B   | B  | . | .  | .   | 0.32  | 0.23  | * | .  | .   | -0.30 | 0.56 |
|    | Ala | 109      | . | .  | B   | B  | . | .  | .   | -0.24 | 0.71  | * | .  | .   | -0.60 | 0.80 |
|    | Ser | 110      | A | .  | .   | B  | . | .  | .   | -0.89 | 0.69  | * | .  | .   | -0.60 | 0.91 |
|    | Arg | 111      | A | .  | .   | B  | . | .  | .   | -0.28 | 0.54  | * | .  | .   | -0.60 | 0.47 |
| 55 | Trp | 112      | . | .  | B   | B  | . | .  | .   | 0.18  | 0.81  | * | .  | .   | -0.60 | 0.73 |
|    | Met | 113      | . | .  | B   | B  | . | .  | .   | -0.32 | 1.07  | * | .  | .   | -0.60 | 0.85 |
|    | Val | 114      | . | .  | B   | B  | . | .  | .   | 0.31  | 1.37  | * | .  | .   | -0.60 | 0.36 |
|    | Tyr | 115      | . | .  | B   | B  | . | .  | .   | 0.40  | 1.37  | * | *  | .   | -0.60 | 0.68 |
|    | Tyr | 116      | . | .  | B   | .  | . | .  | .   | -0.01 | 0.89  | * | *  | .   | -0.25 | 1.07 |
| 60 | Leu | 117      | . | .  | B   | .  | . | .  | .   | -0.02 | 0.66  | . | .  | .   | 0.05  | 1.92 |
|    | Lys | 118      | . | .  | .   | .  | . | T  | C   | 0.37  | 0.40  | . | .  | F   | 0.90  | 1.65 |
|    | Pro | 119      | . | .  | .   | .  | T | T  | .   | 1.27  | 0.07  | . | .  | F   | 1.70  | 1.62 |
|    | Ser | 120      | . | .  | .   | .  | . | T  | C   | 1.48  | -0.69 | . | .  | F   | 2.70  | 3.94 |
|    | Ser | 121      | . | .  | .   | .  | . | T  | C   | 0.87  | -0.87 | * | *  | F   | 3.00  | 2.68 |

668780-0619260

Table 1 (continued)

|    | Res | Position | I | II | III | IV | V | VI | VII | VIII  | IX    | X | XI | XII | XIII  | XIV  |
|----|-----|----------|---|----|-----|----|---|----|-----|-------|-------|---|----|-----|-------|------|
| 5  | Pro | 122      | . | .  | B   | .  | . | .  | .   | 1.79  | -0.23 | * | *  | F   | 2.00  | 1.29 |
|    | Lys | 123      | . | .  | B   | .  | T | .  | .   | 1.04  | -0.66 | * | *  | F   | 2.40  | 1.88 |
|    | His | 124      | . | .  | B   | .  | . | .  | .   | 0.96  | -0.26 | * | *  | .   | 1.25  | 1.21 |
|    | Val | 125      | . | .  | B   | .  | . | .  | .   | 0.97  | -0.26 | * | *  | .   | 0.95  | 1.05 |
|    | Arg | 126      | . | .  | B   | .  | . | .  | .   | 1.23  | 0.23  | * | *  | .   | -0.10 | 0.55 |
| 10 | Phe | 127      | . | .  | B   | .  | . | .  | .   | 1.44  | 0.73  | * | *  | .   | -0.38 | 0.55 |
|    | Ser | 128      | . | .  | .   | .  | T | .  | .   | 1.40  | 0.63  | * | *  | .   | 0.19  | 1.29 |
|    | Trp | 129      | . | .  | .   | .  | T | .  | .   | 0.84  | -0.01 | . | *  | .   | 1.11  | 1.10 |
|    | His | 130      | . | .  | .   | .  | . | .  | C   | 0.84  | 0.49  | . | *  | .   | 0.03  | 1.28 |
|    | Gln | 131      | . | .  | .   | B  | T | .  | .   | 0.42  | 0.34  | . | *  | .   | 0.20  | 0.71 |
| 15 | Asp | 132      | . | .  | .   | B  | T | .  | .   | 0.27  | 0.44  | . | .  | .   | -0.12 | 0.98 |
|    | Ala | 133      | . | .  | B   | B  | . | .  | .   | 0.26  | 0.17  | . | .  | .   | -0.24 | 0.53 |
|    | Val | 134      | . | .  | B   | B  | . | .  | .   | -0.12 | 0.16  | . | .  | .   | -0.26 | 0.44 |
|    | Thr | 135      | . | .  | B   | B  | . | .  | .   | -0.39 | 0.33  | . | .  | .   | -0.28 | 0.14 |
|    | Val | 136      | . | .  | B   | B  | . | .  | .   | -0.39 | 0.71  | * | .  | .   | -0.60 | 0.19 |
| 20 | Thr | 137      | . | .  | B   | B  | . | .  | .   | -1.20 | 0.21  | * | .  | .   | -0.30 | 0.43 |
|    | Cys | 138      | . | .  | B   | .  | . | T  | .   | -0.91 | 0.26  | . | .  | .   | 0.10  | 0.24 |
|    | Ser | 139      | . | .  | B   | .  | . | T  | .   | -0.30 | 0.16  | . | .  | F   | 0.47  | 0.44 |
|    | Asp | 140      | . | .  | B   | .  | . | T  | .   | -0.33 | 0.27  | . | .  | F   | 0.69  | 0.48 |
|    | Leu | 141      | . | .  | B   | .  | . | T  | .   | 0.52  | 0.21  | . | .  | .   | 0.76  | 0.88 |
| 25 | Ser | 142      | . | .  | .   | .  | T | T  | .   | 0.02  | -0.36 | . | .  | .   | 2.13  | 1.10 |
|    | Tyr | 143      | . | .  | .   | .  | T | T  | .   | -0.12 | -0.06 | . | .  | .   | 2.20  | 0.54 |
|    | Gly | 144      | . | .  | .   | .  | T | T  | .   | -0.07 | 0.63  | . | .  | .   | 1.08  | 0.54 |
|    | Asp | 145      | . | .  | B   | .  | . | T  | .   | -0.07 | 0.70  | . | *  | .   | 0.46  | 0.63 |
|    | Leu | 146      | . | A  | B   | B  | . | .  | .   | -0.11 | 0.31  | * | *  | .   | 0.14  | 0.70 |
| 30 | Leu | 147      | . | A  | B   | B  | . | .  | .   | 0.19  | 0.20  | * | *  | .   | -0.08 | 0.52 |
|    | Tyr | 148      | . | A  | B   | B  | . | .  | .   | 0.19  | 0.17  | * | *  | .   | -0.30 | 0.54 |
|    | Glu | 149      | . | A  | B   | B  | . | .  | .   | 0.64  | 0.93  | * | *  | .   | -0.45 | 1.03 |
|    | Val | 150      | . | A  | B   | B  | . | .  | .   | 0.34  | 0.24  | * | *  | .   | -0.15 | 2.45 |
|    | Gln | 151      | . | .  | B   | B  | . | .  | .   | 0.94  | -0.06 | . | *  | .   | 0.45  | 2.10 |
| 35 | Tyr | 152      | . | .  | .   | .  | T | .  | .   | 1.06  | -0.39 | * | *  | .   | 1.39  | 1.87 |
|    | Arg | 153      | . | .  | B   | .  | . | .  | .   | 1.30  | 0.40  | . | *  | F   | 0.58  | 2.19 |
|    | Ser | 154      | . | .  | .   | .  | . | T  | C   | 0.99  | -0.24 | . | *  | F   | 2.22  | 2.11 |
|    | Pro | 155      | . | .  | .   | .  | . | T  | C   | 1.84  | -0.16 | . | *  | F   | 2.56  | 1.94 |
|    | Phe | 156      | . | .  | .   | .  | T | T  | .   | 1.56  | -0.91 | * | *  | F   | 3.40  | 1.72 |
| 40 | Asp | 157      | . | .  | .   | .  | . | T  | C   | 1.80  | 0.00  | * | *  | F   | 1.96  | 1.35 |
|    | Thr | 158      | . | .  | .   | .  | . | .  | C   | 1.39  | 0.01  | . | *  | F   | 1.76  | 1.51 |
|    | Glu | 159      | A | .  | .   | .  | . | .  | .   | 1.73  | -0.03 | . | *  | F   | 2.16  | 2.33 |
|    | Trp | 160      | A | .  | .   | .  | . | T  | .   | 1.94  | -0.81 | . | *  | F   | 2.66  | 2.80 |
|    | Gln | 161      | A | .  | .   | .  | . | T  | .   | 2.64  | -0.41 | . | *  | F   | 2.36  | 3.36 |
| 45 | Ser | 162      | . | .  | .   | .  | T | T  | .   | 2.64  | -0.90 | . | *  | F   | 3.40  | 3.36 |
|    | Lys | 163      | . | .  | .   | .  | T | T  | .   | 2.64  | -0.50 | . | *  | F   | 2.76  | 5.13 |
|    | Gln | 164      | . | .  | .   | .  | T | .  | .   | 1.98  | -0.93 | . | *  | F   | 2.69  | 4.28 |
|    | Glu | 165      | . | .  | .   | .  | T | .  | .   | 2.27  | -0.76 | . | *  | F   | 2.52  | 1.71 |
|    | Asn | 166      | . | .  | .   | .  | T | T  | .   | 1.41  | -0.74 | . | *  | F   | 2.55  | 1.38 |
| 50 | Thr | 167      | . | .  | .   | .  | T | T  | .   | 1.40  | -0.10 | . | *  | F   | 1.93  | 0.59 |
|    | Cys | 168      | . | .  | B   | .  | . | T  | .   | 0.47  | -0.01 | . | *  | F   | 1.70  | 0.49 |
|    | Asn | 169      | . | .  | B   | .  | . | T  | .   | 0.47  | 0.67  | . | *  | .   | 0.48  | 0.21 |
|    | Val | 170      | . | .  | B   | B  | . | .  | .   | 0.12  | 0.27  | * | *  | .   | 0.21  | 0.26 |
|    | Thr | 171      | . | .  | B   | B  | . | .  | .   | -0.69 | 0.21  | * | *  | .   | 0.04  | 0.47 |
| 55 | Ile | 172      | . | A  | B   | B  | . | .  | .   | -0.38 | 0.33  | * | *  | .   | -0.13 | 0.24 |
|    | Glu | 173      | A | A  | .   | B  | . | .  | .   | -0.30 | -0.07 | . | *  | .   | 0.30  | 0.55 |
|    | Gly | 174      | A | A  | .   | .  | . | .  | .   | -0.30 | -0.21 | . | *  | F   | 0.45  | 0.38 |
|    | Leu | 175      | A | A  | .   | .  | . | .  | .   | 0.60  | -0.70 | . | .  | .   | 0.60  | 0.95 |
|    | Asp | 176      | A | A  | .   | .  | . | .  | .   | 0.24  | -1.39 | . | .  | F   | 0.90  | 1.09 |
| 60 | Ala | 177      | A | A  | .   | .  | . | .  | .   | 0.89  | -0.81 | . | .  | F   | 0.75  | 0.59 |
|    | Glu | 178      | A | A  | .   | .  | . | .  | .   | 0.59  | -0.49 | . | .  | F   | 0.60  | 1.13 |
|    | Lys | 179      | A | A  | .   | B  | . | .  | .   | 0.23  | -0.79 | . | .  | .   | 0.60  | 0.90 |
|    | Cys | 180      | A | A  | .   | B  | . | .  | .   | 0.76  | 0.00  | . | .  | .   | -0.30 | 0.77 |
|    | Tyr | 181      | A | A  | .   | B  | . | .  | .   | -0.10 | 0.41  | * | *  | .   | -0.60 | 0.47 |

0937430-0219460

Table 1 (continued)

|    | Res | Position | I | II | III | IV | V | VI | VII | VIII  | IX    | X | XI | XII | XIII  | XIV  |
|----|-----|----------|---|----|-----|----|---|----|-----|-------|-------|---|----|-----|-------|------|
| 5  | Ser | 182      | A | .  | .   | B  | . | .  | .   | 0.60  | 1.06  | * | *  | .   | -0.60 | 0.17 |
|    | Phe | 183      | A | .  | .   | B  | . | .  | .   | -0.26 | 1.06  | * | *  | .   | -0.60 | 0.64 |
|    | Trp | 184      | A | .  | .   | B  | . | .  | .   | -0.26 | 1.13  | * | *  | .   | -0.60 | 0.30 |
|    | Val | 185      | A | .  | .   | B  | . | .  | .   | -0.18 | 0.37  | . | *  | .   | -0.30 | 0.45 |
|    | Arg | 186      | A | .  | .   | B  | . | .  | .   | -0.53 | 0.49  | . | *  | .   | -0.60 | 0.53 |
| 10 | Val | 187      | A | .  | .   | B  | . | .  | .   | -0.23 | 0.31  | . | *  | .   | -0.30 | 0.49 |
|    | Lys | 188      | A | .  | .   | B  | . | .  | .   | 0.47  | -0.60 | . | *  | .   | 0.75  | 1.15 |
|    | Ala | 189      | A | A  | .   | .  | . | .  | .   | -0.10 | -1.24 | . | *  | .   | 0.60  | 0.98 |
|    | Met | 190      | A | A  | .   | .  | . | .  | .   | 0.51  | -0.60 | . | *  | .   | 0.60  | 0.98 |
|    | Glu | 191      | . | A  | B   | .  | . | .  | .   | 0.06  | -0.49 | . | *  | .   | 0.30  | 0.77 |
| 15 | Asp | 192      | . | A  | B   | .  | . | .  | .   | 0.70  | -0.06 | * | *  | .   | 0.30  | 0.76 |
|    | Val | 193      | . | A  | B   | .  | . | .  | .   | 0.66  | -0.13 | * | .  | .   | 0.45  | 1.18 |
|    | Tyr | 194      | . | A  | B   | .  | . | .  | .   | 0.93  | -0.74 | . | .  | .   | 0.75  | 1.14 |
|    | Gly | 195      | . | .  | B   | .  | . | T  | .   | 1.29  | -0.26 | . | .  | F   | 0.85  | 0.98 |
|    | Pro | 196      | . | .  | .   | .  | T | T  | .   | 1.08  | 0.50  | . | .  | F   | 0.50  | 2.08 |
| 20 | Asp | 197      | . | .  | .   | .  | T | T  | .   | 0.78  | 0.29  | . | .  | F   | 0.80  | 2.05 |
|    | Thr | 198      | . | .  | .   | .  | . | T  | C   | 1.63  | -0.09 | . | .  | F   | 1.48  | 2.77 |
|    | Tyr | 199      | . | .  | B   | .  | . | .  | .   | 1.59  | -0.51 | . | .  | F   | 1.66  | 3.00 |
|    | Pro | 200      | . | .  | .   | .  | . | T  | C   | 1.63  | -0.03 | * | .  | F   | 2.04  | 1.89 |
|    | Ser | 201      | . | .  | .   | .  | T | T  | .   | 1.84  | 0.36  | * | .  | F   | 1.92  | 1.75 |
| 25 | Asp | 202      | . | .  | .   | .  | T | T  | .   | 0.99  | -0.13 | * | .  | F   | 2.80  | 1.94 |
|    | Trp | 203      | . | .  | .   | .  | T | T  | .   | 0.99  | -0.24 | . | .  | F   | 2.37  | 0.93 |
|    | Ser | 204      | . | .  | B   | B  | . | .  | .   | 0.57  | -0.19 | . | .  | F   | 1.44  | 1.00 |
|    | Glu | 205      | . | .  | B   | B  | . | .  | .   | 0.49  | 0.00  | . | .  | .   | 0.26  | 0.32 |
|    | Val | 206      | . | .  | B   | B  | . | .  | .   | 0.79  | 0.91  | * | .  | .   | -0.32 | 0.32 |
| 30 | Thr | 207      | . | .  | B   | B  | . | .  | .   | 0.90  | 0.40  | * | .  | .   | -0.60 | 0.42 |
|    | Cys | 208      | . | .  | .   | B  | T | .  | .   | 0.84  | 0.01  | * | *  | .   | 0.10  | 0.47 |
|    | Trp | 209      | . | .  | .   | .  | T | T  | .   | 1.14  | 0.44  | . | *  | .   | 0.20  | 0.63 |
|    | Gln | 210      | A | .  | .   | .  | T | T  | .   | 0.26  | -0.20 | * | *  | .   | 0.70  | 0.75 |
|    | Arg | 211      | . | .  | .   | .  | T | T  | .   | 1.22  | 0.00  | * | *  | F   | 0.65  | 0.98 |
| 35 | Gly | 212      | . | .  | .   | .  | T | T  | .   | 1.53  | -0.57 | * | *  | F   | 1.70  | 1.83 |
|    | Glu | 213      | . | A  | .   | .  | T | .  | .   | 1.61  | -1.49 | * | *  | F   | 1.30  | 1.76 |
|    | Ile | 214      | . | A  | .   | .  | T | .  | .   | 1.23  | -1.39 | * | *  | F   | 1.15  | 0.91 |
|    | Arg | 215      | . | A  | .   | .  | T | .  | .   | 0.64  | -0.81 | . | *  | F   | 1.15  | 0.49 |
| 40 | Asp | 216      | . | A  | .   | .  | T | .  | .   | 0.53  | -0.74 | * | *  | F   | 1.15  | 0.29 |
|    | Ala | 217      | . | A  | .   | .  | T | .  | .   | 0.57  | -0.74 | * | *  | .   | 1.00  | 0.71 |
|    | Cys | 218      | . | A  | B   | .  | . | .  | .   | 0.36  | -0.94 | * | *  | .   | 0.60  | 0.52 |
|    | Ala | 219      | . | A  | .   | .  | T | .  | .   | 0.93  | -0.51 | * | .  | .   | 1.00  | 0.48 |
|    | Glu | 220      | . | A  | .   | .  | T | .  | .   | 0.61  | -0.03 | . | .  | F   | 0.85  | 0.69 |
| 45 | Thr | 221      | . | A  | .   | .  | . | .  | C   | 0.40  | -0.10 | . | *  | F   | 1.14  | 2.00 |
|    | Pro | 222      | . | A  | .   | .  | T | .  | .   | 1.03  | -0.24 | . | .  | F   | 1.68  | 3.06 |
|    | Thr | 223      | . | .  | .   | .  | . | .  | C   | 1.49  | -0.74 | . | *  | F   | 2.32  | 3.53 |
|    | Pro | 224      | . | .  | .   | .  | . | T  | C   | 2.12  | -0.31 | . | *  | F   | 2.56  | 3.78 |
|    | Pro | 225      | . | .  | .   | .  | T | T  | .   | 1.31  | -0.80 | . | *  | F   | 3.40  | 4.89 |
|    | Lys | 226      | . | .  | .   | .  | . | T  | C   | 1.32  | -0.54 | . | *  | F   | 2.86  | 2.80 |
| 50 | Pro | 227      | A | .  | .   | .  | . | T  | .   | 1.58  | -0.64 | * | .  | F   | 2.32  | 2.42 |
|    | Lys | 228      | A | .  | .   | .  | . | .  | .   | 1.19  | -1.07 | * | *  | F   | 1.78  | 3.13 |
|    | Leu | 229      | A | .  | .   | B  | . | .  | .   | 0.51  | -0.71 | * | *  | F   | 1.24  | 1.36 |
|    | Ser | 230      | . | .  | B   | B  | . | .  | .   | -0.09 | -0.03 | . | *  | F   | 0.45  | 0.62 |
| 55 | Lys | 231      | . | .  | B   | B  | . | .  | .   | -1.02 | 0.23  | . | *  | .   | -0.30 | 0.25 |
|    | Phe | 232      | . | .  | B   | B  | . | .  | .   | -1.11 | 0.91  | * | .  | .   | -0.60 | 0.22 |
|    | Ile | 233      | . | .  | B   | B  | . | .  | .   | -1.46 | 0.61  | . | .  | .   | -0.60 | 0.22 |
|    | Leu | 234      | . | .  | B   | B  | . | .  | .   | -1.46 | 0.61  | . | .  | .   | -0.60 | 0.14 |
|    | Ile | 235      | . | .  | B   | B  | . | .  | .   | -1.74 | 1.30  | . | .  | .   | -0.60 | 0.14 |
| 60 | Ser | 236      | A | .  | .   | B  | . | .  | .   | -2.68 | 1.01  | . | .  | .   | -0.60 | 0.20 |
|    | Ser | 237      | A | .  | .   | B  | . | .  | .   | -2.79 | 1.01  | . | .  | .   | -0.60 | 0.17 |
|    | Leu | 238      | A | .  | .   | B  | . | .  | .   | -2.71 | 1.01  | . | .  | .   | -0.60 | 0.20 |
|    | Ala | 239      | A | .  | .   | B  | . | .  | .   | -2.50 | 1.01  | . | .  | .   | -0.60 | 0.12 |
|    | Ile | 240      | A | .  | .   | B  | . | .  | .   | -2.47 | 1.24  | . | .  | .   | -0.60 | 0.09 |
|    | Leu | 241      | A | .  | .   | B  | . | .  | .   | -2.47 | 1.50  | . | *  | .   | -0.60 | 0.08 |

668780\*0249460

Table 1 (continued)

|    | Res | Position | I | II | III | IV | V | VI | VII | VIII  | IX    | X | XI | XII | XIII  | XIV  |
|----|-----|----------|---|----|-----|----|---|----|-----|-------|-------|---|----|-----|-------|------|
| 5  | Leu | 242      | A | .  | .   | B  | . | .  | .   | -2.98 | 1.20  | . | .  | .   | -0.60 | 0.11 |
|    | Met | 243      | . | .  | B   | B  | . | .  | .   | -2.98 | 1.39  | . | *  | .   | -0.60 | 0.13 |
|    | Val | 244      | A | .  | .   | B  | . | .  | .   | -3.20 | 1.39  | . | .  | .   | -0.60 | 0.13 |
|    | Ser | 245      | A | .  | .   | B  | . | .  | .   | -3.12 | 1.39  | . | .  | .   | -0.60 | 0.13 |
|    | Leu | 246      | A | .  | .   | B  | . | .  | .   | -2.61 | 1.39  | . | .  | .   | -0.60 | 0.11 |
| 10 | Leu | 247      | A | .  | .   | B  | . | .  | .   | -2.61 | 1.16  | . | .  | .   | -0.60 | 0.19 |
|    | Leu | 248      | A | .  | .   | B  | . | .  | .   | -2.30 | 1.20  | * | .  | .   | -0.60 | 0.12 |
|    | Leu | 249      | A | .  | .   | B  | . | .  | .   | -1.40 | 1.73  | * | .  | .   | -0.60 | 0.15 |
|    | Ser | 250      | A | .  | .   | B  | . | .  | .   | -1.91 | 1.04  | * | .  | .   | -0.60 | 0.36 |
|    | Leu | 251      | A | .  | .   | B  | . | .  | .   | -1.39 | 1.04  | * | .  | .   | -0.60 | 0.36 |
| 15 | Trp | 252      | A | .  | .   | B  | . | .  | .   | -0.47 | 1.27  | * | .  | .   | -0.60 | 0.46 |
|    | Lys | 253      | A | .  | .   | B  | . | .  | .   | -0.51 | 0.59  | * | *  | .   | -0.60 | 0.68 |
|    | Leu | 254      | A | .  | .   | B  | . | .  | .   | 0.34  | 0.84  | * | *  | .   | -0.60 | 0.61 |
|    | Trp | 255      | A | .  | .   | B  | . | .  | .   | 0.69  | 0.16  | * | .  | .   | -0.15 | 1.16 |
|    | Arg | 256      | A | .  | .   | B  | . | .  | .   | 0.80  | -0.76 | * | .  | .   | 0.75  | 1.16 |
| 20 | Val | 257      | A | .  | .   | B  | . | .  | .   | 0.28  | 0.03  | * | .  | .   | -0.15 | 1.21 |
|    | Lys | 258      | . | .  | B   | B  | . | .  | .   | -0.66 | 0.03  | * | .  | .   | -0.30 | 0.95 |
|    | Lys | 259      | . | .  | B   | B  | . | .  | .   | -0.06 | -0.20 | * | .  | .   | 0.30  | 0.34 |
|    | Phe | 260      | . | .  | B   | B  | . | .  | .   | -0.07 | 0.23  | * | *  | .   | -0.30 | 0.71 |
|    | Leu | 261      | . | .  | B   | B  | . | .  | .   | -1.03 | -0.03 | * | *  | .   | 0.30  | 0.48 |
| 25 | Ile | 262      | . | .  | B   | B  | . | .  | .   | -0.39 | 0.61  | * | .  | .   | -0.60 | 0.18 |
|    | Pro | 263      | . | .  | B   | .  | . | .  | .   | -0.43 | 1.04  | . | .  | .   | -0.10 | 0.32 |
|    | Ser | 264      | . | .  | B   | .  | . | .  | .   | -0.69 | 0.26  | * | *  | F   | 0.65  | 0.64 |
|    | Val | 265      | . | .  | .   | .  | . | .  | C   | 0.06  | 0.00  | . | .  | F   | 1.30  | 1.41 |
|    | Pro | 266      | . | .  | .   | .  | . | .  | C   | 0.57  | -0.69 | . | .  | F   | 2.50  | 1.82 |
| 30 | Asp | 267      | . | .  | .   | .  | . | T  | C   | 0.57  | -0.73 | . | .  | F   | 3.00  | 1.82 |
|    | Pro | 268      | . | .  | .   | .  | T | T  | .   | 0.08  | -0.43 | . | .  | F   | 2.60  | 1.72 |
|    | Lys | 269      | . | .  | .   | .  | T | T  | .   | 0.17  | -0.29 | . | .  | F   | 2.15  | 0.96 |
|    | Ser | 270      | . | .  | B   | .  | . | T  | .   | 0.68  | -0.29 | * | .  | F   | 1.45  | 0.89 |
|    | He  | 271      | . | .  | B   | .  | . | .  | .   | 0.08  | 0.14  | * | .  | .   | 0.20  | 0.57 |
| 35 | Phe | 272      | . | .  | B   | .  | . | T  | .   | -0.62 | 0.40  | * | .  | .   | -0.20 | 0.24 |
|    | Pro | 273      | . | .  | B   | .  | . | T  | .   | -0.41 | 1.19  | * | *  | .   | -0.20 | 0.15 |
|    | Gly | 274      | . | .  | .   | .  | . | T  | C   | -1.34 | 0.80  | * | *  | .   | 0.00  | 0.38 |
|    | Leu | 275      | . | .  | B   | .  | . | T  | .   | -1.08 | 0.80  | * | .  | .   | -0.20 | 0.30 |
|    | Phe | 276      | . | .  | B   | .  | . | .  | .   | -0.19 | 0.51  | * | .  | .   | -0.40 | 0.27 |
| 40 | Glu | 277      | A | .  | .   | .  | . | .  | .   | 0.17  | 0.49  | . | *  | .   | -0.40 | 0.47 |
|    | Ile | 278      | A | .  | .   | .  | . | .  | .   | 0.38  | 0.49  | . | .  | .   | -0.40 | 0.56 |
|    | His | 279      | A | .  | .   | .  | . | T  | .   | 0.02  | 0.20  | * | *  | .   | 0.25  | 1.04 |
|    | Gln | 280      | . | .  | .   | .  | . | T  | C   | 0.83  | 0.20  | * | .  | F   | 0.45  | 0.52 |
|    | Gly | 281      | . | .  | .   | .  | . | T  | C   | 1.53  | 0.60  | * | .  | F   | 0.30  | 1.29 |
| 45 | Asn | 282      | . | .  | .   | .  | . | T  | C   | 1.24  | -0.09 | . | .  | F   | 1.20  | 1.64 |
|    | Phe | 283      | . | A  | .   | .  | . | .  | C   | 1.24  | 0.33  | . | .  | F   | 0.05  | 1.00 |
|    | Gln | 284      | . | A  | .   | .  | . | .  | C   | 0.97  | 0.61  | * | *  | .   | -0.16 | 0.71 |
|    | Glu | 285      | . | A  | B   | .  | . | .  | .   | 0.97  | 0.67  | * | *  | .   | -0.12 | 0.63 |
|    | Trp | 286      | . | A  | B   | .  | . | .  | .   | 1.00  | 0.27  | * | *  | .   | 0.57  | 1.22 |
| 50 | Ile | 287      | . | A  | B   | .  | . | .  | .   | 1.00  | -0.03 | * | .  | .   | 1.41  | 1.02 |
|    | Thr | 288      | . | .  | .   | .  | T | .  | .   | 1.70  | -0.03 | * | .  | F   | 2.40  | 1.02 |
|    | Asp | 289      | . | .  | .   | .  | T | .  | .   | 0.84  | 0.37  | * | .  | F   | 1.56  | 1.56 |
|    | Thr | 290      | A | .  | .   | .  | . | .  | .   | 0.26  | 0.10  | * | .  | F   | 0.92  | 1.65 |
|    | Gln | 291      | A | A  | .   | .  | . | .  | .   | 0.51  | -0.09 | * | .  | F   | 1.08  | 1.15 |
| 55 | Asn | 292      | A | A  | .   | .  | . | .  | .   | 0.59  | -0.07 | . | .  | F   | 0.69  | 0.94 |
|    | Val | 293      | A | A  | .   | .  | . | .  | .   | 0.87  | 0.61  | * | .  | .   | -0.60 | 0.54 |
|    | Ala | 294      | A | A  | .   | .  | . | .  | .   | 0.91  | 0.63  | * | .  | .   | -0.60 | 0.42 |
|    | His | 295      | A | A  | .   | .  | . | .  | .   | 0.62  | 0.23  | * | .  | .   | -0.30 | 0.52 |
|    | Leu | 296      | A | A  | .   | .  | . | .  | .   | 0.03  | 0.44  | * | .  | .   | -0.60 | 0.70 |
| 60 | His | 297      | A | A  | .   | .  | . | .  | .   | -0.31 | 0.30  | * | .  | .   | -0.30 | 0.70 |
|    | Lys | 298      | A | A  | .   | .  | . | .  | .   | -0.04 | 0.23  | * | .  | .   | -0.30 | 0.51 |
|    | Met | 299      | A | A  | .   | .  | . | .  | .   | 0.54  | 0.23  | . | .  | .   | -0.30 | 0.62 |
|    | Ala | 300      | A | A  | .   | .  | . | .  | .   | 0.58  | -0.46 | * | .  | .   | 0.30  | 0.79 |
|    | Gly | 301      | A | A  | .   | .  | . | .  | .   | 1.39  | -0.56 | * | .  | .   | 0.60  | 0.69 |

553780\*0E492E60

Table 1 (continued)

|    | Res | Position | I | II | III | IV | V | VI | VII | VIII  | IX    | X | XI | XII | XIII  | XIV  |
|----|-----|----------|---|----|-----|----|---|----|-----|-------|-------|---|----|-----|-------|------|
| 5  | Ala | 302      | A | A  | .   | .  | . | .  | .   | 1.12  | -0.56 | . | .  | F   | 0.90  | 1.20 |
|    | Glu | 303      | A | A  | .   | .  | . | .  | .   | 0.73  | -0.79 | . | .  | F   | 1.20  | 1.60 |
|    | Gln | 304      | A | A  | .   | .  | . | .  | .   | 1.12  | -0.86 | . | .  | F   | 1.50  | 1.60 |
|    | Glu | 305      | . | A  | .   | .  | . | .  | C   | 1.71  | -0.86 | . | .  | F   | 2.00  | 2.44 |
|    | Ser | 306      | . | A  | .   | .  | . | .  | C   | 2.06  | -1.36 | . | .  | F   | 2.30  | 2.44 |
| 10 | Gly | 307      | . | .  | .   | .  | . | T  | C   | 2.43  | -1.36 | . | .  | F   | 3.00  | 2.44 |
|    | Pro | 308      | . | .  | .   | .  | . | T  | C   | 1.62  | -1.33 | . | .  | F   | 2.70  | 2.18 |
|    | Glu | 309      | . | .  | .   | .  | . | T  | C   | 0.77  | -0.64 | . | .  | F   | 2.40  | 1.34 |
|    | Glu | 310      | A | .  | .   | .  | . | T  | .   | -0.09 | -0.39 | . | .  | F   | 1.60  | 1.01 |
|    | Pro | 311      | A | .  | .   | .  | . | .  | .   | 0.21  | -0.17 | . | .  | F   | 0.95  | 0.48 |
| 15 | Leu | 312      | A | A  | .   | .  | . | .  | .   | -0.26 | -0.20 | . | .  | .   | 0.30  | 0.48 |
|    | Val | 313      | A | A  | .   | .  | . | .  | .   | -0.63 | 0.49  | * | .  | .   | -0.60 | 0.23 |
|    | Val | 314      | A | A  | .   | .  | . | .  | .   | -0.59 | 0.99  | * | .  | .   | -0.60 | 0.15 |
|    | Gln | 315      | A | A  | .   | .  | . | .  | .   | -0.90 | 0.56  | * | .  | .   | -0.60 | 0.36 |
|    | Leu | 316      | A | A  | .   | .  | . | .  | .   | -0.69 | 0.36  | * | .  | .   | -0.30 | 0.71 |
| 20 | Ala | 317      | A | A  | .   | .  | . | .  | .   | -0.47 | -0.29 | * | *  | .   | 0.45  | 1.65 |
|    | Lys | 318      | A | A  | .   | .  | . | .  | .   | 0.39  | -0.43 | * | .  | F   | 0.45  | 0.96 |
|    | Thr | 319      | A | A  | .   | .  | . | .  | .   | 0.94  | -0.83 | * | .  | F   | 0.90  | 2.03 |
|    | Glu | 320      | A | A  | .   | .  | . | .  | .   | 0.73  | -1.13 | * | .  | F   | 0.90  | 2.69 |
|    | Ala | 321      | A | A  | .   | .  | . | .  | .   | 1.66  | -1.20 | * | .  | F   | 1.24  | 2.08 |
| 25 | Glu | 322      | A | A  | .   | .  | . | .  | .   | 1.64  | -1.20 | * | *  | F   | 1.58  | 2.82 |
|    | Ser | 323      | A | .  | .   | .  | . | T  | .   | 0.79  | -1.07 | * | .  | F   | 2.32  | 1.61 |
|    | Pro | 324      | A | .  | .   | .  | . | T  | .   | 1.10  | -0.39 | * | .  | F   | 2.36  | 1.32 |
|    | Arg | 325      | . | .  | .   | .  | T | T  | .   | 0.89  | -0.89 | * | .  | F   | 3.40  | 1.27 |
|    | Met | 326      | A | .  | .   | .  | . | T  | .   | 1.48  | -0.46 | * | .  | .   | 2.21  | 1.46 |
| 30 | Leu | 327      | A | .  | .   | .  | . | .  | .   | 1.17  | -0.44 | * | .  | F   | 1.82  | 1.64 |
|    | Asp | 328      | A | .  | .   | .  | . | T  | .   | 1.47  | -0.39 | * | .  | F   | 1.68  | 1.21 |
|    | Pro | 329      | A | .  | .   | .  | . | T  | .   | 1.68  | -0.39 | * | *  | F   | 1.34  | 2.11 |
|    | Gln | 330      | A | .  | .   | .  | . | T  | .   | 1.61  | -1.00 | * | .  | F   | 1.30  | 4.44 |
|    | Thr | 331      | A | .  | .   | .  | . | T  | .   | 2.21  | -1.69 | . | *  | F   | 1.30  | 5.31 |
| 35 | Glu | 332      | A | A  | .   | .  | . | .  | .   | 2.43  | -1.69 | . | *  | F   | 0.90  | 5.95 |
|    | Glu | 333      | A | A  | .   | .  | . | .  | .   | 2.13  | -1.61 | . | .  | F   | 0.90  | 3.47 |
|    | Lys | 334      | A | A  | .   | .  | . | .  | .   | 2.00  | -1.63 | . | .  | F   | 1.15  | 3.22 |
|    | Glu | 335      | A | A  | .   | .  | . | .  | .   | 1.66  | -1.69 | . | .  | F   | 1.40  | 1.84 |
|    | Ala | 336      | A | .  | .   | .  | . | T  | .   | 1.67  | -1.26 | . | .  | F   | 2.05  | 1.05 |
| 40 | Ser | 337      | A | .  | .   | .  | . | T  | .   | 0.86  | -0.87 | . | .  | F   | 2.15  | 0.71 |
|    | Gly | 338      | . | .  | .   | .  | T | T  | .   | 0.86  | -0.19 | . | *  | F   | 2.50  | 0.34 |
|    | Gly | 339      | . | .  | .   | .  | T | T  | .   | -0.00 | 0.21  | * | .  | F   | 1.65  | 0.58 |
|    | Ser | 340      | . | .  | .   | .  | . | .  | C   | -0.21 | 0.40  | . | .  | F   | 0.70  | 0.35 |
|    | Leu | 341      | . | .  | .   | .  | . | .  | C   | 0.34  | 0.44  | . | .  | .   | 0.30  | 0.55 |
| 45 | Gln | 342      | . | .  | B   | .  | . | .  | .   | 0.64  | 0.51  | . | *  | .   | -0.15 | 0.76 |
|    | Leu | 343      | . | .  | B   | .  | . | .  | .   | 0.78  | 0.49  | . | *  | .   | -0.40 | 0.98 |
|    | Pro | 344      | . | .  | B   | .  | . | .  | .   | 0.31  | 0.53  | * | *  | .   | -0.25 | 1.84 |
|    | His | 345      | . | .  | B   | .  | . | .  | .   | 0.61  | 0.53  | * | *  | .   | -0.40 | 0.88 |
|    | Gln | 346      | . | .  | B   | .  | . | .  | .   | 1.08  | 0.53  | * | .  | F   | 0.03  | 1.84 |
| 50 | Pro | 347      | . | .  | B   | .  | . | .  | .   | 0.73  | 0.27  | * | .  | F   | 0.46  | 1.18 |
|    | Leu | 348      | . | .  | .   | .  | T | T  | .   | 1.54  | 0.27  | * | .  | F   | 1.04  | 0.86 |
|    | Gln | 349      | . | .  | .   | .  | T | T  | .   | 0.90  | -0.23 | * | .  | F   | 1.77  | 0.83 |
|    | Gly | 350      | . | .  | .   | .  | T | T  | .   | 0.08  | 0.01  | * | .  | F   | 1.30  | 0.40 |
|    | Gly | 351      | . | .  | B   | .  | . | T  | .   | -0.23 | 0.23  | . | .  | F   | 0.77  | 0.36 |
| 55 | Asp | 352      | . | .  | B   | B  | . | .  | .   | -0.91 | 0.03  | . | .  | F   | 0.24  | 0.30 |
|    | Val | 353      | . | .  | B   | B  | . | .  | .   | -0.44 | 0.31  | . | .  | F   | 0.11  | 0.21 |
|    | Val | 354      | . | .  | B   | B  | . | .  | .   | -0.79 | 0.31  | . | .  | .   | -0.17 | 0.21 |
|    | Thr | 355      | . | .  | B   | B  | . | .  | .   | -1.14 | 0.31  | . | .  | .   | -0.30 | 0.13 |
|    | Ile | 356      | . | .  | B   | B  | . | .  | .   | -1.11 | 1.10  | . | .  | .   | -0.60 | 0.15 |
| 60 | Gly | 357      | . | .  | B   | B  | . | .  | .   | -1.81 | 0.94  | . | .  | .   | -0.60 | 0.28 |
|    | Gly | 358      | . | .  | B   | B  | . | .  | .   | -1.81 | 1.09  | . | .  | .   | -0.60 | 0.17 |
|    | Phe | 359      | . | .  | B   | B  | . | .  | .   | -1.56 | 1.24  | . | .  | .   | -0.60 | 0.18 |
|    | Thr | 360      | . | .  | B   | B  | . | .  | .   | -1.24 | 1.17  | . | .  | .   | -0.60 | 0.18 |
|    | Phe | 361      | . | .  | B   | B  | . | .  | .   | -0.36 | 1.14  | . | *  | .   | -0.60 | 0.29 |

66370\*0649460

Table 1 (continued)

|    | Res Position | I | II | III | IV | V | VI | VII | VIII  | IX    | X | XI | XII | XIII  | XIV  |
|----|--------------|---|----|-----|----|---|----|-----|-------|-------|---|----|-----|-------|------|
| 5  | Val 362      | . | .  | B   | B  | . | .  | .   | 0.10  | 0.71  | . | *  | .   | -0.32 | 0.57 |
|    | Met 363      | . | .  | B   | B  | . | .  | .   | 0.14  | -0.07 | . | *  | .   | 0.86  | 0.77 |
|    | Asn 364      | . | .  | .   | B  | T | .  | .   | 0.60  | -0.17 | . | *  | .   | 1.69  | 1.19 |
|    | Asp 365      | . | .  | .   | .  | T | T  | .   | 0.06  | -0.20 | . | *  | .   | 2.37  | 2.51 |
|    | Arg 366      | . | .  | .   | .  | T | T  | .   | 0.17  | -0.20 | . | *  | F   | 2.80  | 1.88 |
| 10 | Ser 367      | A | .  | .   | .  | . | T  | .   | 0.21  | -0.31 | . | *  | .   | 1.97  | 1.18 |
|    | Tyr 368      | A | .  | .   | .  | . | T  | .   | 0.42  | -0.03 | . | *  | .   | 1.54  | 0.58 |
|    | Val 369      | . | A  | B   | .  | . | .  | .   | 0.03  | 0.40  | . | *  | .   | -0.04 | 0.38 |
|    | Ala 370      | . | A  | B   | .  | . | .  | .   | -0.36 | 0.83  | * | .  | .   | -0.32 | 0.36 |
|    | Leu 371      | . | A  | B   | .  | . | .  | .   | -0.86 | 0.87  | * | *  | .   | -0.60 | 0.30 |
| 15 | Ter 372      | . | A  | B   | .  | . | .  | .   | -0.94 | 0.54  | . | .  | .   | -0.60 | 0.51 |

Among highly preferred fragments in this regard are those that comprise regions of CRCGCL that combine several structural features, such as several of the features set out above.

Other preferred fragments are biologically active CRCGCL fragments. Biologically active fragments are those exhibiting activity similar, but not necessarily identical, to an activity of the CRCGCL polypeptide. The biological activity of the fragments may include an improved desired activity, or a decreased undesirable activity.

However, many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:1 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. One embodiment of the present invention excludes Genbank Accession No. X91553 (herein incorporated by reference in its entirety.) Moreover, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1559 of SEQ ID NO:1, b is an integer of 15 to 1573, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:1, and where the b is greater than or equal to a + 14.

### Epitope-Bearing Portions

In another aspect, the invention provides peptides and polypeptides comprising epitope-bearing portions of the polypeptides of the present invention.

These epitopes are immunogenic or antigenic epitopes of the polypeptides of the present invention. An "immunogenic epitope" is defined as a part of a protein that elicits an antibody response in vivo when the whole polypeptide of the present invention, or fragment thereof, is the immunogen. On the other hand, a region of a polypeptide to which an antibody can bind is defined as an "antigenic determinant" or "antigenic epitope." The number of in vivo immunogenic epitopes of a protein generally is less than the number of antigenic epitopes. See, e.g., Geysen, et al. (1983) Proc. Natl. Acad. Sci. USA 81:3998- 4002. However, antibodies can be made to any antigenic epitope, regardless of whether it is an immunogenic epitope, by using methods such as phage display. See e.g., Petersen G. et al. (1995) Mol. Gen. Genet. 249:425-431. Therefore, included in the present invention are both immunogenic epitopes and antigenic epitopes.

A list of exemplified amino acid sequences comprising immunogenic epitopes are shown in Table 1 below. It is pointed out that Table 1 only lists amino acid residues comprising epitopes predicted to have the highest degree of antigenicity using the algorithm of Jameson and Wolf, (1988) Comp. Appl. Biosci. 4:181-186 (said references incorporated by reference in their entireties). The Jameson-Wolf antigenic analysis was performed using the computer program PROTEAN, using default parameters (Version 3.11 for the Power MacIntosh, DNASTAR, Inc., 1228 South Park Street Madison, WI). Table 1 and portions of polypeptides not listed in Table 1 are not considered non-immunogenic. The immunogenic epitopes of Table 1 is an exemplified list, not an exhaustive list, because other immunogenic epitopes are merely not recognized as such by the particular algorithm used. Amino acid residues comprising other immunogenic epitopes may be routinely determined using algorithms similar to the Jameson-Wolf analysis or by in vivo testing for an antigenic response using methods known in the art. See, e.g., Geysen et al., supra; U.S. Patents 4,708,781; 5, 194,392; 4,433,092; and 5,480,971 (said references incorporated by reference in their entireties).

Antigenic epitope-bearing peptides and polypeptides of the invention preferably contain a sequence of at least seven, more preferably at least nine and most preferably between about 15 to about 30 amino acids contained within the amino acid sequence of a polypeptide of the invention. Non-limiting examples antigenic polypeptides or peptides that can be used to CRGCL -specific antibodies include:

22-29; 48-56; 62-73; 78-85; 88-95; 99-105; 118-126; 139-146; 151-169; 188-206;  
208-231; 264-271; 286-293; 300-313; 317-342; 347-353; 363-369 amino acid  
residues of SEQ ID NO:2 These polypeptide fragments have been determined to bear  
antigenic epitopes of the CRCGCL protein by the analysis of the Jameson-Wolf  
5 antigenic index, as shown in Figure 3, above.

It is particularly pointed out that the amino acid sequences of Table 1  
comprise immunogenic epitopes. Table 1 lists only the critical residues of  
immunogenic epitopes determined by the Jameson-Wolf analysis. Thus, additional  
flanking residues on either the N-terminal, C-terminal, or both N- and C-terminal  
10 ends may be added to the sequences of Table 1 to generate an epitope-bearing  
polypeptide of the present invention. Therefore, the immunogenic epitopes of Table 1  
may include additional N-terminal or C-terminal amino acid residues. The additional  
flanking amino acid residues may be contiguous flanking N-terminal and/or C-  
terminal sequences from the polypeptides of the present invention, heterologous  
15 polypeptide sequences, or may include both contiguous flanking sequences from the  
polypeptides of the present invention and heterologous polypeptide sequences.

Polypeptides of the present invention comprising immunogenic or antigenic  
~~epitopes are at least 7 amino acids residues in length.~~ "At least" means that a  
polypeptide of the present invention comprising an immunogenic or antigenic epitope  
20 may be 7 amino acid residues in length or any integer between 7 amino acids and the  
number of amino acid residues of the full length polypeptides of the invention.  
Preferred polypeptides comprising immunogenic or antigenic epitopes are at least 10,  
15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 amino acid  
residues in length. However, it is pointed out that each and every integer between 7  
25 and the number of amino acid residues of the full length polypeptide are included in  
the present invention.

6  
The <sup>immunogenic</sup>~~immune~~ and antigenic epitope-bearing fragments may be specified by  
either the number of contiguous amino acid residues, as described above, or further  
specified by N-terminal and C-terminal positions of these fragments on the amino  
30 acid sequence of SEQ ID NO:2. Every combination of a N-terminal and C-terminal  
position that a fragment of, for example, at least 7 or at least 15 contiguous amino  
acid residues in length could occupy on the amino acid sequence of SEQ ID NO:2 is  
included in the invention. Again, "at least 7 contiguous amino acid residues in



length" means 7 amino acid residues in length or any integer between 7 amino acids and the number of amino acid residues of the full length polypeptide of the present invention. Specifically, each and every integer between 7 and the number of amino acid residues of the full length polypeptide are included in the present invention.

5 Immunogenic and antigenic epitope-bearing polypeptides of the invention are useful, for example, to make antibodies which specifically bind the polypeptides of the invention, and in immunoassays to detect the polypeptides of the present invention. The antibodies are useful, for example, in affinity purification of the polypeptides of the present invention. The antibodies may also routinely be used in  
10 a variety of qualitative or quantitative immunoassays, specifically for the polypeptides of the present invention using methods known in the art. See, e.g., Harlow et al., Antibodies: A Laboratory Manual, (Cold Spring Harbor Laboratory Press; 2nd Ed. 1988).

The epitope-bearing polypeptides of the present invention may be produced by  
15 any conventional means for making polypeptides including synthetic and recombinant methods known in the art. For instance, epitope-bearing peptides may be synthesized using known methods of chemical synthesis. For instance, Houghten has described a ~~simple method for the synthesis of large numbers of peptides, such as 10-20 mgs of~~  
248 individual and distinct 13 residue peptides representing single amino acid variants  
20 of a segment of the HA1 polypeptide, all of which were prepared and characterized (by ELISA-type binding studies) in less than four weeks (Houghten, R. A. Proc. Natl. Acad. Sci. USA 82:5131-5135 (1985)). This "Simultaneous Multiple Peptide Synthesis (SMPS)" process is further described in U.S. Patent No. 4,631,211 to Houghten and coworkers (1986). In this procedure the individual resins for the  
25 solid-phase synthesis of various peptides are contained in separate solvent-permeable packets, enabling the optimal use of the many identical repetitive steps involved in solid-phase methods. A completely manual procedure allows 500-1000 or more syntheses to be conducted simultaneously (Houghten et al. (1985) Proc. Natl. Acad. Sci. 82:5131-5135 at 5134.

30 Epitope-bearing polypeptides of the present invention are used to induce antibodies according to methods well known in the art including, but not limited to, in vivo immunization, in vitro immunization, and phage display methods. See, e.g., Sutcliffe, et al., supra; Wilson, et al., supra, and Bittle, et al. (1985) J. Gen. Virol.

668180-02492660

66:2347-2354. If in vivo immunization is used, animals may be immunized with free peptide; however, anti-peptide antibody titer may be boosted by coupling of the peptide to a macromolecular carrier, such as keyhole limpet hemacyanin (KLH) or tetanus toxoid. For instance, peptides containing cysteine residues may be coupled to  
5 a carrier using a linker such as -maleimidobenzoyl- N-hydroxysuccinimide ester (MBS), while other peptides may be coupled to carriers using a more general linking agent such as glutaraldehyde. Animals such as rabbits, rats and mice are immunized with either free or carrier-coupled peptides, for instance, by intraperitoneal and/or intradermal injection of emulsions containing about 100 µgs of peptide or carrier  
10 protein and Freund's adjuvant. Several booster injections may be needed, for instance, at intervals of about two weeks, to provide a useful titer of anti-peptide antibody which can be detected, for example, by ELISA assay using free peptide adsorbed to a solid surface. The titer of anti-peptide antibodies in serum from an immunized animal may be increased by selection of anti-peptide antibodies, for  
15 instance, by adsorption to the peptide on a solid support and elution of the selected antibodies according to methods well known in the art.

As one of skill in the art will appreciate, and discussed above, the polypeptides of the present invention comprising an immunogenic or antigenic epitope can be fused to heterologous polypeptide sequences. For example, the polypeptides of the  
20 present invention may be fused with the constant domain of immunoglobulins (IgA, IgE, IgG, IgM), or portions thereof (CH1, CH2, CH3, any combination thereof including both entire domains and portions thereof) resulting in chimeric polypeptides. These fusion proteins facilitate purification, and show an increased half-life in vivo. This has been shown, e.g., for chimeric proteins consisting of the  
25 first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. See, e.g., EPA 0,394,827; Traunecker et al. (1988) Nature 331:84-86. Fusion proteins that have a disulfide-linked dimeric structure due to the IgG portion can also be more efficient in binding and neutralizing other molecules than monomeric polypeptides or fragments  
30 thereof alone. See, e.g., Fountoulakis et al. (1995) J. Biochem. 270:3958-3964. Nucleic acids encoding the above epitopes can also be recombined with a gene of interest as an epitope tag to aid in detection and purification of the expressed polypeptide.

66:2347-2354

### Antibodies

The present invention further relates to antibodies and T-cell antigen receptors (TCR) which specifically bind the polypeptides of the present invention. The antibodies of the present invention include IgG (including IgG1, IgG2, IgG3, and IgG4), IgA (including IgA1 and IgA2), IgD, IgE, or IgM, and IgY. As used herein, the term "antibody" (Ab) is meant to include whole antibodies, including single-chain whole antibodies, and antigen-binding fragments thereof. Most preferably the antibodies are human antigen binding antibody fragments of the present invention include, but are not limited to, Fab, Fab' and F(ab')<sub>2</sub>, Fd, single-chain Fvs (scFv), single-chain antibodies, disulfide-linked Fvs (sdFv) and fragments comprising either a V<sub>L</sub> or V<sub>H</sub> domain. The antibodies may be from any animal origin including birds and mammals. Preferably, the antibodies are human, murine, rabbit, goat, guinea pig, camel, horse, or chicken.

Antigen-binding antibody fragments, including single-chain antibodies, may comprise the variable region(s) alone or in combination with the entire or partial of the following: hinge region, CH1, CH2, and CH3 domains. Also included in the invention are any combinations of variable region(s) and hinge region, CH1, CH2, and CH3 domains. The present invention further includes monoclonal, polyclonal, chimeric, humanized, and human monoclonal and polyclonal antibodies which specifically bind the polypeptides of the present invention. The present invention further includes antibodies which are anti-idiotypic to the antibodies of the present invention.

The antibodies of the present invention may be monospecific, bispecific, trispecific or of greater multispecificity. Multispecific antibodies may be specific for different epitopes of a polypeptide of the present invention or may be specific for both a polypeptide of the present invention as well as for heterologous compositions, such as a heterologous polypeptide or solid support material. See, e.g., WO 93/17715; WO 92/08802; WO 91/00360; WO 92/05793; Tutt, A. et al. (1991) J. Immunol. 147:60-69; US Patents 5,573,920, 4,474,893, 5,601,819, 4,714,681, 4,925,648; Kostelny, S.A. et al. (1992) J. Immunol. 148:1547-1553.

Antibodies of the present invention may be described or specified in terms of the epitope(s) or portion(s) of a polypeptide of the present invention which are

recognized or specifically bound by the antibody. The epitope(s) or polypeptide portion(s) may be specified as described herein, e.g., by N-terminal and C-terminal positions, by size in contiguous amino acid residues, or listed in the Tables and Figures. Antibodies which specifically bind any epitope or polypeptide of the present invention may also be excluded. Therefore, the present invention includes antibodies that specifically bind polypeptides of the present invention, and allows for the exclusion of the same.

Antibodies of the present invention may also be described or specified in terms of their cross-reactivity. Antibodies that do not bind any other analog, ortholog, or homolog of the polypeptides of the present invention are included. Antibodies that do not bind polypeptides with less than 95%, less than 90%, less than 85%, less than 80%, less than 75%, less than 70%, less than 65%, less than 60%, less than 55%, and less than 50% identity (as calculated using methods known in the art and described herein) to a polypeptide of the present invention are also included in the present invention. Further included in the present invention are antibodies which only bind polypeptides encoded by polynucleotides which hybridize to a polynucleotide of the present invention under stringent hybridization conditions (as described herein). Antibodies of the present invention may also be described or specified in terms of

their binding affinity. Preferred binding affinities include those with a dissociation constant or  $K_d$  less than  $5 \times 10^{-6}M$ ,  $10^{-6}M$ ,  $5 \times 10^{-7}M$ ,  $10^{-7}M$ ,  $5 \times 10^{-8}M$ ,  $10^{-8}M$ ,  $5 \times 10^{-9}M$ ,  $10^{-9}M$ ,  $5 \times 10^{-10}M$ ,  $10^{-10}M$ ,  $5 \times 10^{-11}M$ ,  $10^{-11}M$ ,  $5 \times 10^{-12}M$ ,  $10^{-12}M$ ,  $5 \times 10^{-13}M$ ,  $10^{-13}M$ ,  $5 \times 10^{-14}M$ ,  $10^{-14}M$ ,  $5 \times 10^{-15}M$ , and  $10^{-15}M$ .

Antibodies of the present invention have uses that include, but are not limited to, methods known in the art to purify, detect, and target the polypeptides of the present invention including both in vitro and in vivo diagnostic and therapeutic methods. For example, the antibodies have use in immunoassays for qualitatively and quantitatively measuring levels of the polypeptides of the present invention in biological samples. See, e.g., Harlow et al., ANTIBODIES: A LABORATORY MANUAL, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988) (incorporated by reference in the entirety).

The antibodies of the present invention may be used either alone or in combination with other compositions. The antibodies may further be recombinantly fused to a heterologous polypeptide at the N- or C-terminus or chemically conjugated

(including covalently and non-covalently conjugations) to polypeptides or other compositions. For example, antibodies of the present invention may be recombinantly fused or conjugated to molecules useful as labels in detection assays and effector molecules such as heterologous polypeptides, drugs, or toxins. See, e.g.,

5 WO 92/08495; WO 91/14438; WO 89/12624; US Patent 5,314,995; and EP 0 396 387.

The antibodies of the present invention may be prepared by any suitable method known in the art. For example, a polypeptide of the present invention or an antigenic fragment thereof can be administered to an animal in order to induce the

10 production of sera containing polyclonal antibodies. The term "monoclonal antibody" is not limited to antibodies produced through hybridoma technology. The term "monoclonal antibody" refers to an antibody that is derived from a single clone, including any eukaryotic, prokaryotic, or phage clone, and not the method by which it is produced. Monoclonal antibodies can be prepared using a wide variety of

15 techniques known in the art including the use of hybridoma, recombinant, and phage display technology.

Hybridoma techniques include those known in the art and taught in Harlow et al., ANTIBODIES: A LABORATORY MANUAL, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988); Hammerling, et al., in: MONOCLONAL ANTIBODIES AND

20 T-CELL HYBRIDOMAS 563-681 (Elsevier, N.Y., 1981) (said references incorporated by reference in their entireties). Fab and F(ab')<sub>2</sub> fragments may be produced by proteolytic cleavage, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')<sub>2</sub> fragments).

Alternatively, antibodies of the present invention can be produced through the

25 application of recombinant DNA and phage display technology or through synthetic chemistry using methods known in the art. For example, the antibodies of the present invention can be prepared using various phage display methods known in the art. In phage display methods, functional antibody domains are displayed on the surface of a phage particle which carries polynucleotide sequences encoding them. Phage with a

30 desired binding property are selected from a repertoire or combinatorial antibody library (e.g. human or murine) by selecting directly with antigen, typically antigen bound or captured to a solid surface or bead. Phage used in these methods are typically filamentous phage including fd and M13 with Fab, Fv or disulfide stabilized

0937430-0E492E60

Fv antibody domains recombinantly fused to either the phage gene III or gene VIII protein. Examples of phage display methods that can be used to make the antibodies of the present invention include those disclosed in Brinkman U. et al. (1995) J. Immunol. Methods 182:41-50; Ames, R.S. et al. (1995) J. Immunol. Methods 184:177-186; Kettleborough, C.A. et al. (1994) Eur. J. Immunol. 24:952-958; Persic, L. et al. (1997) Gene 187 9-18; Burton, D.R. et al. (1994) Advances in Immunology 57:191-280; PCT/GB91/01134; WO 90/02809; WO 91/10737; WO 92/01047; WO 92/18619; WO 93/11236; WO 95/15982; WO 95/20401; and US Patents 5,698,426, 5,223,409, 5,403,484, 5,580,717, 5,427,908, 5,750,753, 5,821,047, 5,571,698, 5,427,908, 5,516,637, 5,780,225, 5,658,727 and 5,733,743 (said references incorporated by reference in their entireties).

As described in the above references, after phage selection, the antibody coding regions from the phage can be isolated and used to generate whole antibodies, including human antibodies, or any other desired antigen binding fragment, and expressed in any desired host including mammalian cells, insect cells, plant cells, yeast, and bacteria. For example, techniques to recombinantly produce Fab, Fab' and F(ab')<sub>2</sub> fragments can also be employed using methods known in the art such as those disclosed in WO 92/22324; Mullinax, R.L. et al. (1992) BioTechniques 12(6):864-869; and Sawai, H. et al. (1995) AJRI 34:26-34; and Better, M. et al. (1988) Science 240:1041-1043 (said references incorporated by reference in their entireties).

Examples of techniques which can be used to produce single-chain Fvs and antibodies include those described in U.S. Patents 4,946,778 and 5,258,498; Huston et al. (1991) Methods in Enzymology 203:46-88; Shu, L. et al. (1993) PNAS 90:7995-7999; and Skerra, A. et al. (1988) Science 240:1038-1040. For some uses, including in vivo use of antibodies in humans and in vitro detection assays, it may be preferable to use chimeric, humanized, or human antibodies. Methods for producing chimeric antibodies are known in the art. See e.g., Morrison, Science 229:1202 (1985); Oi et al., BioTechniques 4:214 (1986); Gillies, S.D. et al. (1989) J. Immunol. Methods 125:191-202; and US Patent 5,807,715. Antibodies can be humanized using a variety of techniques including CDR-grafting (EP 0 239 400; WO 91/09967; US Patent 5,530,101; and 5,585,089), veneering or resurfacing (EP 0 592 106; EP 0 519 596; Padlan E.A., (1991) Molecular Immunology 28(4/5):489-498; Studnicka G.M. et al. (1994) Protein Engineering 7(6):805-814; Roguska M.A. et al. (1994) PNAS 91:969-

973), and chain shuffling (US Patent 5,565,332). Human antibodies can be made by a variety of methods known in the art including phage display methods described above. See also, US Patents 4,444,887, 4,716,111, 5,545,806, and 5,814,318; and WO 98/46645, WO 98/50433, WO 98/24893, WO 98/16654, WO 96/34096, WO 96/33735, and WO 91/10741 (said references incorporated by reference in their entireties).

Further included in the present invention are antibodies recombinantly fused or chemically conjugated (including both covalently and non-covalently conjugations) to a polypeptide of the present invention. The antibodies may be specific for antigens other than polypeptides of the present invention. For example, antibodies may be used to target the polypeptides of the present invention to particular cell types, either in vitro or in vivo, by fusing or conjugating the polypeptides of the present invention to antibodies specific for particular cell surface receptors. Antibodies fused or conjugated to the polypeptides of the present invention may also be used in in vitro immunoassays and purification methods using methods known in the art. See e.g., Harbor et al. supra and WO 93/21232; EP 0 439 095; Naramura, M. et al. (1994) Immunol. Lett. 39:91-99; US Patent 5,474,981; Gillies, S.O. et al. (1992) PNAS 89:1428-1432; Fell, H.P. et al. (1991) J. Immunol. 146:2446-2452 (said references incorporated by reference in their entireties).

The present invention further includes compositions comprising the polypeptides of the present invention fused or conjugated to antibody domains other than the variable regions. For example, the polypeptides of the present invention may be fused or conjugated to an antibody Fc region, or portion thereof. The antibody portion fused to a polypeptide of the present invention may comprise the hinge region, CH1 domain, CH2 domain, and CH3 domain or any combination of whole domains or portions thereof. The polypeptides of the present invention may be fused or conjugated to the above antibody portions to increase the in vivo half life of the polypeptides or for use in immunoassays using methods known in the art. The polypeptides may also be fused or conjugated to the above antibody portions to form multimers. For example, Fc portions fused to the polypeptides of the present invention can form dimers through disulfide bonding between the Fc portions. Higher multimeric forms can be made by fusing the polypeptides to portions of IgA and IgM. Methods for fusing or conjugating the polypeptides of the present invention to

antibody portions are known in the art. See e.g., US Patents 5,336,603, 5,622,929, 5,359,046, 5,349,053, 5,447,851, 5,112,946; EP 0 307 434, EP 0 367 166; WO 96/04388, WO 91/06570; Ashkenazi, A. et al. (1991) PNAS 88:10535-10539; Zheng, X.X. et al. (1995) J. Immunol. 154:5590-5600; and Vil, H. et al. (1992) PNAS 89:11337-11341 (said references incorporated by reference in their entireties).

The invention further relates to antibodies that act as agonists or antagonists of the polypeptides of the present invention. Antibodies which act as agonists or antagonists of the polypeptides of the present invention include, for example, antibodies which disrupt receptor/ligand interactions with the polypeptides of the invention either partially or fully. For example, the present invention includes antibodies that disrupt the ability of the proteins of the invention to multimerize. In another example, the present invention includes antibodies which allow the proteins of the invention to multimerize, but disrupts the ability of the proteins of the invention to bind one or more CRCGCL receptor(s)/ligand(s). In yet another example, the present invention includes antibodies which allow the proteins of the invention to multimerize, and bind CRCGCL receptor(s)/ligand(s), but blocks biological activity associated with the CRCGCL/receptor/ligand complex.

~~Antibodies which act as agonists or antagonists of the polypeptides of the~~  
present invention also include, both receptor-specific antibodies and ligand-specific antibodies. Included are receptor-specific antibodies that do not prevent ligand binding but prevent receptor activation. Receptor activation (i.e., signaling) may be determined by techniques described herein or otherwise known in the art. Also included are receptor-specific antibodies which both prevent ligand binding and receptor activation. Likewise, included are neutralizing antibodies which bind the ligand and prevent binding of the ligand to the receptor, as well as antibodies which bind the ligand, thereby preventing receptor activation, but do not prevent the ligand from binding the receptor. Further included are antibodies that activate the receptor. These antibodies may act as agonists for either all or less than all of the biological activities affected by ligand-mediated receptor activation. The antibodies may be specified as agonists or antagonists for biological activities comprising specific activities disclosed herein. The above antibody agonists can be made using methods known in the art. See e.g., WO 96/40281; US Patent Number 5,811,097; Deng, B. et al., Blood 92(6):1981-1988 (1998); Chen, Z. et al., Cancer Res. 58(16):3668-3678



(1998); Harrop, J.A. et al., J. Immunol. 161(4):1786-1794 (1998); Zhu, Z. et al., Cancer Res. 58(15):3209-3214 (1998); Yoon, D.Y. et al., J. Immunol. 160(7):3170-3179 (1998); Prat, M. et al., J. Cell. Sci. 111(Pt2):237-247 (1998); Pitard, V. et al., J. Immunol. Methods 205(2):177-190 (1997); Liautard, J. et al., Cytokine 9(4):233-241 (1997); Carlson, N.G. et al., J. Biol. Chem. 272(17):11295-11301 (1997); Taryman, R.E. et al., Neuron 14(4):755-762 (1995); Muller, Y.A. et al., Structure 6(9):1153-1167 (1998); Bartunek, P. et al., Cytokine 8(1):14-20 (1996)(said references incorporated by reference in their entireties).

As discussed above, antibodies to the CRCGCL proteins of the invention can, in turn, be utilized to generate anti-idiotypic antibodies that "mimic" CRCGCL using techniques well known to those skilled in the art. (See, e.g., Greenspan & Bona, FASEB J. 7(5):437-444; (1989) and Nissinoff, J. Immunol. 147(8):2429-2438 (1991)). For example, antibodies which bind to CRCGCL and competitively inhibit CRCGCL multimerization and/or binding to ligand can be used to generate anti-idiotypes that "mimic" the CRCGCL multimerization and/or binding domain and, as a consequence, bind to and neutralize CRCGCL and/or its ligand. Such neutralizing anti-idiotypes or Fab fragments of such anti-idiotypes can be used in therapeutic regimens to neutralize CRCGCL ligand. For example, such anti-idiotypic antibodies can be used to bind CRCGCL, or to bind CRCGCL ligands/receptors, and thereby block CRCGCL biological activity.

### **Fusion Proteins**

Any CRCGCL polypeptide can be used to generate fusion proteins. For example, the CRCGCL polypeptide, when fused to a second protein, can be used as an antigenic tag. Antibodies raised against the CRCGCL polypeptide can be used to indirectly detect the second protein by binding to the CRCGCL. Moreover, because secreted proteins target cellular locations based on trafficking signals, the CRCGCL polypeptides can be used as a targeting molecule once fused to other proteins.

Examples of domains that can be fused to CRCGCL polypeptides include not only heterologous signal sequences, but also other heterologous functional regions. The fusion does not necessarily need to be direct, but may occur through linker sequences.

In certain preferred embodiments, CRCGCL proteins of the invention

comprise fusion proteins wherein the CRCGCL polypeptides are those described above as m-n. In preferred embodiments, the application is directed to nucleic acid molecules at least 90%, 95%, 96%, 97%, 98% or 99% identical to the nucleic acid sequences encoding polypeptides having the amino acid sequence of the specific N- and C-terminal deletions recited herein. Polynucleotides encoding these polypeptides are also encompassed by the invention.

Moreover, fusion proteins may also be engineered to improve characteristics of the CRCGCL polypeptide. For instance, a region of additional amino acids, particularly charged amino acids, may be added to the N-terminus of the CRCGCL polypeptide to improve stability and persistence during purification from the host cell or subsequent handling and storage. Also, peptide moieties may be added to the CRCGCL polypeptide to facilitate purification. Such regions may be removed prior to final preparation of the CRCGCL polypeptide. The addition of peptide moieties to facilitate handling of polypeptides are familiar and routine techniques in the art.

Moreover, CRCGCL polypeptides, including fragments, and specifically epitopes, can be combined with parts of the constant domain of immunoglobulins (IgG), resulting in chimeric polypeptides. These fusion proteins facilitate purification and show an increased half-life in vivo. One reported example describes chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. (EP A 394,827; Traunecker et al., Nature 331:84-86 (1988).) Fusion proteins having disulfide-linked dimeric structures (due to the IgG) can also be more efficient in binding and neutralizing other molecules, than the monomeric secreted protein or protein fragment alone. (Fountoulakis et al., J. Biochem. 270:3958-3964 (1995).)

Similarly, EP-A-O 464 533 (Canadian counterpart 2045869) discloses fusion proteins comprising various portions of constant region of immunoglobulin molecules together with another human protein or part thereof. In many cases, the Fc part in a fusion protein is beneficial in therapy and diagnosis, and thus can result in, for example, improved pharmacokinetic properties. (EP-A 0232 262.) Alternatively, deleting the Fc part after the fusion protein has been expressed, detected, and purified, would be desired. For example, the Fc portion may hinder therapy and diagnosis if the fusion protein is used as an antigen for immunizations. In drug discovery, for

Moreover, the CRCGCL polypeptides can be fused to marker sequences, such as a peptide which facilitates purification of CRCGCL. In preferred embodiments, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311), among others, many of which are commercially available. As described in Gentz et al., Proc. Natl. Acad. Sci. USA 86:821-824 (1989), for instance, hexa-histidine provides for convenient purification of the fusion protein. Another peptide tag useful for purification, the "HA" tag, corresponds to an epitope derived from the influenza hemagglutinin protein. (Wilson et al., Cell 37:767 (1984).)

## Vectors, Host Cells, and Protein Production

CRGCL polynucleotides may be joined to a vector containing a selectable  
25 marker for propagation in a host. Generally, a plasmid vector is introduced in a  
precipitate, such as a calcium phosphate precipitate, or in a complex with a charged  
lipid. If the vector is a virus, it may be packaged in vitro using an appropriate  
packaging cell line and then transduced into host cells.

The CRCGCL polynucleotide insert should be operatively linked to an appropriate promoter, such as the phage lambda PL promoter, the E. coli lac, trp, phoA and tac promoters, the SV40 early and late promoters and promoters of retroviral LTRs, to name a few. Other suitable promoters will be known to the skilled artisan. The expression constructs will further contain sites for transcription

initiation, termination, and, in the transcribed region, a ribosome binding site for translation. The coding portion of the transcripts expressed by the constructs will preferably include a translation initiating codon at the beginning and a termination codon (UAA, UGA or UAG) appropriately positioned at the end of the polypeptide to be translated.

As indicated, the expression vectors will preferably include at least one selectable marker. Such markers include dihydrofolate reductase, G418 or neomycin resistance for eukaryotic cell culture and tetracycline, kanamycin or ampicillin resistance genes for culturing in *E. coli* and other bacteria. Representative examples of appropriate hosts include, but are not limited to, bacterial cells, such as *E. coli*, *Streptomyces* and *Salmonella typhimurium* cells; fungal cells, such as yeast cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, 293, and Bowes melanoma cells; and plant cells. Appropriate culture mediums and conditions for the above-described host cells are known in the art.

Among vectors preferred for use in bacteria include pQE70, pQE60 and pQE-9, available from QIAGEN, Inc.; pBluescript vectors, Phagescript vectors, pNH8A, pNH16a, pNH18A, pNH46A, available from Stratagene Cloning Systems, Inc.; and ptre99a, pKK223-3, pKK233-3, pDR540, pRIT5 available from Pharmacia Biotech, Inc. Among preferred eukaryotic vectors are pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; and pSVK3, pBPV, pMSG and pSVL available from Pharmacia. Other suitable vectors will be readily apparent to the skilled artisan.

Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection, or other methods. Such methods are described in many standard laboratory manuals, such as Davis et al., Basic Methods In Molecular Biology (1986). It is specifically contemplated that CRCGCL polypeptides may in fact be expressed by a host cell lacking a recombinant vector.

CRCGCL polypeptides can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most

preferably, high performance liquid chromatography ("HPLC") is employed for purification.

CRCGCL polypeptides, and preferably the secreted form, can also be recovered from: products purified from natural sources, including bodily fluids, tissues and cells, whether directly isolated or cultured; products of chemical synthetic procedures; and products produced by recombinant techniques from a prokaryotic or eukaryotic host, including, for example, bacterial, yeast, higher plant, insect, and mammalian cells. Depending upon the host employed in a recombinant production procedure, the CRCGCL polypeptides may be glycosylated or may be non-glycosylated. In addition, CRCGCL polypeptides may also include an initial modified methionine residue, in some cases as a result of host-mediated processes. Thus, it is well known in the art that the N-terminal methionine encoded by the translation initiation codon generally is removed with high efficiency from any protein after translation in all eukaryotic cells. While the N-terminal methionine on most proteins also is efficiently removed in most prokaryotes, for some proteins, this prokaryotic removal process is inefficient, depending on the nature of the amino acid to which the N-terminal methionine is covalently linked.

In addition to encompassing host cells containing the vector constructs discussed herein, the invention also encompasses primary, secondary, and immortalized host cells of vertebrate origin, particularly mammalian origin, that have been engineered to delete or replace endogenous genetic material (e.g., CRCGCL coding sequence), and/or to include genetic material (e.g., heterologous polynucleotide sequences) that is operably associated with CRCGCL polynucleotides of the invention, and which activates, alters, and/or amplifies endogenous CRCGCL polynucleotides. For example, techniques known in the art may be used to operably associate heterologous control regions (e.g., promoter and/or enhancer) and endogenous CRCGCL polynucleotide sequences via homologous recombination (see, e.g., U.S. Patent No. 5,641,670, issued June 24, 1997; International Publication No. WO 96/29411, published September 26, 1996; International Publication No. WO 94/12650, published August 4, 1994; Koller et al., Proc. Natl. Acad. Sci. USA 86:8932-8935 (1989); and Zijlstra et al., Nature 342:435-438 (1989), the disclosures of each of which are incorporated by reference in their entireties).

In addition, polypeptides of the invention can be chemically synthesized using

5  
10  
15

20

20

25

30

Also provided by the invention are chemically modified derivatives of CRCGCL which may provide additional advantages such as increased solubility,

5 The polypeptides may be modified at random positions within the molecule, or at predetermined positions within the molecule and may include one, two, three or more attached chemical moieties.

The polymer may be of any molecular weight, and may be branched or unbranched. For polyethylene glycol, the preferred molecular weight is between about 1 kDa and about 100 kDa (the term "about" indicating that in preparations of polyethylene glycol, some molecules will weigh more, some less, than the stated molecular weight) for ease in handling and manufacturing. Other sizes may be used, depending on the desired therapeutic profile (e.g., the duration of sustained release desired, the effects, if any on biological activity, the ease in handling, the degree or lack of antigenicity and other known effects of the polyethylene glycol to a therapeutic protein or analog).

The polyethylene glycol molecules (or other chemical moieties) should be attached to the protein with consideration of effects on functional or antigenic domains of the protein. There are a number of attachment methods available to those skilled in the art, e.g., EP 0 401 384, herein incorporated by reference (coupling PEG to G-CSF), see also Malik et al., Exp. Hematol. 20:1028-1035 (1992) (reporting pegylation of GM-CSF using tresyl chloride). For example, polyethylene glycol may be covalently bound through amino acid residues via a reactive group, such as, a free amino or carboxyl group. Reactive groups are those to which an activated polyethylene glycol molecule may be bound. The amino acid residues having a free amino group may include lysine residues and the N-terminal amino acid residues; those having a free carboxyl group may include aspartic acid residues glutamic acid residues and the C-terminal amino acid residue. Sulfhydryl groups may also be used as a reactive group for attaching the polyethylene glycol molecules. Preferred for therapeutic purposes is attachment at an amino group, such as attachment at the N-terminus or lysine group.

One may specifically desire proteins chemically modified at the N-terminus. Using polyethylene glycol as an illustration of the present composition, one may

select from a variety of polyethylene glycol molecules (by molecular weight, branching, etc.), the proportion of polyethylene glycol molecules to protein (or peptide) molecules in the reaction mix, the type of pegylation reaction to be performed, and the method of obtaining the selected N-terminally pegylated protein.

- 5 The method of obtaining the N-terminally pegylated preparation (i.e., separating this moiety from other monopegylated moieties if necessary) may be by purification of the N-terminally pegylated material from a population of pegylated protein molecules. Selective proteins chemically modified at the N-terminus modification may be accomplished by reductive alkylation which exploits differential reactivity of different
- 10 types of primary amino groups (lysine versus the N-terminal) available for derivatization in a particular protein. Under the appropriate reaction conditions, substantially selective derivatization of the protein at the N-terminus with a carbonyl group containing polymer is achieved.

- The CRCGCL polypeptides of the invention may be in monomers or
- 15 multimers (i.e., dimers, trimers, tetramers and higher multimers). Accordingly, the present invention relates to monomers and multimers of the CRCGCL polypeptides of the invention, their preparation, and compositions (preferably, pharmaceutical compositions) containing them. In specific embodiments, the polypeptides of the invention are monomers, dimers, trimers or tetramers. In additional embodiments, the
- 20 multimers of the invention are at least dimers, at least trimers, or at least tetramers.

- Multimers encompassed by the invention may be homomers or heteromers. As used herein, the term homomer, refers to a multimer containing only CRCGCL polypeptides of the invention (including CRCGCL fragments, variants, splice variants, and fusion proteins, as described herein). These homomers may contain
- 25 CRCGCL polypeptides having identical or different amino acid sequences. In a specific embodiment, a homomer of the invention is a multimer containing only CRCGCL polypeptides having an identical amino acid sequence. In another specific embodiment, a homomer of the invention is a multimer containing CRCGCL polypeptides having different amino acid sequences. In specific embodiments, the
- 30 multimer of the invention is a homodimer (e.g., containing CRCGCL polypeptides having identical or different amino acid sequences) or a homotrimer (e.g., containing CRCGCL polypeptides having identical and/or different amino acid sequences). In

093649-0819260



additional embodiments, the homomeric multimer of the invention is at least a homodimer, at least a homotrimer, or at least a homotetramer.

As used herein, the term heteromer refers to a multimer containing one or more heterologous polypeptides (i.e., polypeptides of different proteins) in addition to the CRCGCL polypeptides of the invention. In a specific embodiment, the multimer of the invention is a heterodimer, a heterotrimer, or a heterotetramer. In additional embodiments, the homomeric multimer of the invention is at least a homodimer, at least a homotrimer, or at least a homotetramer.

Multimers of the invention may be the result of hydrophobic, hydrophilic, ionic and/or covalent associations and/or may be indirectly linked, by for example, liposome formation. Thus, in one embodiment, multimers of the invention, such as, for example, homodimers or homotrimers, are formed when polypeptides of the invention contact one another in solution. In another embodiment, heteromultimers of the invention, such as, for example, heterotrimers or heterotetramers, are formed when polypeptides of the invention contact antibodies to the polypeptides of the invention (including antibodies to the heterologous polypeptide sequence in a fusion protein of the invention) in solution. In other embodiments, multimers of the invention are formed by covalent associations with and/or between the CRCGCL polypeptides of the invention. Such covalent associations may involve one or more amino acid residues contained in the polypeptide sequence ( e.g., that recited in SEQ ID NO:2, or contained in the polypeptide encoded by the clone HTAEK53). In one instance, the covalent associations are cross-linking between cysteine residues located within the polypeptide sequences which interact in the native (i.e., naturally occurring) polypeptide. In another instance, the covalent associations are the consequence of chemical or recombinant manipulation. Alternatively, such covalent associations may involve one or more amino acid residues contained in the heterologous polypeptide sequence in a CRCGCL fusion protein. In one example, covalent associations are between the heterologous sequence contained in a fusion protein of the invention (see, e.g., US Patent Number 5,478,925). In a specific example, the covalent associations are between the heterologous sequence contained in a CRCGCL-Fc fusion protein of the invention (as described herein). In another specific example, covalent associations of fusion proteins of the invention are between heterologous polypeptide sequence from another Cytokine Receptor family



herein or otherwise known in the art are applied to generate recombinant polypeptides of the invention which contain a transmembrane domain (or hydrophobic or signal peptide) and which can be incorporated by membrane reconstitution techniques into liposomes (see, e.g., US Patent Number 5,478,925, which is herein incorporated by  
5 reference in its entirety).

#### Uses of the CRCGCL Polynucleotides

The CRCGCL polynucleotides identified herein can be used in numerous ways as reagents. The following description should be considered exemplary and  
10 utilizes known techniques.

There exists an ongoing need to identify new chromosome markers, since few chromosome marking reagents, based on actual sequence data (repeat polymorphisms), are presently available. Using a panel of radiation hybrids, CRCGCL maps to the pseudoautosomal region (PAR) of the sex chromosomes,  
15 which is located on both X (Xp22.3) and Y (Yp13.3). Interestingly, two other cytokine receptors map to this region (IL3Ra, and GMCSFRa). See, Kremer et al. "A Cytokine Receptor Gene Cluster in the X-Y pseudoautosomal region ?" Blood 82(1) 22-28 (1993). Thus, CRCGCL polynucleotides can be used in linkage analysis as a  
marker for the pseudoautosomal region on the X and Y chromosomes.

Briefly, sequences can be mapped to chromosomes by preparing PCR primers  
20 (preferably 15-25 bp) from the sequences shown in SEQ ID NO:1. Primers can be selected using computer analysis so that primers do not span more than one predicted exon in the genomic DNA. These primers are then used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids  
25 containing the human CRCGCL gene corresponding to the SEQ ID NO:1 will yield an amplified fragment.

Similarly, somatic hybrids provide a rapid method of PCR mapping the polynucleotides to particular chromosomes. Three or more clones can be assigned per day using a single thermal cycler. Moreover, sublocalization of the CRCGCL  
30 polynucleotides can be achieved with panels of specific chromosome fragments. Other gene mapping strategies that can be used include in situ hybridization, prescreening with labeled flow-sorted chromosomes, and preselection by hybridization to construct chromosome specific-cDNA libraries.

668180-0E192E60

Precise chromosomal location of the CRCGCL polynucleotides can also be achieved using fluorescence in situ hybridization (FISH) of a metaphase chromosomal spread. This technique uses polynucleotides as short as 500 or 600 bases; however, polynucleotides 2,000-4,000 bp are preferred. For a review of this technique, see Verma et al., "Human Chromosomes: a Manual of Basic Techniques," Pergamon Press, New York (1988).

For chromosome mapping, the CRCGCL polynucleotides can be used individually (to mark a single chromosome or a single site on that chromosome) or in panels (for marking multiple sites and/or multiple chromosomes). Preferred polynucleotides correspond to the noncoding regions of the cDNAs because the coding sequences are more likely conserved within gene families, thus increasing the chance of cross hybridization during chromosomal mapping.

Once a polynucleotide has been mapped to a precise chromosomal location, the physical position of the polynucleotide can be used in linkage analysis. Linkage analysis establishes coinheritance between a chromosomal location and presentation of a particular disease. (Disease mapping data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available on line through Johns Hopkins University Welch Medical Library) .) Assuming 1 megabase mapping resolution and one gene per 20 kb, a cDNA precisely localized to a chromosomal region associated with the disease could be one of 50-500 potential causative genes.

Thus, once coinheritance is established, differences in the CRCGCL polynucleotide and the corresponding gene between affected and unaffected individuals can be examined. First, visible structural alterations in the chromosomes, such as deletions or translocations, are examined in chromosome spreads or by PCR. If no structural alterations exist, the presence of point mutations are ascertained. Mutations observed in some or all affected individuals, but not in normal individuals, indicates that the mutation may cause the disease. However, complete sequencing of the CRCGCL polypeptide and the corresponding gene from several normal individuals is required to distinguish the mutation from a polymorphism. If a new polymorphism is identified, this polymorphic polypeptide can be used for further linkage analysis.

Furthermore, increased or decreased expression of the gene in affected individuals as compared to unaffected individuals can be assessed using CRCGCL

polynucleotides. Any of these alterations (altered expression, chromosomal rearrangement, or mutation) can be used as a diagnostic or prognostic marker.

5 In addition to the foregoing, a CRCGCL polynucleotide can be used to control gene expression through triple helix formation or antisense DNA or RNA. Both methods rely on binding of the polynucleotide to DNA or RNA. For these techniques, preferred polynucleotides are usually 20 to 40 bases in length and complementary to either the region of the gene involved in transcription (triple helix - see Lee et al., Nucl. Acids Res. 6:3073 (1979); Cooney et al., Science 241:456 (1988); and Dervan et al., Science 251:1360 (1991) ) or to the mRNA itself (antisense - Okano, J. Neurochem. 56:560 (1991); Oligodeoxy-nucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988).) Triple helix formation optimally results in a shut-off of RNA transcription from DNA, while antisense RNA hybridization blocks translation of an mRNA molecule into polypeptide. Both techniques are effective in model systems, and the information disclosed herein can be used to design antisense or triple helix polynucleotides in an effort to treat disease.

CRCGCL polynucleotides are also useful in gene therapy. One goal of gene therapy is to insert a normal gene into an organism having a defective gene, in an effort to correct the genetic defect. CRCGCL offers a means of targeting such genetic defects in a highly accurate manner. Another goal is to insert a new gene that was not present in the host genome, thereby producing a new trait in the host cell.

The CRCGCL polynucleotides are also useful for identifying individuals from minute biological samples. The United States military, for example, is considering the use of restriction fragment length polymorphism (RFLP) for identification of its personnel. In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, and probed on a Southern blot to yield unique bands for identifying personnel. This method does not suffer from the current limitations of "Dog Tags" which can be lost, switched, or stolen, making positive identification difficult. The CRCGCL polynucleotides can be used as additional DNA markers for RFLP.

The CRCGCL polynucleotides can also be used as an alternative to RFLP, by determining the actual base-by-base DNA sequence of selected portions of an individual's genome. These sequences can be used to prepare PCR primers for

5

10

20

25

expression level in healthy tissue from an individual not having the immune system disorder.

Thus, the invention provides a diagnostic method of a disorder, which involves: (a) assaying CRCGCL gene expression level in cells or body fluid of an individual; (b) comparing the CRCGCL gene expression level with a standard CRCGCL gene expression level, whereby an increase or decrease in the assayed CRCGCL gene expression level compared to the standard expression level is indicative of disorder in the immune system.

In the very least, the CRCGCL polynucleotides can be used as molecular weight markers on Southern gels, as diagnostic probes for the presence of a specific mRNA in a particular cell type, as a probe to "subtract-out" known sequences in the process of discovering novel polynucleotides, for selecting and making oligomers for attachment to a "gene chip" or other support, to raise anti-DNA antibodies using DNA immunization techniques, and as an antigen to elicit an immune response.

#### Uses of CRCGCL Polypeptides

CRCGCL polypeptides can be used in numerous ways. The following description should be considered exemplary and utilizes known techniques.

CRCGCL polypeptides can be used to assay protein levels in a biological sample using antibody-based techniques. For example, protein expression in tissues can be studied with classical immunohistological methods. (Jalkanen, M., et al., J. Cell. Biol. 101:976-985 (1985); Jalkanen, M., et al., J. Cell . Biol. 105:3087-3096 (1987).) Other antibody-based methods useful for detecting protein gene expression include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA). Suitable antibody assay labels are known in the art and include enzyme labels, such as, glucose oxidase, and radioisotopes, such as iodine (125I, 121I), carbon (14C), sulfur (35S), tritium (3H), indium (112In), and technetium (99mTc), and fluorescent labels, such as fluorescein and rhodamine, and biotin.

In addition to assaying secreted protein levels in a biological sample, proteins can also be detected in vivo by imaging. Antibody labels or markers for in vivo imaging of protein include those detectable by X-radiography, NMR or ESR. For X-radiography, suitable labels include radioisotopes such as barium or cesium, which emit detectable radiation but are not overtly harmful to the subject. Suitable markers

for NMR and ESR include those with a detectable characteristic spin, such as deuterium, which may be incorporated into the antibody by labeling of nutrients for the relevant hybridoma.

5 A protein-specific antibody or antibody fragment which has been labeled with an appropriate detectable imaging moiety, such as a radioisotope (for example,  $^{131}\text{I}$ ,  $^{112}\text{In}$ ,  $^{99\text{m}}\text{Tc}$ ), a radio-opaque substance, or a material detectable by nuclear magnetic resonance, is introduced (for example, parenterally, subcutaneously, or intraperitoneally) into the mammal. It will be understood in the art that the size of the subject and the imaging system used will determine the quantity of imaging moiety  
10 needed to produce diagnostic images. In the case of a radioisotope moiety, for a human subject, the quantity of radioactivity injected will normally range from about 5 to 20 millicuries of  $^{99\text{m}}\text{Tc}$ . The labeled antibody or antibody fragment will then preferentially accumulate at the location of cells which contain the specific protein. In vivo tumor imaging is described in S.W. Burchiel et al., "Immunopharmacokinetics of Radiolabeled Antibodies and Their Fragments." (Chapter 13 in Tumor Imaging:  
15 The Radiochemical Detection of Cancer, S.W. Burchiel and B. A. Rhodes, eds., Masson Publishing Inc. (1982).)

Thus, the invention provides a diagnostic method of a disorder, which involves (a) assaying the expression of CRCGCL polypeptide in cells or body fluid of  
20 an individual; (b) comparing the level of gene expression with a standard gene expression level, whereby an increase or decrease in the assayed CRCGCL polypeptide gene expression level compared to the standard expression level is indicative of a disorder.

Moreover, CRCGCL polypeptides can be used to treat disease. For example,  
25 patients can be administered CRCGCL polypeptides in an effort to replace absent or decreased levels of the CRCGCL polypeptide (e.g., insulin), to supplement absent or decreased levels of a different polypeptide (e.g., hemoglobin S for hemoglobin B), to inhibit the activity of a polypeptide (e.g., an oncogene), to activate the activity of a polypeptide (e.g., by binding to a receptor), to reduce the activity of a membrane  
30 bound receptor by competing with it for free ligand (e.g., soluble TNF receptors used in reducing inflammation), or to bring about a desired response (e.g., blood vessel growth).



Similarly, antibodies directed to CRCGCL polypeptides can also be used to treat disease. For example, administration of an antibody directed to a CRCGCL polypeptide can bind and reduce overproduction of the polypeptide. Similarly, administration of an antibody can activate the polypeptide, such as by binding to a polypeptide bound to a membrane (receptor).

At the very least, the CRCGCL polypeptides can be used as molecular weight markers on SDS-PAGE gels or on molecular sieve gel filtration columns using methods well known to those of skill in the art. CRCGCL polypeptides can also be used to raise antibodies, which in turn are used to measure protein expression from a recombinant cell, as a way of assessing transformation of the host cell. Moreover, CRCGCL polypeptides can be used to test the following biological activities.

### **Gene Therapy Methods**

Another aspect of the present invention is to gene therapy methods for treating disorders, diseases and conditions. The gene therapy methods relate to the introduction of nucleic acid (DNA, RNA and antisense DNA or RNA) sequences into an animal to achieve expression of the CRCGCL polypeptide of the present invention. ~~This method requires a polynucleotide that codes for a CRCGCL polypeptide~~ operatively linked to a promoter and any other genetic elements necessary for the expression of the polypeptide by the target tissue. Such gene therapy and delivery techniques are known in the art, see, for example, WO90/11092, which is herein incorporated by reference.

Thus, for example, cells from a patient may be engineered with a polynucleotide (DNA or RNA) comprising a promoter operably linked to a CRCGCL polynucleotide ex vivo, with the engineered cells then being provided to a patient to be treated with the polypeptide. Such methods are well-known in the art. For example, see Belldgrun, A., et al., J. Natl. Cancer Inst. 85: 207-216 (1993); Ferrantini, M. et al., Cancer Research 53: 1107-1112 (1993); Ferrantini, M. et al., J. Immunology 153: 4604-4615 (1994); Kaido, T., et al., Int. J. Cancer 60: 221-229 (1995); Ogura, H., et al., Cancer Research 50: 5102-5106 (1990); Santodonato, L., et al., Human Gene Therapy 7:1-10 (1996); Santodonato, L., et al., Gene Therapy 4:1246-1255 (1997); and Zhang, J.-F. et al., Cancer Gene Therapy 3: 31-38 (1996)), which are herein incorporated by reference. In one embodiment, the cells that are

engineered are arterial cells. The arterial cells may be reintroduced into the patient through direct injection to the artery, the tissues surrounding the artery, or through catheter injection.

As discussed in more detail below, the CRCGCL polynucleotide constructs  
5 can be delivered by any method that delivers injectable materials to the cells of an animal, such as, injection into the interstitial space of tissues (heart, muscle, skin, lung, liver, and the like). The CRCGCL polynucleotide constructs may be delivered in a pharmaceutically acceptable liquid or aqueous carrier.

In one embodiment, the CRCGCL polynucleotide is delivered as a naked  
10 polynucleotide. The term "naked" polynucleotide, DNA or RNA refers to sequences that are free from any delivery vehicle that acts to assist, promote or facilitate entry into the cell, including viral sequences, viral particles, liposome formulations, lipofectin or precipitating agents and the like. However, the CRCGCL polynucleotides can also be delivered in liposome formulations and lipofectin  
15 formulations and the like can be prepared by methods well known to those skilled in the art. Such methods are described, for example, in U.S. Patent Nos. 5,593,972, 5,589,466, and 5,580,859, which are herein incorporated by reference.

~~The CRCGCL polynucleotide vector constructs used in the gene therapy~~  
method are preferably constructs that will not integrate into the host genome nor will  
20 they contain sequences that allow for replication. Appropriate vectors include pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; pSVK3, pBPV, pMSG and pSVL available from Pharmacia; and pEF1/V5, pcDNA3.1, and pRc/CMV2 available from Invitrogen. Other suitable vectors will be readily apparent to the skilled artisan.

Any strong promoter known to those skilled in the art can be used for driving  
25 the expression of CRCGCL DNA. Suitable promoters include adenoviral promoters, such as the adenoviral major late promoter; or heterologous promoters, such as the cytomegalovirus (CMV) promoter; the respiratory syncytial virus (RSV) promoter; inducible promoters, such as the MMT promoter, the metallothionein promoter; heat  
30 shock promoters; the albumin promoter; the ApoAI promoter; human globin promoters; viral thymidine kinase promoters, such as the Herpes Simplex thymidine kinase promoter; retroviral LTRs; the b-actin promoter; and human growth hormone promoters. The promoter also may be the native promoter for CRCGCL.

Unlike other gene therapy techniques, one major advantage of introducing naked nucleic acid sequences into target cells is the transitory nature of the polynucleotide synthesis in the cells. Studies have shown that non-replicating DNA sequences can be introduced into cells to provide production of the desired polypeptide for periods of up to six months.

The CRCGCL polynucleotide construct can be delivered to the interstitial space of tissues within the an animal, including of muscle, skin, brain, lung, liver, spleen, bone marrow, thymus, heart, lymph, blood, bone, cartilage, pancreas, kidney, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland, and connective tissue. Interstitial space of the tissues comprises the intercellular, fluid, mucopolysaccharide matrix among the reticular fibers of organ tissues, elastic fibers in the walls of vessels or chambers, collagen fibers of fibrous tissues, or that same matrix within connective tissue ensheathing muscle cells or in the lacunae of bone. It is similarly the space occupied by the plasma of the circulation and the lymph fluid of the lymphatic channels. Delivery to the interstitial space of muscle tissue is preferred for the reasons discussed below. They may be conveniently delivered by injection into the tissues comprising these cells. They are preferably delivered to and ~~expressed in persistent, non-dividing cells which are differentiated, although delivery~~ and expression may be achieved in non-differentiated or less completely differentiated cells, such as, for example, stem cells of blood or skin fibroblasts. In vivo muscle cells are particularly competent in their ability to take up and express polynucleotides.

For the naked acid sequence injection, an effective dosage amount of DNA or RNA will be in the range of from about 0.05 mg/kg body weight to about 50 mg/kg body weight. Preferably the dosage will be from about 0.005 mg/kg to about 20 mg/kg and more preferably from about 0.05 mg/kg to about 5 mg/kg. Of course, as the artisan of ordinary skill will appreciate, this dosage will vary according to the tissue site of injection. The appropriate and effective dosage of nucleic acid sequence can readily be determined by those of ordinary skill in the art and may depend on the condition being treated and the route of administration.

The preferred route of administration is by the parenteral route of injection into the interstitial space of tissues. However, other parenteral routes may also be used, such as, inhalation of an aerosol formulation particularly for delivery to lungs or bronchial tissues, throat or mucous membranes of the nose. In addition, naked

09376430-091899

CRCGCL DNA constructs can be delivered to arteries during angioplasty by the catheter used in the procedure.

5 The naked polynucleotides are delivered by any method known in the art, including, but not limited to, direct needle injection at the delivery site, intravenous injection, topical administration, catheter infusion, and so-called "gene guns". These delivery methods are known in the art.

As is evidenced in the Examples, naked CRCGCL nucleic acid sequences can be administered in vivo results in the successful expression of CRCGCL polypeptide in the femoral arteries of rabbits.

10 The constructs may also be delivered with delivery vehicles such as viral sequences, viral particles, liposome formulations, lipofectin, precipitating agents, etc. Such methods of delivery are known in the art.

15 In certain embodiments, the CRCGCL polynucleotide constructs are complexed in a liposome preparation. Liposomal preparations for use in the instant invention include cationic (positively charged), anionic (negatively charged) and neutral preparations. However, cationic liposomes are particularly preferred because a tight charge complex can be formed between the cationic liposome and the ~~polyanionic nucleic acid~~. ~~Cationic liposomes have been shown to mediate~~ intracellular delivery of plasmid DNA (Felgner et al., Proc. Natl. Acad. Sci. USA  
20 (1987) 84:7413-7416, which is herein incorporated by reference); mRNA (Malone et al., Proc. Natl. Acad. Sci. USA (1989) 86:6077-6081, which is herein incorporated by reference); and purified transcription factors (Debs et al., J. Biol. Chem. (1990) 265:10189-10192, which is herein incorporated by reference), in functional form.

25 Cationic liposomes are readily available. For example, N[1-2,3-dioleoyloxy)propyl]-N,N,N-triethylammonium (DOTMA) liposomes are particularly useful and are available under the trademark Lipofectin, from GIBCO BRL, Grand Island, N.Y. (See, also, Felgner et al., Proc. Natl Acad. Sci. USA (1987) 84:7413-7416, which is herein incorporated by reference). Other commercially available liposomes include transfectace (DDAB/DOPE) and DOTAP/DOPE  
30 (Boehringer).

Other cationic liposomes can be prepared from readily available materials using techniques well known in the art. See, e.g. PCT Publication No. WO 90/11092 (which is herein incorporated by reference) for a description of the synthesis of

DOTAP (1,2-bis(oleoyloxy)-3-(trimethylammonio)propane) liposomes. Preparation of DOTMA liposomes is explained in the literature, see, e.g., P. Felgner et al., Proc. Natl. Acad. Sci. USA 84:7413-7417, which is herein incorporated by reference. Similar methods can be used to prepare liposomes from other cationic lipid materials.

5 Similarly, anionic and neutral liposomes are readily available, such as from Avanti Polar Lipids (Birmingham, Ala.), or can be easily prepared using readily available materials. Such materials include phosphatidyl, choline, cholesterol, phosphatidyl ethanolamine, dioleoylphosphatidyl choline (DOPC),  
10 dioleoylphosphatidyl glycerol (DOPG), dioleoylphosphatidyl ethanolamine (DOPE), among others. These materials can also be mixed with the DOTMA and DOTAP starting materials in appropriate ratios. Methods for making liposomes using these materials are well known in the art.

For example, commercially dioleoylphosphatidyl choline (DOPC),  
15 dioleoylphosphatidyl glycerol (DOPG), and dioleoylphosphatidyl ethanolamine (DOPE) can be used in various combinations to make conventional liposomes, with or without the addition of cholesterol. Thus, for example, DOPG/DOPC vesicles can be prepared by drying 50 mg each of DOPG and DOPC under a stream of nitrogen gas  
into a sonication vial. The sample is placed under a vacuum pump overnight and is  
hydrated the following day with deionized water. The sample is then sonicated for 2  
20 hours in a capped vial, using a Heat Systems model 350 sonicator equipped with an inverted cup (bath type) probe at the maximum setting while the bath is circulated at 15EC. Alternatively, negatively charged vesicles can be prepared without sonication to produce multilamellar vesicles or by extrusion through nucleopore membranes to produce unilamellar vesicles of discrete size. Other methods are known and available  
25 to those of skill in the art.

The liposomes can comprise multilamellar vesicles (MLVs), small unilamellar vesicles (SUVs), or large unilamellar vesicles (LUVs), with SUVs being preferred. The various liposome-nucleic acid complexes are prepared using methods well known in the art. See, e.g., Straubinger et al., Methods of Immunology (1983), 101:512-527,  
30 which is herein incorporated by reference. For example, MLVs containing nucleic acid can be prepared by depositing a thin film of phospholipid on the walls of a glass tube and subsequently hydrating with a solution of the material to be encapsulated. SUVs are prepared by extended sonication of MLVs to produce a homogeneous

66780-06492660

population of unilamellar liposomes. The material to be entrapped is added to a suspension of preformed MLVs and then sonicated. When using liposomes containing cationic lipids, the dried lipid film is resuspended in an appropriate solution such as sterile water or an isotonic buffer solution such as 10 mM Tris/NaCl, sonicated, and then the preformed liposomes are mixed directly with the DNA. The liposome and DNA form a very stable complex due to binding of the positively charged liposomes to the cationic DNA. SUVs find use with small nucleic acid fragments. LUVs are prepared by a number of methods, well known in the art. Commonly used methods include  $\text{Ca}^{2+}$ -EDTA chelation (Papahadjopoulos et al., Biochim. Biophys. Acta (1975) 394:483; Wilson et al., Cell (1979) 17:77); ether injection (Deamer, D. and Bangham, A., Biochim. Biophys. Acta (1976) 443:629; Ostro et al., Biochem. Biophys. Res. Commun. (1977) 76:836; Fraley et al., Proc. Natl. Acad. Sci. USA (1979) 76:3348); detergent dialysis (Enoch, H. and Strittmatter, P., Proc. Natl. Acad. Sci. USA (1979) 76:145); and reverse-phase evaporation (REV) (Fraley et al., J. Biol. Chem. (1980) 255:10431; Szoka, F. and Papahadjopoulos, D., Proc. Natl. Acad. Sci. USA (1978) 75:145; Schaefer-Ridder et al., Science (1982) 215:166), which are herein incorporated by reference.

Generally, the ratio of DNA to liposomes will be from about 10:1 to about 1:10. Preferably, the ratio will be from about 5:1 to about 1:5. More preferably, the ratio will be about 3:1 to about 1:3. Still more preferably, the ratio will be about 1:1.

U.S. Patent No. 5,676,954 (which is herein incorporated by reference) reports on the injection of genetic material, complexed with cationic liposomes carriers, into mice. U.S. Patent Nos. 4,897,355, 4,946,787, 5,049,386, 5,459,127, 5,589,466, 5,693,622, 5,580,859, 5,703,055, and international publication no. WO 94/9469 (which are herein incorporated by reference) provide cationic lipids for use in transfecting DNA into cells and mammals. U.S. Patent Nos. 5,589,466, 5,693,622, 5,580,859, 5,703,055, and international publication no. WO 94/9469 (which are herein incorporated by reference) provide methods for delivering DNA-cationic lipid complexes to mammals.

In certain embodiments, cells are be engineered, ex vivo or in vivo, using a retroviral particle containing RNA which comprises a sequence encoding CRGCL. Retroviruses from which the retroviral plasmid vectors may be derived include, but are not limited to, Moloney Murine Leukemia Virus, spleen necrosis virus, Rous

sarcoma Virus, Harvey Sarcoma Virus, avian leukosis virus, gibbon ape leukemia virus, human immunodeficiency virus, Myeloproliferative Sarcoma Virus, and mammary tumor virus.

5 The retroviral plasmid vector is employed to transduce packaging cell lines to form producer cell lines. Examples of packaging cells which may be transfected include, but are not limited to, the PE501, PA317, R-2, R-AM, PA12, T19-14X, VT-19-17-H2, RCRE, RCRIP, GP+E-86, GP+envAm12, and DAN cell lines as described in Miller, Human Gene Therapy 1:5-14 (1990), which is incorporated herein by reference in its entirety. The vector may transduce the packaging cells through any  
10 means known in the art. Such means include, but are not limited to, electroporation, the use of liposomes, and  $\text{CaPO}_4$  precipitation. In one alternative, the retroviral plasmid vector may be encapsulated into a liposome, or coupled to a lipid, and then administered to a host.

The producer cell line generates infectious retroviral vector particles which  
15 include polynucleotide encoding CRCGCL. Such retroviral vector particles then may be employed, to transduce eukaryotic cells, either in vitro or in vivo. The transduced eukaryotic cells will express CRCGCL.

~~In certain other embodiments, cells are engineered, ex vivo or in vivo, with~~  
CRCGCL polynucleotide contained in an adenovirus vector. Adenovirus can be  
20 manipulated such that it encodes and expresses CRCGCL, and at the same time is inactivated in terms of its ability to replicate in a normal lytic viral life cycle. Adenovirus expression is achieved without integration of the viral DNA into the host cell chromosome, thereby alleviating concerns about insertional mutagenesis. Furthermore, adenoviruses have been used as live enteric vaccines for many years  
25 with an excellent safety profile (Schwartz, A. R. et al. (1974) Am. Rev. Respir. Dis.109:233-238). Finally, adenovirus mediated gene transfer has been demonstrated in a number of instances including transfer of alpha-1-antitrypsin and CFTR to the lungs of cotton rats (Rosenfeld, M. A. et al. (1991) Science 252:431-434; Rosenfeld et al., (1992) Cell 68:143-155). Furthermore, extensive studies to attempt to establish  
30 adenovirus as a causative agent in human cancer were uniformly negative (Green, M. et al. (1979) Proc. Natl. Acad. Sci. USA 76:6606).

Suitable adenoviral vectors useful in the present invention are described, for example, in Kozarsky and Wilson, Curr. Opin. Genet. Devel. 3:499-503 (1993);

663100-0249260

Rosenfeld et al., Cell 68:143-155 (1992); Engelhardt et al., Human Genet. Ther. 4:759-769 (1993); Yang et al., Nature Genet. 7:362-369 (1994); Wilson et al., Nature 365:691-692 (1993); and U.S. Patent No. 5,652,224, which are herein incorporated by reference. For example, the adenovirus vector Ad2 is useful and can be grown in  
5 human 293 cells. These cells contain the E1 region of adenovirus and constitutively express Ela and Elb, which complement the defective adenoviruses by providing the products of the genes deleted from the vector. In addition to Ad2, other varieties of adenovirus (e.g., Ad3, Ad5, and Ad7) are also useful in the present invention.

Preferably, the adenoviruses used in the present invention are replication  
10 deficient. Replication deficient adenoviruses require the aid of a helper virus and/or packaging cell line to form infectious particles. The resulting virus is capable of infecting cells and can express a polynucleotide of interest which is operably linked to a promoter, for example, the HARP promoter of the present invention, but cannot replicate in most cells. Replication deficient adenoviruses may be deleted in one or  
15 more of all or a portion of the following genes: E1a, E1b, E3, E4, E2a, or L1 through L5.

In certain other embodiments, the cells are engineered, ex vivo or in vivo, using an adeno-associated virus (AAV). AAVs are naturally occurring defective  
viruses that require helper viruses to produce infectious particles (Muzyczka, N.,  
20 Curr. Topics in Microbiol. Immunol. 158:97 (1992)). It is also one of the few viruses that may integrate its DNA into non-dividing cells. Vectors containing as little as 300 base pairs of AAV can be packaged and can integrate, but space for exogenous DNA is limited to about 4.5 kb. Methods for producing and using such AAVs are known in the art. See, for example, U.S. Patent Nos. 5,139,941, 5,173,414, 5,354,678,  
25 5,436,146, 5,474,935, 5,478,745, and 5,589,377.

For example, an appropriate AAV vector for use in the present invention will include all the sequences necessary for DNA replication, encapsidation, and host-cell integration. The CRCGCL polynucleotide construct is inserted into the AAV vector using standard cloning methods, such as those found in Sambrook et al., Molecular  
30 Cloning: A Laboratory Manual, Cold Spring Harbor Press (1989). The recombinant AAV vector is then transfected into packaging cells which are infected with a helper virus, using any standard technique, including lipofection, electroporation, calcium phosphate precipitation, etc. Appropriate helper viruses include adenoviruses,

68T80\*0E49Z60



cytomegaloviruses, vaccinia viruses, or herpes viruses. Once the packaging cells are transfected and infected, they will produce infectious AAV viral particles which contain the CRCGCL polynucleotide construct. These viral particles are then used to transduce eukaryotic cells, either ex vivo or in vivo. The transduced cells will contain  
5 the CRCGCL polynucleotide construct integrated into its genome, and will express CRCGCL.

Another method of gene therapy involves operably associating heterologous control regions and endogenous polynucleotide sequences (e.g. encoding CRCGCL) via homologous recombination (see, e.g., U.S. Patent No. 5,641,670, issued June 24,  
10 1997; International Publication No. WO 96/29411, published September 26, 1996; International Publication No. WO 94/12650, published August 4, 1994; Koller et al., Proc. Natl. Acad. Sci. USA 86:8932-8935 (1989); and Zijlstra et al., Nature 342:435-438 (1989). This method involves the activation of a gene which is present in the target cells, but which is not normally expressed in the cells, or is expressed at a  
15 lower level than desired.

Polynucleotide constructs are made, using standard techniques known in the art, which contain the promoter with targeting sequences flanking the promoter. ~~Suitable promoters are described herein. The targeting sequence is sufficiently~~  
complementary to an endogenous sequence to permit homologous recombination of  
20 the promoter-targeting sequence with the endogenous sequence. The targeting sequence will be sufficiently near the 5' end of the CRCGCL desired endogenous polynucleotide sequence so the promoter will be operably linked to the endogenous sequence upon homologous recombination.

The promoter and the targeting sequences can be amplified using PCR.  
25 Preferably, the amplified promoter contains distinct restriction enzyme sites on the 5' and 3' ends. Preferably, the 3' end of the first targeting sequence contains the same restriction enzyme site as the 5' end of the amplified promoter and the 5' end of the second targeting sequence contains the same restriction site as the 3' end of the amplified promoter. The amplified promoter and targeting sequences are digested  
30 and ligated together.

The promoter-targeting sequence construct is delivered to the cells, either as naked polynucleotide, or in conjunction with transfection-facilitating agents, such as liposomes, viral sequences, viral particles, whole viruses, lipofection, precipitating

663130-0213260

5           The promoter-targeting sequence construct is taken up by cells. Homologous recombination between the construct and the endogenous sequence takes place, such that an endogenous CRCGCL sequence is placed under the control of the promoter. The promoter then drives the expression of the endogenous CRCGCL sequence.

Preferably, the polynucleotide encoding CRCGCL contains a secretory signal sequence that facilitates secretion of the protein. Typically, the signal sequence is positioned in the coding region of the polynucleotide to be expressed towards or at the 5' end of the coding region. The signal sequence may be homologous or heterologous to the polynucleotide of interest and may be homologous or heterologous to the cells to be transfected. Additionally, the signal sequence may be chemically synthesized using methods known in the art.

Any mode of administration of any of the above-described polynucleotides constructs can be used so long as the mode results in the expression of one or more molecules in an amount sufficient to provide a therapeutic effect. This includes direct needle injection, systemic injection, catheter infusion, biolistic injectors, particle accelerators (i.e., "gene guns"), gelfoam sponge depots, other commercially available depot materials, osmotic pumps (e.g., Alza minipumps), oral or suppository solid (tablet or pill) pharmaceutical formulations, and decanting or topical applications during surgery. For example, direct injection of naked calcium phosphate-precipitated plasmid into rat liver and rat spleen or a protein-coated

plasmid into the portal vein has resulted in gene expression of the foreign gene in the rat livers (Kaneda et al., Science 243:375 (1989)).

5 A preferred method of local administration is by direct injection. Preferably, a recombinant molecule of the present invention complexed with a delivery vehicle is administered by direct injection into or locally within the area of arteries. Administration of a composition locally within the area of arteries refers to injecting the composition centimeters and preferably, millimeters within arteries.

10 Another method of local administration is to contact a polynucleotide construct of the present invention in or around a surgical wound. For example, a patient can undergo surgery and the polynucleotide construct can be coated on the surface of tissue inside the wound or the construct can be injected into areas of tissue inside the wound.

15 Therapeutic compositions useful in systemic administration, include recombinant molecules of the present invention complexed to a targeted delivery vehicle of the present invention. Suitable delivery vehicles for use with systemic administration comprise liposomes comprising ligands for targeting the vehicle to a particular site.

Preferred methods of systemic administration, include intravenous injection, aerosol, oral and percutaneous (topical) delivery. Intravenous injections can be performed using methods standard in the art. Aerosol delivery can also be performed using methods standard in the art (see, for example, Stribling et al., Proc. Natl. Acad. Sci. USA 189:11277-11281, 1992, which is incorporated herein by reference). Oral delivery can be performed by complexing a polynucleotide construct of the present invention to a carrier capable of withstanding degradation by digestive enzymes in the gut of an animal. Examples of such carriers, include plastic capsules or tablets, such as those known in the art. Topical delivery can be performed by mixing a polynucleotide construct of the present invention with a lipophilic reagent (e.g., DMSO) that is capable of passing into the skin.

30 Determining an effective amount of substance to be delivered can depend upon a number of factors including, for example, the chemical structure and biological activity of the substance, the age and weight of the animal, the precise condition requiring treatment and its severity, and the route of administration. The frequency of treatments depends upon a number of factors, such as the amount of

669180-043460

polynucleotide constructs administered per dose, as well as the health and history of the subject. The precise amount, number of doses, and timing of doses will be determined by the attending physician or veterinarian.

Therapeutic compositions of the present invention can be administered to any animal, preferably to mammals and birds. Preferred mammals include humans, dogs, cats, mice, rats, rabbits sheep, cattle, horses and pigs, with humans being particularly preferred.

#### 10 **Biological Activities of CRCGCL**

CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, can be used in assays to test for one or more biological activities. If CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, do exhibit activity in a particular assay, it is likely that CRCGCL may be involved in the diseases associated with the biological activity. Therefore, CRCGCL could be used to treat the associated disease.

#### **Immune Activity**

CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, may be useful in treating deficiencies or disorders of the immune system, by activating or inhibiting the proliferation, differentiation, or mobilization (chemotaxis) of immune cells. Immune cells develop through a process called hematopoiesis, producing myeloid (platelets, red blood cells, neutrophils, and macrophages) and lymphoid (B and T lymphocytes) cells from pluripotent stem cells. The etiology of these immune deficiencies or disorders may be genetic, somatic, such as cancer or some autoimmune disorders, acquired (e.g., by chemotherapy or toxins), or infectious. Moreover, CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, can be used as a marker or detector of a particular immune system disease or disorder.

Interestingly, CRCGCL maps to the pseudoautosomal regions on the X and Y chromosomes. It is likely that mutations in CRCGCL may also lead to immune disorders, especially those involving activated T cells. Moreover, mutations in

CRCGCL may be involved in autoimmune diseases, especially X-linked autoimmune diseases.

CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, may be useful in treating or detecting deficiencies or disorders of hematopoietic cells. CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, could be used to increase differentiation and proliferation of hematopoietic cells, including the pluripotent stem cells, in an effort to treat those disorders associated with a decrease in certain (or many) types hematopoietic cells. Examples of immunologic deficiency syndromes include, but are not limited to: blood protein disorders (e.g. agammaglobulinemia, dysgammaglobulinemia), ataxia telangiectasia, common variable immunodeficiency, Digeorge Syndrome, HIV infection, HTLV-BLV infection, leukocyte adhesion deficiency syndrome, lymphopenia, phagocyte bactericidal dysfunction, severe combined immunodeficiency (SCIDs), Wiskott-Aldrich Disorder, anemia, thrombocytopenia, or hemoglobinuria.

Moreover, CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, can also be used to modulate hemostatic (the stopping of bleeding) or thrombolytic activity (clot formation). For example, by increasing hemostatic or thrombolytic activity, CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, could be used to treat blood coagulation disorders (e.g., afibrinogenemia, factor deficiencies), blood platelet disorders (e.g. thrombocytopenia), or wounds resulting from trauma, surgery, or other causes. Alternatively, CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, that can decrease hemostatic or thrombolytic activity could be used to inhibit or dissolve clotting, important in the treatment of heart attacks (infarction), strokes, or scarring.

CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, may also be useful in treating or detecting autoimmune disorders. Many autoimmune disorders result from inappropriate recognition of self as foreign material by immune cells. This inappropriate recognition results in an immune response leading to the destruction of the host tissue. Therefore, the administration of CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, that can inhibit an immune response, particularly the proliferation, differentiation, or

09375430-081899

chemotaxis of T-cells, may be an effective therapy in preventing autoimmune disorders.

Examples of autoimmune disorders that can be treated or detected include, but are not limited to: Addison's Disease, hemolytic anemia, antiphospholipid syndrome, rheumatoid arthritis, dermatitis, allergic encephalomyelitis, glomerulonephritis, Goodpasture's Syndrome, Graves' Disease, Multiple Sclerosis, Myasthenia Gravis, Neuritis, Ophthalmia, Bullous Pemphigoid, Pemphigus, Polyendocrinopathies, Purpura, Reiter's Disease, Stiff-Man Syndrome, Autoimmune Thyroiditis, Systemic Lupus Erythematosus, Autoimmune Pulmonary Inflammation, Guillain-Barre Syndrome, insulin dependent diabetes mellitis, and autoimmune inflammatory eye disease.

Similarly, allergic reactions and conditions, such as asthma (particularly allergic asthma) or other respiratory problems, may also be treated by CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL. Moreover, these molecules can be used to treat anaphylaxis, hypersensitivity to an antigenic molecule, or blood group incompatibility.

CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, may also be used to treat and/or prevent organ rejection or graft-versus-host disease (GVHD). Organ rejection occurs by host immune cell destruction of the transplanted tissue through an immune response. Similarly, an immune response is also involved in GVHD, but, in this case, the foreign transplanted immune cells destroy the host tissues. The administration of CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, that inhibits an immune response, particularly the proliferation, differentiation, or chemotaxis of T-cells, may be an effective therapy in preventing organ rejection or GVHD.

Similarly, CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, may also be used to modulate inflammation. For example, CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, may inhibit the proliferation and differentiation of cells involved in an inflammatory response. These molecules can be used to treat inflammatory conditions, both chronic and acute conditions, including inflammation associated with infection (e.g., septic shock, sepsis, or systemic inflammatory response syndrome (SIRS)), ischemia-reperfusion injury, endotoxin lethality, arthritis, complement-mediated hyperacute

66730-049260

rejection, nephritis, cytokine or chemokine induced lung injury, inflammatory bowel disease, Crohn's disease, or resulting from over production of cytokines (e.g., TNF or IL-1.)

5           **Hyperproliferative Disorders**

CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, can be used to treat or detect hyperproliferative disorders, including neoplasms. CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, may inhibit the proliferation of the disorder through direct or indirect  
10 interactions. Alternatively, CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, may proliferate other cells which can inhibit the hyperproliferative disorder.

For example, by increasing an immune response, particularly increasing antigenic qualities of the hyperproliferative disorder or by proliferating,  
15 differentiating, or mobilizing T-cells, hyperproliferative disorders can be treated. This immune response may be increased by either enhancing an existing immune response, or by initiating a new immune response. Alternatively, decreasing an immune response may also be a method of treating hyperproliferative disorders, such  
as a chemotherapeutic agent.

20           Examples of hyperproliferative disorders that can be treated or detected by CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, include, but are not limited to neoplasms located in the: abdomen, bone, breast, digestive system, liver, pancreas, peritoneum, endocrine glands (adrenal, parathyroid, pituitary, testicles, ovary, thymus, thyroid), eye, head and neck, nervous (central and  
25 peripheral), lymphatic system, pelvic, skin, soft tissue, spleen, thoracic, and urogenital.

Similarly, other hyperproliferative disorders can also be treated or detected by CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL. Examples of such hyperproliferative disorders include, but are not limited to:  
30 hypergammaglobulinemia, lymphoproliferative disorders, paraproteinemias, purpura, sarcoidosis, Sezary Syndrome, Waldenstrom's Macroglobulinemia, Gaucher's Disease, histiocytosis, and any other hyperproliferative disease, besides neoplasia, located in an organ system listed above.

00376430-001890

### Cardiovascular Disorders

CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, encoding CRCGCL may be used to treat cardiovascular disorders, including peripheral artery disease, such as limb ischemia.

Cardiovascular disorders include cardiovascular abnormalities, such as arterio-arterial fistula, arteriovenous fistula, cerebral arteriovenous malformations, congenital heart defects, pulmonary atresia, and Scimitar Syndrome. Congenital heart defects include aortic coarctation, cor triatriatum, coronary vessel anomalies, crisscross heart, dextrocardia, patent ductus arteriosus, Ebstein's anomaly, Eisenmenger complex, hypoplastic left heart syndrome, levocardia, tetralogy of fallot, transposition of great vessels, double outlet right ventricle, tricuspid atresia, persistent truncus arteriosus, and heart septal defects, such as aortopulmonary septal defect, endocardial cushion defects, Lutembacher's Syndrome, trilogy of Fallot, ventricular heart septal defects.

Cardiovascular disorders also include heart disease, such as arrhythmias, carcinoid heart disease, high cardiac output, low cardiac output, cardiac tamponade, endocarditis (including bacterial), heart aneurysm, cardiac arrest, congestive heart failure, congestive cardiomyopathy, paroxysmal dyspnea, cardiac edema, heart hypertrophy, congestive cardiomyopathy, left ventricular hypertrophy, right ventricular hypertrophy, post-infarction heart rupture, ventricular septal rupture, heart valve diseases, myocardial diseases, myocardial ischemia, pericardial effusion, pericarditis (including constrictive and tuberculous), pneumopericardium, postpericardiotomy syndrome, pulmonary heart disease, rheumatic heart disease, ventricular dysfunction, hyperemia, cardiovascular pregnancy complications, Scimitar Syndrome, cardiovascular syphilis, and cardiovascular tuberculosis.

Arrhythmias include sinus arrhythmia, atrial fibrillation, atrial flutter, bradycardia, extrasystole, Adams-Stokes Syndrome, bundle-branch block, sinoatrial block, long QT syndrome, parasystole, Lown-Ganong-Levine Syndrome, Mahaim-type pre-excitation syndrome, Wolff-Parkinson-White syndrome, sick sinus syndrome, tachycardias, and ventricular fibrillation. Tachycardias include paroxysmal tachycardia, supraventricular tachycardia, accelerated idioventricular rhythm, atrioventricular nodal reentry tachycardia, ectopic atrial tachycardia, ectopic

663730-01492660



Heart valve disease include aortic valve insufficiency, aortic valve stenosis, hear murmurs, aortic valve prolapse, mitral valve prolapse, tricuspid valve prolapse, mitral valve insufficiency, mitral valve stenosis, pulmonary atresia, pulmonary valve insufficiency, pulmonary valve stenosis, tricuspid atresia, tricuspid valve insufficiency, and tricuspid valve stenosis.

Myocardial ischemias include coronary disease, such as angina pectoris, coronary aneurysm, coronary arteriosclerosis, coronary thrombosis, coronary vasospasm, myocardial infarction and myocardial stunning.

Aneurysms include dissecting aneurysms, false aneurysms, infected aneurysms, ruptured aneurysms, aortic aneurysms, cerebral aneurysms, coronary aneurysms, heart aneurysms, and iliac aneurysms.

Arterial occlusive diseases include arteriosclerosis, intermittent claudication, carotid stenosis, fibromuscular dysplasias, mesenteric vascular occlusion, Moyamoya

disease, renal artery obstruction, retinal artery occlusion, and thromboangiitis obliterans.

Cerebrovascular disorders include carotid artery diseases, cerebral amyloid angiopathy, cerebral aneurysm, cerebral anoxia, cerebral arteriosclerosis, cerebral arteriovenous malformation, cerebral artery diseases, cerebral embolism and thrombosis, carotid artery thrombosis, sinus thrombosis, Wallenberg's syndrome, cerebral hemorrhage, epidural hematoma, subdural hematoma, subarachnoid hemorrhage, cerebral infarction, cerebral ischemia (including transient), subclavian steal syndrome, periventricular leukomalacia, vascular headache, cluster headache, migraine, and vertebrobasilar insufficiency.

Embolisms include air embolisms, amniotic fluid embolisms, cholesterol embolisms, blue toe syndrome, fat embolisms, pulmonary embolisms, and thromboembolisms. Thrombosis include coronary thrombosis, hepatic vein thrombosis, retinal vein occlusion, carotid artery thrombosis, sinus thrombosis, Wallenberg's syndrome, and thrombophlebitis.

Ischemia includes cerebral ischemia, ischemic colitis, compartment syndromes, anterior compartment syndrome, myocardial ischemia, reperfusion injuries, and peripheral limb ischemia. Vasculitis includes aortitis, arteritis, Behcet's Syndrome, Churg-Strauss Syndrome, mucocutaneous lymph node syndrome, thromboangiitis obliterans, hypersensitivity vasculitis, Schoenlein-Henoch purpura, allergic cutaneous vasculitis, and Wegener's granulomatosis.

CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, are especially effective for the treatment of critical limb ischemia and coronary disease. As shown in the Examples, administration of CRCGCL polynucleotides and polypeptides to an experimentally induced ischemia rabbit hindlimb may restore blood pressure ratio, blood flow, angiographic score, and capillary density.

CRCGCL polypeptides may be administered using any method known in the art, including, but not limited to, direct needle injection at the delivery site, intravenous injection, topical administration, catheter infusion, biolistic injectors, particle accelerators, gelfoam sponge depots, other commercially available depot materials, osmotic pumps, oral or suppository solid pharmaceutical formulations, decanting or topical applications during surgery, aerosol delivery. Such methods are

558180-0649260

known in the art. CRCGCL polypeptides may be administered as part of a pharmaceutical composition, described in more detail below. Methods of delivering CRCGCL polynucleotides are described in more detail herein.

5 **Anti-Angiogenesis Activity**

The naturally occurring balance between endogenous stimulators and inhibitors of angiogenesis is one in which inhibitory influences predominate. Rastinejad *et al.*, *Cell* 56:345-355 (1989). In those rare instances in which neovascularization occurs under normal physiological conditions, such as wound  
10 healing, organ regeneration, embryonic development, and female reproductive processes, angiogenesis is stringently regulated and spatially and temporally delimited. Under conditions of pathological angiogenesis such as that characterizing solid tumor growth, these regulatory controls fail. Unregulated angiogenesis becomes pathologic and sustains progression of many neoplastic and non-neoplastic diseases.  
15 A number of serious diseases are dominated by abnormal neovascularization including solid tumor growth and metastases, arthritis, some types of eye disorders, and psoriasis. See, e.g., reviews by Moses *et al.*, *Biotech.* 9:630-634 (1991); Folkman *et al.*, *N. Engl. J. Med.*, 333:1757-1763 (1995); Auerbach *et al.*, *J. Microvasc. Res.* 29:401-411 (1985); Folkman, *Advances in Cancer Research*, eds. Klein and  
20 Weinhouse, Academic Press, New York, pp. 175-203 (1985); Patz, *Am. J. Ophthalmol.* 94:715-743 (1982); and Folkman *et al.*, *Science* 221:719-725 (1983). In a number of pathological conditions, the process of angiogenesis contributes to the disease state. For example, significant data have accumulated which suggest that the growth of solid tumors is dependent on angiogenesis. Folkman and Klagsbrun,  
25 *Science* 235:442-447 (1987).

The present invention provides for treatment of diseases or disorders associated with neovascularization by administration of the CRCGCL polynucleotides and/or polypeptides of the invention, as well as agonists or antagonists of CRCGCL. Malignant and metastatic conditions which can be treated with the polynucleotides  
30 and polypeptides, or agonists or antagonists of the invention include, but are not limited to, malignancies, solid tumors, and cancers described herein and otherwise known in the art (for a review of such disorders, see Fishman *et al.*, *Medicine*, 2d Ed., J. B. Lippincott Co., Philadelphia (1985)):

Ocular disorders associated with neovascularization which can be treated with the CRCGCL polynucleotides and polypeptides of the present invention (including CRCGCL agonists and/or antagonists) include, but are not limited to: neovascular glaucoma, diabetic retinopathy, retinoblastoma, retrolental fibroplasia, uveitis, retinopathy of prematurity macular degeneration, corneal graft neovascularization, as well as other eye inflammatory diseases, ocular tumors and diseases associated with choroidal or iris neovascularization. See, e.g., reviews by Waltman *et al.*, *Am. J. Ophthalm.* 85:704-710 (1978) and Gartner *et al.*, *Surv. Ophthalm.* 22:291-312 (1978).

Additionally, disorders which can be treated with the CRCGCL polynucleotides and polypeptides of the present invention (including CRCGCL agonist and/or antagonists) include, but are not limited to, hemangioma, arthritis, psoriasis, angiofibroma, atherosclerotic plaques, delayed wound healing, granulations, hemophilic joints, hypertrophic scars, nonunion fractures, Osler-Weber syndrome, pyogenic granuloma, scleroderma, trachoma, and vascular adhesions.

Moreover, disorders and/or states, which can be treated with be treated with the CRCGCL polynucleotides and polypeptides of the present invention (including CRCGCL agonist and/or antagonists) include, but are not limited to, solid tumors, blood born tumors such as leukemias, tumor metastasis, Kaposi's sarcoma, benign tumors, for example hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas, rheumatoid arthritis, psoriasis, ocular angiogenic diseases, for example, diabetic retinopathy, retinopathy of prematurity, macular degeneration, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, rubeosis, retinoblastoma, and uvietis, delayed wound healing, endometriosis, vascluogenesis, granulations, hypertrophic scars (keloids), nonunion fractures, scleroderma, trachoma, vascular adhesions, myocardial angiogenesis, coronary collaterals, cerebral collaterals, arteriovenous malformations, ischemic limb angiogenesis, Osler-Webber Syndrome, plaque neovascularization, telangiectasia, hemophiliac joints, angiofibroma fibromuscular dysplasia, wound granulation, Crohn's disease, atherosclerosis, birth control agent by preventing vascularization required for embryo implantation controlling menstruation, diseases that have angiogenesis as a pathologic consequence such as cat scratch disease (Rochele minalia quintosa), ulcers (*Helicobacter pylori*), Bartonellosis and bacillary angiomatosis.

**Diseases at the Cellular Level**

Diseases associated with increased cell survival or the inhibition of apoptosis that could be treated or detected by CRCGCL polynucleotides or polypeptides, as well as antagonists or agonists of CRCGCL, include cancers (such as follicular lymphomas, carcinomas with p53 mutations, and hormone-dependent tumors, including, but not limited to colon cancer, cardiac tumors, pancreatic cancer, melanoma, retinoblastoma, glioblastoma, lung cancer, intestinal cancer, testicular cancer, stomach cancer, neuroblastoma, myxoma, myoma, lymphoma, endothelioma, osteoblastoma, osteoclastoma, osteosarcoma, chondrosarcoma, adenoma, breast cancer, prostate cancer, Kaposi's sarcoma and ovarian cancer); autoimmune disorders (such as, multiple sclerosis, Sjogren's syndrome, Hashimoto's thyroiditis, biliary cirrhosis, Behcet's disease, Crohn's disease, polymyositis, systemic lupus erythematosus and immune-related glomerulonephritis and rheumatoid arthritis) and viral infections (such as herpes viruses, pox viruses and adenoviruses), inflammation, graft v. host disease, acute graft rejection, and chronic graft rejection. In preferred embodiments, CRCGCL polynucleotides, polypeptides, and/or antagonists of the invention are used to inhibit growth, progression, and/or metasis of cancers, in particular those listed above.

Additional diseases or conditions associated with increased cell survival that could be treated or detected by CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, include, but are not limited to, progression, and/or metastases of malignancies and related disorders such as leukemia (including acute leukemias (e.g., acute lymphocytic leukemia, acute myelocytic leukemia (including myeloblastic, promyelocytic, myelomonocytic, monocytic, and erythroleukemia)) and chronic leukemias (e.g., chronic myelocytic (granulocytic) leukemia and chronic lymphocytic leukemia)), polycythemia vera, lymphomas (e.g., Hodgkin's disease and non-Hodgkin's disease), multiple myeloma, Waldenstrom's macroglobulinemia, heavy chain disease, and solid tumors including, but not limited to, sarcomas and carcinomas such as fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal

cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilm's tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, menangioma, melanoma, neuroblastoma, and retinoblastoma.

Diseases associated with increased apoptosis that could be treated or detected by CRCGCL polynucleotides or polypeptides, as well as agonists or antagonists of CRCGCL, include AIDS; neurodegenerative disorders (such as Alzheimer's disease, Parkinson's disease, Amyotrophic lateral sclerosis, Retinitis pigmentosa, Cerebellar degeneration and brain tumor or prior associated disease); autoimmune disorders (such as, multiple sclerosis, Sjogren's syndrome, Hashimoto's thyroiditis, biliary cirrhosis, Behcet's disease, Crohn's disease, polymyositis, systemic lupus erythematosus and immune-related glomerulonephritis and rheumatoid arthritis) myelodysplastic syndromes (such as aplastic anemia), graft v. host disease, ischemic injury (such as that caused by myocardial infarction, stroke and reperfusion injury), liver injury (e.g., hepatitis related liver injury, ischemia/reperfusion injury, cholestosis (bile duct injury) and liver cancer); toxin-induced liver disease (such as that caused by alcohol), septic shock, cachexia and anorexia.

### **Wound Healing and Epithelial Cell Proliferation**

In accordance with yet a further aspect of the present invention, there is provided a process for utilizing CRCGCL polynucleotides or polypeptides, as well as agonists or antagonists of CRCGCL, for therapeutic purposes, for example, to stimulate epithelial cell proliferation and basal keratinocytes for the purpose of wound healing, and to stimulate hair follicle production and healing of dermal wounds. CRCGCL polynucleotides or polypeptides, as well as agonists or antagonists of CRCGCL, may be clinically useful in stimulating wound healing including surgical wounds, excisional wounds, deep wounds involving damage of the dermis and epidermis, eye tissue wounds, dental tissue wounds, oral cavity wounds, diabetic ulcers, dermal ulcers, cubitus ulcers, arterial ulcers, venous stasis ulcers, burns

5

10

25

30

have a cytoprotective effect on the small intestine mucosa. CRCGCL polynucleotides or polypeptides, as well as agonists or antagonists of CRCGCL, may also stimulate healing of mucositis (mouth ulcers) that result from chemotherapy and viral infections.

5 CRCGCL polynucleotides or polypeptides, as well as agonists or antagonists of CRCGCL, could further be used in full regeneration of skin in full and partial thickness skin defects, including burns, (i.e., repopulation of hair follicles, sweat glands, and sebaceous glands), treatment of other skin defects such as psoriasis. CRCGCL polynucleotides or polypeptides, as well as agonists or antagonists of  
10 CRCGCL, could be used to treat epidermolysis bullosa, a defect in adherence of the epidermis to the underlying dermis which results in frequent, open and painful blisters by accelerating reepithelialization of these lesions. CRCGCL polynucleotides or polypeptides, as well as agonists or antagonists of CRCGCL, could also be used to treat gastric and duodenal ulcers and help heal by scar formation of the mucosal lining  
15 and regeneration of glandular mucosa and duodenal mucosal lining more rapidly. Inflammatory bowel diseases, such as Crohn's disease and ulcerative colitis, are diseases which result in destruction of the mucosal surface of the small or large intestine, respectively. Thus, CRCGCL polynucleotides or polypeptides, as well as agonists or antagonists of CRCGCL, could be used to promote the resurfacing of the  
20 mucosal surface to aid more rapid healing and to prevent progression of inflammatory bowel disease. Treatment with CRCGCL polynucleotides or polypeptides, agonists or antagonists of CRCGCL, is expected to have a significant effect on the production of mucus throughout the gastrointestinal tract and could be used to protect the intestinal mucosa from injurious substances that are ingested or following surgery. CRCGCL  
25 polynucleotides or polypeptides, as well as agonists or antagonists of CRCGCL, could be used to treat diseases associated with the under expression of CRCGCL.

Moreover, CRCGCL polynucleotides or polypeptides, as well as agonists or antagonists of CRCGCL, could be used to prevent and heal damage to the lungs due to various pathological states. A growth factor such as CRCGCL polynucleotides or  
30 polypeptides, as well as agonists or antagonists of CRCGCL, which could stimulate proliferation and differentiation and promote the repair of alveoli and bronchiolar epithelium to prevent or treat acute or chronic lung damage. For example, emphysema, which results in the progressive loss of alveoli, and inhalation injuries,

668780-02492260



i.e., resulting from smoke inhalation and burns, that cause necrosis of the bronchiolar epithelium and alveoli could be effectively treated using CRCGCL polynucleotides or polypeptides, agonists or antagonists of CRCGCL. Also, CRCGCL polynucleotides or polypeptides, as well as agonists or antagonists of CRCGCL, could be used to stimulate the proliferation of and differentiation of type II pneumocytes, which may help treat or prevent disease such as hyaline membrane diseases, such as infant respiratory distress syndrome and bronchopulmonary dysplasia, in premature infants.

CRCGCL polynucleotides or polypeptides, as well as agonists or antagonists of CRCGCL, could stimulate the proliferation and differentiation of hepatocytes and, thus, could be used to alleviate or treat liver diseases and pathologies such as fulminant liver failure caused by cirrhosis, liver damage caused by viral hepatitis and toxic substances (i.e., acetaminophen, carbon tetrachloride and other hepatotoxins known in the art).

In addition, CRCGCL polynucleotides or polypeptides, as well as agonists or antagonists of CRCGCL, could be used to treat or prevent the onset of diabetes mellitus. In patients with newly diagnosed Types I and II diabetes, where some islet cell function remains, CRCGCL polynucleotides or polypeptides, as well as agonists or antagonists of CRCGCL, could be used to maintain the islet function so as to alleviate, delay or prevent permanent manifestation of the disease. Also, CRCGCL polynucleotides or polypeptides, as well as agonists or antagonists of CRCGCL, could be used as an auxiliary in islet cell transplantation to improve or promote islet cell function.

#### **Infectious Disease**

CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, can be used to treat or detect infectious agents. For example, by increasing the immune response, particularly increasing the proliferation and differentiation of B and/or T cells, infectious diseases may be treated. The immune response may be increased by either enhancing an existing immune response, or by initiating a new immune response. Alternatively, CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, may also directly inhibit the infectious agent, without necessarily eliciting an immune response.

5

10

15

20

25

30

uveitis), gingivitis, opportunistic infections (e.g., AIDS related infections), paronychia, prosthesis-related infections, Reiter's Disease, respiratory tract infections, such as Whooping Cough or Empyema, sepsis, Lyme Disease, Cat-Scratch Disease, Dysentery, Paratyphoid Fever, food poisoning, Typhoid, pneumonia, Gonorrhea, meningitis, Chlamydia, Syphilis, Diphtheria, Leprosy, Paratuberculosis, Tuberculosis, Lupus, Botulism, gangrene, tetanus, impetigo, Rheumatic Fever, Scarlet Fever, sexually transmitted diseases, skin diseases (e.g., cellulitis, dermatocycoses), toxemia, urinary tract infections, wound infections. CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, can be used to treat or detect any of these symptoms or diseases.

Moreover, parasitic agents causing disease or symptoms that can be treated or detected by CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, include, but not limited to, the following families: Amebiasis, Babesiosis, Coccidiosis, Cryptosporidiosis, Dientamoebiasis, Dourine, Ectoparasitic, Giardiasis, Helminthiasis, Leishmaniasis, Theileriasis, Toxoplasmosis, Trypanosomiasis, and Trichomonas. These parasites can cause a variety of diseases or symptoms, including, but not limited to: Scabies, Trombiculiasis, eye infections, intestinal disease (e.g., dysentery, giardiasis), liver disease, lung disease, opportunistic infections (e.g., AIDS related), Malaria, pregnancy complications, and toxoplasmosis. CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, can be used to treat or detect any of these symptoms or diseases.

Preferably, treatment using CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, could either be by administering an effective amount of CRCGCL polypeptide to the patient, or by removing cells from the patient, supplying the cells with CRCGCL polynucleotide, and returning the engineered cells to the patient (ex vivo therapy). Moreover, the CRCGCL polypeptide or polynucleotide can be used as an antigen in a vaccine to raise an immune response against infectious disease.

### Regeneration

CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, can be used to differentiate, proliferate, and attract cells, leading to the regeneration of tissues. (See, Science 276:59-87 (1997).) The regeneration of tissues

could be used to repair, replace, or protect tissue damaged by congenital defects, trauma (wounds, burns, incisions, or ulcers), age, disease (e.g. osteoporosis, osteoarthritis, periodontal disease, liver failure), surgery, including cosmetic plastic surgery, fibrosis, reperfusion injury, or systemic cytokine damage.

5           Tissues that could be regenerated using the present invention include organs (e.g., pancreas, liver, intestine, kidney, skin, endothelium), muscle (smooth, skeletal or cardiac), vasculature (including vascular and lymphatics), nervous, hematopoietic, and skeletal (bone, cartilage, tendon, and ligament) tissue. Preferably, regeneration occurs without or decreased scarring. Regeneration also may include angiogenesis.

10           Moreover, CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, may increase regeneration of tissues difficult to heal. For example, increased tendon/ligament regeneration would quicken recovery time after damage. CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, of the present invention could also be used prophylactically in an effort to  
15           avoid damage. Specific diseases that could be treated include of tendinitis, carpal tunnel syndrome, and other tendon or ligament defects. A further example of tissue regeneration of non-healing wounds includes pressure ulcers, ulcers associated with vascular insufficiency, surgical, and traumatic wounds.

            Similarly, nerve and brain tissue could also be regenerated by using CRCGCL  
20           polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, to proliferate and differentiate nerve cells. Diseases that could be treated using this method include central and peripheral nervous system diseases, neuropathies, or mechanical and traumatic disorders (e.g., spinal cord disorders, head trauma, cerebrovascular disease, and stroke). Specifically, diseases associated with peripheral nerve injuries, peripheral  
25           neuropathy (e.g., resulting from chemotherapy or other medical therapies), localized neuropathies, and central nervous system diseases (e.g., Alzheimer's disease, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome), could all be treated using the CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL.

30

#### Chemotaxis

CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, may have chemotaxis activity. A chemotaxic molecule attracts or

mobilizes cells (e.g., monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils, epithelial and/or endothelial cells) to a particular site in the body, such as inflammation, infection, or site of hyperproliferation. The mobilized cells can then fight off and/or heal the particular trauma or abnormality.

5           CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, may increase chemotactic activity of particular cells. These chemotactic molecules can then be used to treat inflammation, infection, hyperproliferative disorders, or any immune system disorder by increasing the number of cells targeted to a particular location in the body. For example, chemotactic molecules can be used  
10 to treat wounds and other trauma to tissues by attracting immune cells to the injured location. As a chemotactic molecule, CRCGCL could also attract fibroblasts, which can be used to treat wounds.

          It is also contemplated that CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, may inhibit chemotactic activity. These  
15 molecules could also be used to treat disorders. Thus, CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, could be used as an inhibitor of chemotaxis.

#### **Binding Activity**

20           CRCGCL polypeptides may be used to screen for molecules that bind to CRCGCL or for molecules to which CRCGCL binds. The binding of CRCGCL and the molecule may activate (agonist), increase, inhibit (antagonist), or decrease activity of the CRCGCL or the molecule bound. Examples of such molecules include antibodies, oligonucleotides, proteins (e.g., receptors), or small molecules.

25           Preferably, the molecule is closely related to the natural ligand of CRCGCL, e.g., a fragment of the ligand, or a natural substrate, a ligand, a structural or functional mimetic. (See, Coligan et al., Current Protocols in Immunology 1(2):Chapter 5 (1991).) Similarly, the molecule can be closely related to the natural receptor to which CRCGCL binds, or at least, a fragment of the receptor capable of being bound  
30 by CRCGCL (e.g., active site). In either case, the molecule can be rationally designed using known techniques.

          Preferably, the screening for these molecules involves producing appropriate cells which express CRCGCL, either as a secreted protein or on the cell membrane.

56T199-061926A

Preferred cells include cells from mammals, yeast, *Drosophila*, or *E. coli*. Cells expressing CRCGCL(or cell membrane containing the expressed polypeptide) are then preferably contacted with a test compound potentially containing the molecule to observe binding, stimulation, or inhibition of activity of either CRCGCL or the molecule.

The assay may simply test binding of a candidate compound to CRCGCL, wherein binding is detected by a label, or in an assay involving competition with a labeled competitor. Further, the assay may test whether the candidate compound results in a signal generated by binding to CRCGCL.

Alternatively, the assay can be carried out using cell-free preparations, polypeptide/molecule affixed to a solid support, chemical libraries, or natural product mixtures. The assay may also simply comprise the steps of mixing a candidate compound with a solution containing CRCGCL, measuring CRCGCL/molecule activity or binding, and comparing the CRCGCL/molecule activity or binding to a standard.

Preferably, an ELISA assay can measure CRCGCL level or activity in a sample (e.g., biological sample) using a monoclonal or polyclonal antibody. The antibody can measure CRCGCL level or activity by either binding, directly or indirectly, to CRCGCL or by competing with CRCGCL for a substrate.

Additionally, the receptor to which CRCGCL binds can be identified by numerous methods known to those of skill in the art, for example, ligand panning and FACS sorting (Coligan, et al., *Current Protocols in Immun.*, 1(2), Chapter 5, (1991)). For example, expression cloning is employed wherein polyadenylated RNA is prepared from a cell responsive to the polypeptides, for example, NIH3T3 cells which are known to contain multiple receptors for the FGF family proteins, and SC-3 cells, and a cDNA library created from this RNA is divided into pools and used to transfect COS cells or other cells that are not responsive to the polypeptides. Transfected cells which are grown on glass slides are exposed to the polypeptide of the present invention, after they have been labelled. The polypeptides can be labeled by a variety of means including iodination or inclusion of a recognition site for a site-specific protein kinase.

Following fixation and incubation, the slides are subjected to autoradiographic analysis. Positive pools are identified and sub-pools are prepared and re-

09376430-091899

transfected using an iterative sub-pooling and re-screening process, eventually yielding a single clones that encodes the putative receptor.

As an alternative approach for receptor identification, the labeled polypeptides can be photoaffinity linked with cell membrane or extract preparations that express the receptor molecule. Cross-linked material is resolved by PAGE analysis and exposed to X-ray film. The labeled complex containing the receptors of the polypeptides can be excised, resolved into peptide fragments, and subjected to protein microsequencing. The amino acid sequence obtained from microsequencing would be used to design a set of degenerate oligonucleotide probes to screen a cDNA library to identify the genes encoding the putative receptors.

Moreover, the techniques of gene-shuffling, motif-shuffling, exon-shuffling, and/or codon-shuffling (collectively referred to as "DNA shuffling") may be employed to modulate the activities of CRCGCL thereby effectively generating agonists and antagonists of CRCGCL. *See generally*, U.S. Patent Nos. 5,605,793, 5,811,238, 5,830,721, 5,834,252, and 5,837,458, and Patten, P. A., *et al.*, *Curr. Opinion Biotechnol.* 8:724-33 (1997); Harayama, S. *Trends Biotechnol.* 16(2):76-82 (1998); Hansson, L. O., *et al.*, *J. Mol. Biol.* 287:265-76 (1999); and Lorenzo, M. M. and Blasco, R. *Biotechniques* 24(2):308-13 (1998) (each of these patents and publications are hereby incorporated by reference). In one embodiment, alteration of CRCGCL polynucleotides and corresponding polypeptides may be achieved by DNA shuffling. DNA shuffling involves the assembly of two or more DNA segments into a desired CRCGCL molecule by homologous, or site-specific, recombination. In another embodiment, CRCGCL polynucleotides and corresponding polypeptides may be altered by being subjected to random mutagenesis by error-prone PCR, random nucleotide insertion or other methods prior to recombination. In another embodiment, one or more components, motifs, sections, parts, domains, fragments, etc., of CRCGCL may be recombined with one or more components, motifs, sections, parts, domains, fragments, etc. of one or more heterologous molecules. In preferred embodiments, the heterologous molecules are Cytokine Receptor family members. In further preferred embodiments, the heterologous molecule is a growth factor such as, for example, platelet-derived growth factor (PDGF), insulin-like growth factor (IGF-I), transforming growth factor (TGF)-alpha, epidermal growth factor (EGF), fibroblast growth factor (FGF), TGF-beta, bone morphogenetic protein (BMP)-2,

BMP-4, BMP-5, BMP-6, BMP-7, activins A and B, decapentaplegic(dpp), 60A, OP-2, dorsalin, growth differentiation factors (GDFs), nodal, MIS, inhibin-alpha, TGF-beta1, TGF-beta2, TGF-beta3, TGF-beta5, and glial-derived neurotrophic factor (GDNF).

5 Other preferred fragments are biologically active CRCGCL fragments. Biologically active fragments are those exhibiting activity similar, but not necessarily identical, to an activity of the CRCGCL polypeptide. The biological activity of the fragments may include an improved desired activity, or a decreased undesirable activity.

10 Additionally, this invention provides a method of screening compounds to identify those which modulate the action of the polypeptide of the present invention. An example of such an assay comprises combining a mammalian fibroblast cell, a the polypeptide of the present invention, the compound to be screened and  $^3\text{[H]}$  thymidine under cell culture conditions where the fibroblast cell would normally  
15 proliferate. A control assay may be performed in the absence of the compound to be screened and compared to the amount of fibroblast proliferation in the presence of the compound to determine if the compound stimulates proliferation by determining the uptake of  $^3\text{[H]}$  thymidine in each case. ~~The amount of fibroblast cell proliferation is~~ measured by liquid scintillation chromatography which measures the incorporation of  
20  $^3\text{[H]}$  thymidine. Both agonist and antagonist compounds may be identified by this procedure.

In another method, a mammalian cell or membrane preparation expressing a receptor for a polypeptide of the present invention is incubated with a labeled polypeptide of the present invention in the presence of the compound. The ability of  
25 the compound to enhance or block this interaction could then be measured. Alternatively, the response of a known second messenger system following interaction of a compound to be screened and the CRCGCL receptor is measured and the ability of the compound to bind to the receptor and elicit a second messenger response is measured to determine if the compound is a potential agonist or  
30 antagonist. Such second messenger systems include but are not limited to, cAMP guanylate cyclase, ion channels or phosphoinositide hydrolysis.



66T80\*0249460

All of these above assays can be used as diagnostic or prognostic markers. The molecules discovered using these assays can be used to treat disease or to bring about a particular result in a patient (e.g., blood vessel growth) by activating or inhibiting the CRCGCL/molecule. Moreover, the assays can discover agents which may inhibit or enhance the production of CRCGCL from suitably manipulated cells or tissues. Therefore, the invention includes a method of identifying compounds which bind to CRCGCL comprising the steps of: (a) incubating a candidate binding compound with CRCGCL; and (b) determining if binding has occurred. Moreover, the invention includes a method of identifying agonists/antagonists comprising the steps of: (a) incubating a candidate compound with CRCGCL, (b) assaying a biological activity, and (b) determining if a biological activity of CRCGCL has been altered.

Also, one could identify molecules bind CRCGCL experimentally by using the beta-pleated sheet regions disclosed in Figure 3 and Table 1. Accordingly, specific embodiments of the invention are directed to polynucleotides encoding polypeptides which comprise, or alternatively consist of, the amino acid sequence of each beta pleated sheet regions disclosed in Figure 3/Table 1. Additional embodiments of the invention are directed to polynucleotides encoding CRCGCL polypeptides which comprise, or alternatively consist of, any combination or all of the beta pleated sheet regions disclosed in Figure 3/Table 1. Additional preferred embodiments of the invention are directed to polypeptides which comprise, or alternatively consist of, the CRCGCL amino acid sequence of each of the beta pleated sheet regions disclosed in Figure 3/Table 1. Additional embodiments of the invention are directed to CRCGCL polypeptides which comprise, or alternatively consist of, any combination or all of the beta pleated sheet regions disclosed in Figure 3/Table 1.

#### **Antisense And Ribozyme (Antagonists)**

In specific embodiments, antagonists according to the present invention are nucleic acids corresponding to the sequences contained in SEQ ID NO:1, or the complementary strand thereof, and/or to nucleotide sequences contained in the deposited clone 209641 or 209691. In one embodiment, antisense sequence is generated internally by the organism, in another embodiment, the antisense sequence is separately administered (see, for example, O'Connor, J., Neurochem. 56:560

(1991). Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988). Antisense technology can be used to control gene expression through antisense DNA or RNA, or through triple-helix formation. Antisense techniques are discussed for example, in Okano, J., Neurochem. 56:560 (1991); Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988). Triple helix formation is discussed in, for instance, Lee et al., Nucleic Acids Research 6:3073 (1979); Cooney et al., Science 241:456 (1988); and Dervan et al., Science 251:1300 (1991). The methods are based on binding of a polynucleotide to a complementary DNA or RNA.

For example, the 5' coding portion of a polynucleotide that encodes the mature polypeptide of the present invention may be used to design an antisense RNA oligonucleotide of from about 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription thereby preventing transcription and the production of the receptor. The antisense RNA oligonucleotide hybridizes to the mRNA in vivo and blocks translation of the mRNA molecule into receptor polypeptide.

In one embodiment, the CRCGCL antisense nucleic acid of the invention is produced intracellularly by transcription from an exogenous sequence. For example, a vector or a portion thereof, is transcribed, producing an antisense nucleic acid (RNA) of the invention. Such a vector would contain a sequence encoding the CRCGCL antisense nucleic acid. Such a vector can remain episomal or become chromosomally integrated, as long as it can be transcribed to produce the desired antisense RNA. Such vectors can be constructed by recombinant DNA technology methods standard in the art. Vectors can be plasmid, viral, or others known in the art, used for replication and expression in vertebrate cells. Expression of the sequence encoding CRCGCL, or fragments thereof, can be by any promoter known in the art to act in vertebrate, preferably human cells. Such promoters can be inducible or constitutive. Such promoters include, but are not limited to, the SV40 early promoter region (Bernoist and Chambon, Nature 29:304-310 (1981), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto et al., Cell 22:787-797 (1980), the herpes thymidine promoter (Wagner et al., Proc. Natl. Acad. Sci. U.S.A. 78:1441-1445 (1981), the regulatory sequences of the metallothionein gene (Brinster, et al., Nature 296:39-42 (1982)), etc.

00375430-03100

The antisense nucleic acids of the invention comprise a sequence complementary to at least a portion of an RNA transcript of a CRCGCL gene. However, absolute complementarity, although preferred, is not required. A sequence "complementary to at least a portion of an RNA," referred to herein, means a sequence having sufficient complementarity to be able to hybridize with the RNA, forming a stable duplex; in the case of double stranded CRCGCL antisense nucleic acids, a single strand of the duplex DNA may thus be tested, or triplex formation may be assayed. The ability to hybridize will depend on both the degree of complementarity and the length of the antisense nucleic acid. Generally, the larger the hybridizing nucleic acid, the more base mismatches with a CRCGCL RNA it may contain and still form a stable duplex (or triplex as the case may be). One skilled in the art can ascertain a tolerable degree of mismatch by use of standard procedures to determine the melting point of the hybridized complex.

Oligonucleotides that are complementary to the 5' end of the message, e.g., the 5' untranslated sequence up to and including the AUG initiation codon, should work most efficiently at inhibiting translation. However, sequences complementary to the 3' untranslated sequences of mRNAs have been shown to be effective at inhibiting translation of mRNAs as well. See generally, Wagner, R., 1994, Nature 372:333-335.

Thus, oligonucleotides complementary to either the 5'- or 3'- non- translated, non-coding regions of CRCGCL shown in Figures 1A-B could be used in an antisense approach to inhibit translation of endogenous CRCGCL mRNA. Oligonucleotides complementary to the 5' untranslated region of the mRNA should include the complement of the AUG start codon. Antisense oligonucleotides complementary to mRNA coding regions are less efficient inhibitors of translation but could be used in accordance with the invention. Whether designed to hybridize to the 5'-, 3'- or coding region of CRCGCL mRNA, antisense nucleic acids should be at least six nucleotides in length, and are preferably oligonucleotides ranging from 6 to about 50 nucleotides in length. In specific aspects the oligonucleotide is at least 10 nucleotides, at least 17 nucleotides, at least 25 nucleotides or at least 50 nucleotides.

The polynucleotides of the invention can be DNA or RNA or chimeric mixtures or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone, for example, to improve stability of the molecule, hybridization,

etc. The oligonucleotide may include other appended groups such as peptides (e.g., for targeting host cell receptors in vivo), or agents facilitating transport across the cell membrane (see, e.g., Letsinger et al., 1989, Proc. Natl. Acad. Sci. U.S.A. 86:6553-6556; Lemaitre et al., 1987, Proc. Natl. Acad. Sci. 84:648-652; PCT Publication No. WO88/09810, published December 15, 1988) or the blood-brain barrier (see, e.g., PCT Publication No. WO89/10134, published April 25, 1988), hybridization-triggered cleavage agents. (See, e.g., Krol et al., 1988, BioTechniques 6:958-976) or intercalating agents. (See, e.g., Zon, 1988, Pharm. Res. 5:539-549). To this end, the oligonucleotide may be conjugated to another molecule, e.g., a peptide, hybridization triggered cross-linking agent, transport agent, hybridization-triggered cleavage agent, etc.

The antisense oligonucleotide may comprise at least one modified base moiety which is selected from the group including, but not limited to, 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xantine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, ~~2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine,~~ 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine.

The antisense oligonucleotide may also comprise at least one modified sugar moiety selected from the group including, but not limited to, arabinose, 2-fluoroarabinose, xylulose, and hexose.

In yet another embodiment, the antisense oligonucleotide comprises at least one modified phosphate backbone selected from the group including, but not limited to, a phosphorothioate, a phosphorodithioate, a phosphoramidothioate, a

554130-0613260

In yet another embodiment, the antisense oligonucleotide is an a-anomeric oligonucleotide. An a-anomeric oligonucleotide forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual b-units, the strands run parallel to each other (Gautier et al., 1987, Nucl. Acids Res. 15:6625-6641). The oligonucleotide is a 2'-O-methylribonucleotide (Inoue et al., 1987, Nucl. Acids Res. 15:6131-6148), or a chimeric RNA-DNA analogue (Inoue et al., 1987, FEBS Lett. 215:327-330).

Polynucleotides of the invention may be synthesized by standard methods known in the art, e.g. by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). As examples, phosphorothioate oligonucleotides may be synthesized by the method of Stein et al. (1988, Nucl. Acids Res. 16:3209), methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin et al., 1988, Proc. Natl. Acad. Sci. U.S.A. 85:7448-7451), etc.

While antisense nucleotides complementary to the CRCGL coding region ~~sequence could be used, those complementary to the transcribed untranslated region~~ are most preferred.

Potential antagonists according to the invention also include catalytic RNA, or a ribozyme (See, e.g., PCT International Publication WO 90/11364, published October 4, 1990; Sarver et al, Science 247:1222-1225 (1990). While ribozymes that cleave mRNA at site specific recognition sequences can be used to destroy CRCGCL mRNAs, the use of hammerhead ribozymes is preferred. Hammerhead ribozymes cleave mRNAs at locations dictated by flanking regions that form complementary base pairs with the target mRNA. The sole requirement is that the target mRNA have the following sequence of two bases: 5'-UG-3'. The construction and production of hammerhead ribozymes is well known in the art and is described more fully in Haseloff and Gerlach, Nature 334:585-591 (1988). There are numerous potential hammerhead ribozyme cleavage sites within the nucleotide sequence of CRCGCL (Figures 1A-B). Preferably, the ribozyme is engineered so that the cleavage recognition site is located near the 5' end of the CRCGCL mRNA; i.e., to increase

efficiency and minimize the intracellular accumulation of non-functional mRNA transcripts.

As in the antisense approach, the ribozymes of the invention can be composed of modified oligonucleotides (e.g. for improved stability, targeting, etc.) and should be delivered to cells which express CRCGCL in vivo. DNA constructs encoding the ribozyme may be introduced into the cell in the same manner as described above for the introduction of antisense encoding DNA. A preferred method of delivery involves using a DNA construct "encoding" the ribozyme under the control of a strong constitutive promoter, such as, for example, pol III or pol II promoter, so that transfected cells will produce sufficient quantities of the ribozyme to destroy endogenous CRCGCL messages and inhibit translation. Since ribozymes unlike antisense molecules, are catalytic, a lower intracellular concentration is required for efficiency.

Antagonist/agonist compounds may be employed to inhibit the cell growth and proliferation effects of the polypeptides of the present invention on neoplastic cells and tissues, i.e. stimulation of angiogenesis of tumors, and, therefore, retard or prevent abnormal cellular growth and proliferation, for example, in tumor formation or growth.

The antagonist/agonist may also be employed to prevent hyper-vascular diseases, and prevent the proliferation of epithelial lens cells after extracapsular cataract surgery. Prevention of the mitogenic activity of the polypeptides of the present invention may also be desirable in cases such as restenosis after balloon angioplasty.

The antagonist/agonist may also be employed to prevent the growth of scar tissue during wound healing.

The antagonist/agonist may also be employed to treat the diseases described herein.

#### **Other Activities**

The polypeptide of the present invention, as a result of the ability to stimulate vascular endothelial cell growth, may be employed in treatment for stimulating re-vascularization of ischemic tissues due to various disease conditions such as thrombosis, arteriosclerosis, and other cardiovascular conditions. These polypeptide

The polypeptide may also be employed for treating wounds due to injuries, burns, post-operative tissue repair, and ulcers since they are mitogenic to various cells of different origins, such as fibroblast cells and skeletal muscle cells, and therefore, facilitate the repair or replacement of damaged or diseased tissue.

The polypeptide of the present invention may be also be employed to prevent skin aging due to sunburn by stimulating keratinocyte growth.

20           The CRCGCL polypeptide may also be employed to maintain organs before transplantation or for supporting cell culture of primary tissues.

CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, may also increase or decrease the differentiation or proliferation of embryonic stem cells, besides, as discussed above, hematopoietic lineage.

CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, may also be used to modulate mammalian characteristics, such as body height, weight, hair color, eye color, skin, percentage of adipose tissue, pigmentation, size, and shape (e.g., cosmetic surgery). Similarly, CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, may be used to modulate mammalian metabolism affecting catabolism, anabolism, processing, utilization, and storage of energy.

CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, may be used to change a mammal's mental state or physical state by influencing biorhythms, cardiac rhythms, depression (including depressive disorders), tendency for violence, tolerance for pain, reproductive capabilities (preferably by Activin or Inhibin-like activity), hormonal or endocrine levels, appetite, libido, memory, stress, or other cognitive qualities.

CRCGCL polynucleotides or polypeptides, or agonists or antagonists of CRCGCL, may also be used as a food additive or preservative, such as to increase or decrease storage capabilities, fat content, lipid, protein, carbohydrate, vitamins, minerals, cofactors or other nutritional components.

The above-recited applications have uses in a wide variety of hosts. Such hosts include, but are not limited to, human, murine, rabbit, goat, guinea pig, camel, horse, mouse, rat, hamster, pig, micro-pig, chicken, goat, cow, sheep, dog, cat, non-human primate, and human. In specific embodiments, the host is a mouse, rabbit, goat, guinea pig, chicken, rat, hamster, pig, sheep, dog or cat. In preferred embodiments, the host is a mammal. In most preferred embodiments, the host is a human.

~~Having generally described the invention, the same will be more readily understood by reference to the following examples, which are provided by way of illustration and are not intended as limiting.~~

Having generally described the invention, the same will be more readily understood by reference to the following examples, which are provided by way of illustration and are not intended as limiting.

## Examples

### *Example 1: Isolation of the CRCGCL cDNA Clone From the Deposited Sample*

The cDNA for CRCGCL is inserted into the EcoRI/XhoI multiple cloning site of Uni-ZAP XR (Stratagene). Uni-ZAP XR contains an ampicillin resistance gene and may be transformed into E. coli strain DH10B, available from Life Technologies. (See, for instance, Gruber, C. E., et al., Focus 15:59- (1993).)



Two approaches can be used to isolate CRCGCL from the deposited sample. First, the deposited clone is transformed into a suitable host (such as XL-1 Blue (Stratagene)) using techniques known to those of skill in the art, such as those provided by the vector supplier or in related publications or patents. The transformants are plated on 1.5% agar plates (containing the appropriate selection agent, e.g., ampicillin) to a density of about 150 transformants (colonies) per plate. A single colony is then used to generate DNA using nucleic acid isolation techniques well known to those skilled in the art. (e.g., Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd Edit., (1989), Cold Spring Harbor Laboratory Press.)

Alternatively, two primers of 17-20 nucleotides derived from both ends of the SEQ ID NO:1 (i.e., within the region of SEQ ID NO:1 bounded by the 5' NT and the 3' NT of the clone) are synthesized and used to amplify the CRCGCL cDNA using the deposited cDNA plasmid as a template. The polymerase chain reaction is carried out under routine conditions, for instance, in 25 ul of reaction mixture with 0.5 ug of the above cDNA template. A convenient reaction mixture is 1.5-5 mM MgCl<sub>2</sub>, 0.01% (w/v) gelatin, 20 uM each of dATP, dCTP, dGTP, dTTP, 25 pmol of each primer and 0.25 Unit of Taq polymerase. Thirty five cycles of PCR (denaturation at 94 degree C for 1 min; annealing at 55 degree C for 1 min; elongation at 72 degree C for 1 min) are performed with a Perkin-Elmer Cetus automated thermal cycler. The amplified product is analyzed by agarose gel electrophoresis and the DNA band with expected molecular weight is excised and purified. The PCR product is verified to be the selected sequence by subcloning and sequencing the DNA product.

Several methods are available for the identification of the 5' or 3' non-coding portions of the CRCGCL gene which may not be present in the deposited clone. These methods include but are not limited to, filter probing, clone enrichment using specific probes, and protocols similar or identical to 5' and 3' "RACE" protocols which are well known in the art. For instance, a method similar to 5' RACE is available for generating the missing 5' end of a desired full-length transcript. (Fromont-Racine et al., Nucleic Acids Res. 21(7):1683-1684 (1993).)

Briefly, a specific RNA oligonucleotide is ligated to the 5' ends of a population of RNA presumably containing full-length gene RNA transcripts. A primer set containing a primer specific to the ligated RNA oligonucleotide and a primer specific to a known sequence of the CRCGCL gene of interest is used to PCR

amplify the 5' portion of the CRCGCL full-length gene. This amplified product may then be sequenced and used to generate the full length gene.

This above method starts with total RNA isolated from the desired source, although poly-A+ RNA can be used. The RNA preparation can then be treated with phosphatase if necessary to eliminate 5' phosphate groups on degraded or damaged RNA which may interfere with the later RNA ligase step. The phosphatase should then be inactivated and the RNA treated with tobacco acid pyrophosphatase in order to remove the cap structure present at the 5' ends of messenger RNAs. This reaction leaves a 5' phosphate group at the 5' end of the cap cleaved RNA which can then be ligated to an RNA oligonucleotide using T4 RNA ligase.

This modified RNA preparation is used as a template for first strand cDNA synthesis using a gene specific oligonucleotide. The first strand synthesis reaction is used as a template for PCR amplification of the desired 5' end using a primer specific to the ligated RNA oligonucleotide and a primer specific to the known sequence of the gene of interest. The resultant product is then sequenced and analyzed to confirm that the 5' end sequence belongs to the CRCGCL gene.

---

### *Example 2: Isolation of CRCGCL Genomic Clones*

A human genomic P1 library (Genomic Systems, Inc.) is screened by PCR using primers selected for the cDNA sequence corresponding to SEQ ID NO:1., according to the method described in Example 1. (See also, Sambrook.)

### *Example 3: Tissue Distribution of CRCGCL Polypeptides*

Tissue distribution of mRNA expression of CRCGCL is determined using protocols for Northern blot analysis, described by, among others, Sambrook et al. For example, a CRCGCL probe produced by the method described in Example 1 is labeled with P<sup>32</sup> using the rediprime™ DNA labeling system (Amersham Life Science), according to manufacturer's instructions. After labeling, the probe is purified using CHROMA SPIN-100™ column (Clontech Laboratories, Inc.), according to manufacturer's protocol number PT1200-1. The purified labeled probe is then used to examine various human tissues for mRNA expression.

Multiple Tissue Northern (MTN) blots containing various human tissues (H) or human immune system tissues (IM) (Clontech) are examined with the labeled probe using ExpressHyb™ hybridization solution (Clontech) according to manufacturer's protocol number PT1190-1. Following hybridization and washing, the blots are mounted and exposed to film at -70 degree C overnight, and the films developed according to standard procedures.

#### ***Example 4: Chromosomal Mapping of CRCGCL***

10 An oligonucleotide primer set is designed according to the sequence at the 5' end of SEQ ID NO:1. This primer preferably spans about 100 nucleotides. This primer set is then used in a polymerase chain reaction under the following set of conditions : 30 seconds, 95 degree C; 1 minute, 56 degree C; 1 minute, 70 degree C. This cycle is repeated 32 times followed by one 5 minute cycle at 70 degree C.

15 Human, mouse, and hamster DNA is used as template in addition to a somatic cell hybrid panel containing individual chromosomes or chromosome fragments (Bios, Inc). The reactions is analyzed on either 8% polyacrylamide gels or 3.5 % agarose gels. Chromosome mapping is determined by the presence of an approximately 100 bp PCR fragment in the particular somatic cell hybrid.

20

#### ***Example 5: Bacterial Expression of CRCGCL***

CRCGCL polynucleotide encoding a CRCGCL polypeptide invention is amplified using PCR oligonucleotide primers corresponding to the 5' and 3' ends of the DNA sequence, as outlined in Example 1, to synthesize insertion fragments. The primers used to amplify the cDNA insert should preferably contain restriction sites, such as BamHI and XbaI, at the 5' end of the primers in order to clone the amplified product into the expression vector. For example, BamHI and XbaI correspond to the restriction enzyme sites on the bacterial expression vector pQE-9. (Qiagen, Inc., Chatsworth, CA). This plasmid vector encodes antibiotic resistance (Amp<sup>r</sup>), a bacterial origin of replication (ori), an IPTG-regulatable promoter/operator (P/O), a ribosome binding site (RBS), a 6-histidine tag (6-His), and restriction enzyme cloning sites.

Specifically, to clone the CRCGCL protein in a bacterial vector, the 5' primer has the sequence 5' GTTAGGCCATGGGAGGAGCAGCAGAAGGA 3' (SEQ ID NO: 14) containing the Nco I restriction site followed a number of nucleotides of the amino terminal coding sequence of the CRCGCL sequence in SEQ ID NO:1. One of ordinary skill in the art would appreciate, of course, that the point in the protein coding sequence where the 5' primer begins may be varied to amplify a DNA segment encoding any desired portion of the complete CRCGCL protein shorter or longer than the the portion described above. The 3' primer has the sequence 5' GGTTAAAGATCTCAACGCCACGTAGGAGCGGTC 3' (SEQ ID NO: 15) containing the BglII restriction site followed by a number nucleotides complementary to the 3' end of the coding sequence of the CRCGCL DNA sequence of SEQ ID NO:1.

The pQE-9 vector is digested with BamHI and XbaI and the amplified fragment is ligated into the pQE-9 vector maintaining the reading frame initiated at the bacterial RBS. The ligation mixture is then used to transform the E. coli strain M15/rep4 (Qiagen, Inc.) which contains multiple copies of the plasmid pREP4, which expresses the lacI repressor and also confers kanamycin resistance (Kan<sup>r</sup>). ~~Transformants are identified by their ability to grow on LB plates and~~ ampicillin/kanamycin resistant colonies are selected. Plasmid DNA is isolated and confirmed by restriction analysis.

Clones containing the desired constructs are grown overnight (O/N) in liquid culture in LB media supplemented with both Amp (100 ug/ml) and Kan (25 ug/ml). The O/N culture is used to inoculate a large culture at a ratio of 1:100 to 1:250. The cells are grown to an optical density 600 (O.D.<sup>600</sup>) of between 0.4 and 0.6. IPTG (Isopropyl-B-D-thiogalacto pyranoside) is then added to a final concentration of 1 mM. IPTG induces by inactivating the lacI repressor, clearing the P/O leading to increased gene expression.

Cells are grown for an extra 3 to 4 hours. Cells are then harvested by centrifugation (20 mins at 6000Xg). The cell pellet is solubilized in the chaotropic agent 6 Molar Guanidine HCl by stirring for 3-4 hours at 4 degree C. The cell debris is removed by centrifugation, and the supernatant containing the polypeptide is loaded onto a nickel-nitrilo-tri-acetic acid ("Ni-NTA") affinity resin column (available from

QIAGEN, Inc., supra). Proteins with a 6 x His tag bind to the Ni-NTA resin with high affinity and can be purified in a simple one-step procedure (for details see: The QIAexpressionist (1995) QIAGEN, Inc., supra).

5 Briefly, the supernatant is loaded onto the column in 6 M guanidine-HCl, pH 8, the column is first washed with 10 volumes of 6 M guanidine-HCl, pH 8, then washed with 10 volumes of 6 M guanidine-HCl pH 6, and finally the polypeptide is eluted with 6 M guanidine-HCl, pH 5.

10 The purified CRCGCL protein is then renatured by dialyzing it against phosphate-buffered saline (PBS) or 50 mM Na-acetate, pH 6 buffer plus 200 mM NaCl. Alternatively, the CRCGCL protein can be successfully refolded while immobilized on the Ni-NTA column. The recommended conditions are as follows: renature using a linear 6M-1M urea gradient in 500 mM NaCl, 20% glycerol, 20 mM Tris/HCl pH 7.4, containing protease inhibitors. The renaturation should be performed over a period of 1.5 hours or more. After renaturation the proteins are  
15 eluted by the addition of 250 mM imidazole. Imidazole is removed by a final dialyzing step against PBS or 50 mM sodium acetate pH 6 buffer plus 200 mM NaCl. The purified CRCGCL protein is stored at 4 degree C or frozen at -80 degree C.

In addition to the above expression vector, the present invention further  
includes an expression vector comprising phage operator and promoter elements  
20 operatively linked to a CRCGCL polynucleotide, called pHE4a. (ATCC Accession Number 209645, deposited February 25, 1998.) This vector contains: 1) a neomycinphosphotransferase gene as a selection marker, 2) an E. coli origin of replication, 3) a T5 phage promoter sequence, 4) two lac operator sequences, 5) a Shine-Delgarno sequence, and 6) the lactose operon repressor gene (lacIq). The  
25 origin of replication (oriC) is derived from pUC19 (LTI, Gaithersburg, MD). The promoter sequence and operator sequences are made synthetically.

30 DNA can be inserted into the pHEa by restricting the vector with NdeI and XbaI, BamHI, XhoI, or Asp718, running the restricted product on a gel, and isolating the larger fragment (the stuffer fragment should be about 310 base pairs). The DNA insert is generated according to the PCR protocol described in Example 1, using PCR primers having restriction sites for NdeI (5' primer) and XbaI, BamHI, XhoI, or Asp718 (3' primer). The PCR insert is gel purified and restricted with compatible enzymes. The insert and vector are ligated according to standard protocols.

0376430-0349260

The engineered vector could easily be substituted in the above protocol to express protein in a bacterial system. More preferably, the bacterial expression vector, pQE60 can also be used to express CRCGCL.

5

***Example 6: Purification of CRCGCL Polypeptide  
from an Inclusion Body***

10 The following alternative method can be used to purify CRCGCL polypeptide expressed in E coli when it is present in the form of inclusion bodies. Unless otherwise specified, all of the following steps are conducted at 4-10 degree C.

15 Upon completion of the production phase of the E. coli fermentation, the cell culture is cooled to 4-10 degree C and the cells harvested by continuous centrifugation at 15,000 rpm (Heraeus Sepatech). On the basis of the expected yield of protein per unit weight of cell paste and the amount of purified protein required, an appropriate amount of cell paste, by weight, is suspended in a buffer solution containing 100 mM Tris, 50 mM EDTA, pH 7.4. The cells are dispersed to a homogeneous suspension using a high shear mixer.

20 The cells are then lysed by passing the solution through a microfluidizer (Microfluidics, Corp. or APV Gaulin, Inc.) twice at 4000-6000 psi. The homogenate is then mixed with NaCl solution to a final concentration of 0.5 M NaCl, followed by centrifugation at 7000 xg for 15 min. The resultant pellet is washed again using 0.5M NaCl, 100 mM Tris, 50 mM EDTA, pH 7.4.

25 The resulting washed inclusion bodies are solubilized with 1.5 M guanidine hydrochloride (GuHCl) for 2-4 hours. After 7000 xg centrifugation for 15 min., the pellet is discarded and the polypeptide containing supernatant is incubated at 4 degree C overnight to allow further GuHCl extraction.

30 Following high speed centrifugation (30,000 xg) to remove insoluble particles, the GuHCl solubilized protein is refolded by quickly mixing the GuHCl extract with 20 volumes of buffer containing 50 mM sodium, pH 4.5, 150 mM NaCl, 2 mM EDTA by vigorous stirring. The refolded diluted protein solution is kept at 4 degree C without mixing for 12 hours prior to further purification steps.

To clarify the refolded polypeptide solution, a previously prepared tangential filtration unit equipped with 0.16 um membrane filter with appropriate surface area

05375430-03459

(e.g., Filtron), equilibrated with 40 mM sodium acetate, pH 6.0 is employed. The filtered sample is loaded onto a cation exchange resin (e.g., Poros HS-50, Perseptive Biosystems). The column is washed with 40 mM sodium acetate, pH 6.0 and eluted with 250 mM, 500 mM, 1000 mM, and 1500 mM NaCl in the same buffer, in a stepwise manner. The absorbance at 280 nm of the effluent is continuously monitored. Fractions are collected and further analyzed by SDS-PAGE.

Fractions containing the CRCGCL polypeptide are then pooled and mixed with 4 volumes of water. The diluted sample is then loaded onto a previously prepared set of tandem columns of strong anion (Poros HQ-50, Perseptive Biosystems) and weak anion (Poros CM-20, Perseptive Biosystems) exchange resins. The columns are equilibrated with 40 mM sodium acetate, pH 6.0. Both columns are washed with 40 mM sodium acetate, pH 6.0, 200 mM NaCl. The CM-20 column is then eluted using a 10 column volume linear gradient ranging from 0.2 M NaCl, 50 mM sodium acetate, pH 6.0 to 1.0 M NaCl, 50 mM sodium acetate, pH 6.5. Fractions are collected under constant  $A_{280}$  monitoring of the effluent. Fractions containing the polypeptide (determined, for instance, by 16% SDS-PAGE) are then pooled.

The resultant CRCGCL polypeptide should exhibit greater than 95% purity after the above refolding and purification steps. No major contaminant bands should be observed from Commassie blue stained 16% SDS-PAGE gel when 5 ug of purified protein is loaded. The purified CRCGCL protein can also be tested for endotoxin/LPS contamination, and typically the LPS content is less than 0.1 ng/ml according to LAL assays.

### ***Example 7: Cloning and Expression of CRCGCL in a Baculovirus Expression System***

In this example, the plasmid shuttle vector pA2 is used to insert CRCGCL polynucleotide into a baculovirus to express CRCGCL. This expression vector contains the strong polyhedrin promoter of the Autographa californica nuclear polyhedrosis virus (AcMNPV) followed by convenient restriction sites such as BamHI, Xba I and Asp718. The polyadenylation site of the simian virus 40 ("SV40") is used for efficient polyadenylation. For easy selection of recombinant virus, the plasmid contains the beta-galactosidase gene from E. coli under control of a weak

Drosophila promoter in the same orientation, followed by the polyadenylation signal of the polyhedrin gene. The inserted genes are flanked on both sides by viral sequences for cell-mediated homologous recombination with wild-type viral DNA to generate a viable virus that express the cloned CRCGCL polynucleotide.

5 Many other baculovirus vectors can be used in place of the vector above, such as pAc373, pVL941, and pAcIM1, as one skilled in the art would readily appreciate, as long as the construct provides appropriately located signals for transcription, translation, secretion and the like, including a signal peptide and an in-frame AUG as required. Such vectors are described, for instance, in Luckow et al., Virology 170:31-10 39 (1989).

Specifically, the CRCGCL cDNA sequence contained in the deposited clone, including the AUG initiation codon and any naturally associated leader sequence, is amplified using the PCR protocol described in Example 1. If the naturally occurring signal sequence is used to produce the secreted protein, the pA2 vector does not need 15 a second signal peptide. Alternatively, the vector can be modified (pA2 GP) to include a baculovirus leader sequence, using the standard methods described in Summers et al., "A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures," Texas Agricultural Experimental Station Bulletin No. 1555 (1987).

20 More specifically, the cDNA sequence encoding the full length CRCGCL protein in the deposited clone, including the AUG initiation codon and the naturally associated leader sequence shown in SEQ ID NO:1, is amplified using PCR oligonucleotide primers corresponding to the 5' and 3' sequences of the gene. For example, the 5' primer could have the sequence 5' 25 CCGGTTAGATCTGCCATCATGGCTTTGGGGCAAGGAGG 3' (SEQ ID NO: 16) containing the BglII restriction enzyme site, an efficient signal for initiation of translation in eukaryotic cells (Kozak, M., J. Mol. Biol. 196:947-950 (1987)), followed by a number of nucleotides of the sequence of the complete CRCGCL protein shown in Figures 1A-1B. Alternatively, the 5' primer: 5' 30 CCGGTTAGATCTGCCATCATGGGGCGGCTGGTTCTG 3' (SEQ ID NO:28), also having a Bgl II restriction site could also be used. The 3' primer has the sequence 5' CCGGTTTCTAGATCACAAGGCCACGTAGGAGCGGTC 3' (SEQ ID NO: 17)



containing the XbaI restriction site followed by a number of nucleotides complementary to the 3' noncoding sequence in Figures 1A-1B.

The amplified fragment is isolated from a 1% agarose gel using a commercially available kit ("GeneClean," BIO 101 Inc., La Jolla, Ca.). The fragment  
5 then is digested with appropriate restriction enzymes and again purified on a 1% agarose gel.

The plasmid is digested with the corresponding restriction enzymes and optionally, can be dephosphorylated using calf intestinal phosphatase, using routine procedures known in the art. The DNA is then isolated from a 1% agarose gel using a  
10 commercially available kit ("GeneClean" BIO 101 Inc., La Jolla, Ca.).

The fragment and the dephosphorylated plasmid are ligated together with T4 DNA ligase. E. coli HB101 or other suitable E. coli hosts such as XL-1 Blue (Stratagene Cloning Systems, La Jolla, CA) cells are transformed with the ligation mixture and spread on culture plates. Bacteria containing the plasmid are identified  
15 by digesting DNA from individual colonies and analyzing the digestion product by gel electrophoresis. The sequence of the cloned fragment is confirmed by DNA sequencing.

~~Five ug of a plasmid containing the polynucleotide is co-transfected with 1.0~~  
20 ug of a commercially available linearized baculovirus DNA ("BaculoGold™ baculovirus DNA", Pharmingen, San Diego, CA), using the lipofection method described by Felgner et al., Proc. Natl. Acad. Sci. USA 84:7413-7417 (1987). One ug of BaculoGold™ virus DNA and 5 ug of the plasmid are mixed in a sterile well of a microtiter plate containing 50 ul of serum-free Grace's medium (Life Technologies Inc., Gaithersburg, MD). Afterwards, 10 ul Lipofectin plus 90 ul Grace's medium are  
25 added, mixed and incubated for 15 minutes at room temperature. Then the transfection mixture is added drop-wise to Sf9 insect cells (ATCC CRL 1711) seeded in a 35 mm tissue culture plate with 1 ml Grace's medium without serum. The plate is then incubated for 5 hours at 27 degrees C. The transfection solution is then removed from the plate and 1 ml of Grace's insect medium supplemented with 10% fetal calf  
30 serum is added. Cultivation is then continued at 27 degrees C for four days.

After four days the supernatant is collected and a plaque assay is performed, as described by Summers and Smith, supra. An agarose gel with "Blue Gal" (Life Technologies Inc., Gaithersburg) is used to allow easy identification and isolation of

00375430-001000

gal-expressing clones, which produce blue-stained plaques. (A detailed description of a "plaque assay" of this type can also be found in the user's guide for insect cell culture and baculovirology distributed by Life Technologies Inc., Gaithersburg, page 9-10.) After appropriate incubation, blue stained plaques are picked with the tip of a micropipettor (e.g., Eppendorf). The agar containing the recombinant viruses is then resuspended in a microcentrifuge tube containing 200 ul of Grace's medium and the suspension containing the recombinant baculovirus is used to infect Sf9 cells seeded in 35 mm dishes. Four days later the supernatants of these culture dishes are harvested and then they are stored at 4 degree C.

To verify the expression of the polypeptide, Sf9 cells are grown in Grace's medium supplemented with 10% heat-inactivated FBS. The cells are infected with the recombinant baculovirus containing the polynucleotide at a multiplicity of infection ("MOI") of about 2. If radiolabeled proteins are desired, 6 hours later the medium is removed and is replaced with SF900 II medium minus methionine and cysteine (available from Life Technologies Inc., Rockville, MD). After 42 hours, 5 uCi of <sup>35</sup>S-methionine and 5 uCi <sup>35</sup>S-cysteine (available from Amersham) are added. The cells are further incubated for 16 hours and then are harvested by centrifugation. ~~The proteins in the supernatant as well as the intracellular proteins are analyzed by~~ SDS-PAGE followed by autoradiography (if radiolabeled).

Microsequencing of the amino acid sequence of the amino terminus of purified protein may be used to determine the amino terminal sequence of the produced CRCGCL protein.

### ***Example 8: Expression of CRCGCL in Mammalian Cells***

CRCGCL polypeptide can be expressed in a mammalian cell. A typical mammalian expression vector contains a promoter element, which mediates the initiation of transcription of mRNA, a protein coding sequence, and signals required for the termination of transcription and polyadenylation of the transcript. Additional elements include enhancers, Kozak sequences and intervening sequences flanked by donor and acceptor sites for RNA splicing. Highly efficient transcription is achieved with the early and late promoters from SV40, the long terminal repeats (LTRs) from Retroviruses, e.g., RSV, HTLVI, HIVI and the early promoter of the cytomegalovirus

(CMV). However, cellular elements can also be used (e.g., the human actin promoter).

Suitable expression vectors for use in practicing the present invention include, for example, vectors such as pSVL and pMSG (Pharmacia, Uppsala, Sweden), pRSVcat (ATCC 37152), pSV2DHFR (ATCC 37146), pBC12MI (ATCC 67109), pCMVSPORT 2.0, and pCMVSPORT 3.0. Mammalian host cells that could be used include, human Hela, 293, H9 and Jurkat cells, mouse NIH3T3 and C127 cells, Cos 1, Cos 7 and CV1, quail QC1-3 cells, mouse L cells and Chinese hamster ovary (CHO) cells.

Alternatively, CRCGCL polypeptide can be expressed in stable cell lines containing the CRCGCL polynucleotide integrated into a chromosome. The co-transfection with a selectable marker such as DHFR, gpt, neomycin, hygromycin allows the identification and isolation of the transfected cells.

The transfected CRCGCL gene can also be amplified to express large amounts of the encoded protein. The DHFR (dihydrofolate reductase) marker is useful in developing cell lines that carry several hundred or even several thousand copies of the gene of interest. (See, e.g., Alt, F. W., et al., J. Biol. Chem. 253:1357-1370 (1978); Hamlin, J. L. and Ma, C., Biochem. et Biophys. Acta, 1097:107-143 (1990); Page, M. J. and Sydenham, M. A., Biotechnology 9:64-68 (1991).) Another useful selection marker is the enzyme glutamine synthase (GS) (Murphy et al., Biochem J. 227:277-279 (1991); Bebbington et al., Bio/Technology 10:169-175 (1992). Using these markers, the mammalian cells are grown in selective medium and the cells with the highest resistance are selected. These cell lines contain the amplified gene(s) integrated into a chromosome. Chinese hamster ovary (CHO) and NSO cells are often used for the production of proteins.

Derivatives of the plasmid pSV2-DHFR (ATCC Accession No. 37146), the expression vectors pC4 (ATCC Accession No. 209646) and pC6 (ATCC Accession No. 209647) contain the strong promoter (LTR) of the Rous Sarcoma Virus (Cullen et al., Molecular and Cellular Biology, 438-447 (March, 1985)) plus a fragment of the CMV-enhancer (Boshart et al., Cell 41:521-530 (1985).) Multiple cloning sites, e.g., with the restriction enzyme cleavage sites BamHI, XbaI and Asp718, facilitate the cloning of CRCGCL. The vectors also contain the 3' intron, the polyadenylation and

0349260

termination signal of the rat preproinsulin gene, and the mouse DHFR gene under control of the SV40 early promoter.

Specifically, the plasmid pC6 or pC4 is digested with appropriate restriction enzymes and then dephosphorylated using calf intestinal phosphates by procedures known in the art. The vector is then isolated from a 1% agarose gel. Also preferred is the pcDNA3 vector (Life Technologies).

If a naturally occurring signal sequence is used to produce a secreted protein, the vector does not need a second signal peptide. Alternatively, if a naturally occurring signal sequence is not used, the vector can be modified to include a heterologous signal sequence in an effort to secrete the protein from the cell. (See, e.g., WO 96/34891.)

The amplified fragment is then digested with the same restriction enzyme and purified on a 1% agarose gel using a commercially available kit ("GeneClean," BIO 101 Inc., La Jolla, Ca.). The isolated fragment and the dephosphorylated vector are then ligated with T4 DNA ligase. E. coli HB101 or XL-1 Blue cells are then transformed and bacteria are identified that contain the fragment inserted into plasmid pC6 or pC4 using, for instance, restriction enzyme analysis.

For example, a soluble CRCGCL polypeptide, such as amino acids Met 1 to Lys 231, could also be expressed. A 5' primer: 5'

CCGGTTAGATCTGCCATCATGGGGCGGCTGGTTCTG 3' (SEQ ID NO:28), having a Bgl II restriction site and a 3' primer: 5'

GGCCGGTCTAGATTATTTGGACAGCTTTGGTTTG 3' (SEQ ID NO:31) could be used to PCR amino acids Met 1 to Lys 231. The amplified product could be inserted into a mammalian expression vector, such as pC4 or pC6.

Chinese hamster ovary cells lacking an active DHFR gene is used for transfection. Five µg of the expression plasmid pC6 or pC4 is cotransfected with 0.5 µg of the plasmid pSVneo using lipofectin (Felgner et al., supra). The plasmid pSV2-neo contains a dominant selectable marker, the neo gene from Tn5 encoding an enzyme that confers resistance to a group of antibiotics including G418. The cells are seeded in alpha minus MEM supplemented with 1 mg/ml G418. After 2 days, the cells are trypsinized and seeded in hybridoma cloning plates (Greiner, Germany) in alpha minus MEM supplemented with 10, 25, or 50 ng/ml of metothrexate plus 1 mg/ml G418. After about 10-14 days single clones are trypsinized and then seeded in

037430-037430

5 The same procedure is repeated until clones are obtained which grow at a concentration of 100 - 200 uM. Expression of CRCGCL is analyzed, for instance, by SDS-PAGE and Western blot or by reversed phase HPLC analysis.

## 10

20

25

As a means of exemplifying but not limiting the present invention, the polynucleotide encoding the CRCGCL polypeptide fragment I-35 to F-276 is

amplified and cloned as follows: A 5' primer is generated comprising a restriction enzyme site followed by an initiation codon in frame with the polynucleotide sequence encoding the N-terminal portion of the polypeptide fragment beginning with I-35. A complementary 3' primer is generated comprising a restriction enzyme site followed by a stop codon in frame with the polynucleotide sequence encoding C-terminal portion of the CRCGCL polypeptide fragment ending with F-276.

The amplified polynucleotide fragment and the expression vector are digested with restriction enzymes which recognize the sites in the primers. The digested polynucleotides are then ligated together. The CRCGCL polynucleotide fragment is inserted into the restricted expression vector, preferably in a manner which places the CRCGCL polypeptide fragment coding region downstream from the promoter. The ligation mixture is transformed into competent E. coli cells using standard procedures and as described in the Examples herein. Plasmid DNA is isolated from resistant colonies and the identity of the cloned DNA confirmed by restriction analysis, PCR and DNA sequencing.

### ***Example 10: Protein Fusions of CRCGCL***

CRCGCL polypeptides are preferably fused to other proteins. These fusion proteins can be used for a variety of applications. For example, fusion of CRCGCL polypeptides to His-tag, HA-tag, protein A, IgG domains, and maltose binding protein facilitates purification. (See Example 5; see also EP A 394,827; Traunecker, et al., Nature 331:84-86 (1988).) Similarly, fusion to IgG-1, IgG-3, and albumin increases the halflife time in vivo. Nuclear localization signals fused to CRCGCL polypeptides can target the protein to a specific subcellular localization, while covalent heterodimer or homodimers can increase or decrease the activity of a fusion protein. Fusion proteins can also create chimeric molecules having more than one function. Finally, fusion proteins can increase solubility and/or stability of the fused protein compared to the non-fused protein. All of the types of fusion proteins described above can be made by modifying the following protocol, which outlines the fusion of a polypeptide to an IgG molecule, or the protocol described in Example 5.

Briefly, the human Fc portion of the IgG molecule can be PCR amplified, using primers that span the 5' and 3' ends of the sequence described below. These

00000-00000-00000

primers also should have convenient restriction enzyme sites that will facilitate cloning into an expression vector, preferably a mammalian expression vector.

For example, if pC4 (Accession No. 209646) is used, the human Fc portion can be ligated into the BamHI cloning site. Note that the 3' BamHI site should be destroyed. Next, the vector containing the human Fc portion is re-restricted with BamHI, linearizing the vector, and CRCGCL polynucleotide, isolated by the PCR protocol described in Example 1, is ligated into this BamHI site.

Alternatively, a soluble CRCGCL polypeptide, such as amino acids Met 1 to Lys 231, could also be fused to the Fc portion. For Example, a 5' primer: 5' CCGGTTAGATCTGCCATCATGGGGCGGCTGGTTCTG 3' (SEQ ID NO:28), having a Bgl II restriction site and a 3' primer: 5' GGCCGGTCTAGATTTGGACAGCTTTGGTTTG 3' (SEQ ID NO:29) could be used to PCR amino acids Met 1 to Lys 231. The amplified product could be fused to Fc to produce a Fc fusion protein, as set forth above, and ligated to the pC4 vector for mammalian expression or pA2 for baculovirus expression.

In either case, note that the polynucleotide is cloned without a stop codon, otherwise a fusion protein will not be produced. Moreover, if the naturally occurring signal sequence is used to produce the secreted protein, pC4 does not need a second signal peptide. Alternatively, if the naturally occurring signal sequence is not used, the vector can be modified to include a heterologous signal sequence. (See, e.g., WO 96/34891.)

The Fc fusions described above could also be inserted into the pA2 vector to express in Baculovirus systems, as set forth in Example 7, using techniques known in the art and described herein.

#### Human IgG Fc region:

```
GGGATCCGGAGCCCAAATCTTCTGACAAAACTCACACATGCCCACCGTGCCCAGCACCTGAATTCGAGG
GTGCACCGTCAGTCTTCTCTTCCCCCAAAACCCAAGGACACCCTCATGATCTCCCGGACTCCTGAGG
TCACATGCGTGGTGGTGGACGTAAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCG
TGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAGCG
TCCTCACCGTCTCTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCC
TCCCAACCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCC
TGCCCCCATCCCGGGATGAGCTGACCAAGAACCAGGTGACCTGACCTGCCTGGTCAAAGGCTTCTATC
CAAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACACTACAAGACCACGCCCTCCCG
TGCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGG
```

GGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCC  
TGTCTCCGGGTAAATGAGTGCGACGGCCGCGACTCTAGAGGAT (SEQ ID NO:4)

### ***Example 11: Production of an Antibody***

5

#### **a) Hybridoma Technology**

The antibodies of the present invention can be prepared by a variety of methods. (See, Current Protocols, Chapter 2.) As one example of such methods, cells  
10 expressing CRCGCL is administered to an animal to induce the production of sera containing polyclonal antibodies. In a preferred method, a preparation of CRCGCL protein is prepared and purified to render it substantially free of natural contaminants. Such a preparation is then introduced into an animal in order to produce polyclonal antisera of greater specific activity.

15 In the most preferred method, the antibodies of the present invention are monoclonal antibodies (or protein binding fragments thereof). Such monoclonal antibodies can be prepared using hybridoma technology. (Köhler et al., Nature 256:495 (1975); Köhler et al., Eur. J. Immunol. 6:511 (1976); Köhler et al., Eur. J. Immunol. 6:292 (1976); Hammerling et al., in: Monoclonal Antibodies and T-Cell  
20 Hybridomas, Elsevier, N.Y., pp. 563-681 (1981).) In general, such procedures involve immunizing an animal (preferably a mouse) with CRCGCL polypeptide or, more preferably, with a secreted CRCGCL polypeptide-expressing cell. Such cells may be cultured in any suitable tissue culture medium; however, it is preferable to culture cells in Earle's modified Eagle's medium supplemented with 10% fetal bovine  
25 serum (inactivated at about 56 degree C), and supplemented with about 10 g/l of nonessential amino acids, about 1,000 U/ml of penicillin, and about 100 ug/ml of streptomycin.

The splenocytes of such mice are extracted and fused with a suitable myeloma cell line. Any suitable myeloma cell line may be employed in accordance with the  
30 present invention; however, it is preferable to employ the parent myeloma cell line (SP2O), available from the ATCC. After fusion, the resulting hybridoma cells are selectively maintained in HAT medium, and then cloned by limiting dilution as described by Wands et al. (Gastroenterology 80:225-232 (1981).) The hybridoma

0036430-0019660



cells obtained through such a selection are then assayed to identify clones which secrete antibodies capable of binding the CRCGCL polypeptide.

Alternatively, additional antibodies capable of binding to CRCGCL polypeptide can be produced in a two-step procedure using anti-idiotypic antibodies.

5 Such a method makes use of the fact that antibodies are themselves antigens, and therefore, it is possible to obtain an antibody which binds to a second antibody. In accordance with this method, protein specific antibodies are used to immunize an animal, preferably a mouse. The splenocytes of such an animal are then used to produce hybridoma cells, and the hybridoma cells are screened to identify clones  
10 which produce an antibody whose ability to bind to the CRCGCL protein-specific antibody can be blocked by CRCGCL. Such antibodies comprise anti-idiotypic antibodies to the CRCGCL protein-specific antibody and can be used to immunize an animal to induce formation of further CRCGCL protein-specific antibodies.

It will be appreciated that Fab and F(ab')<sub>2</sub> and other fragments of the  
15 antibodies of the present invention may be used according to the methods disclosed herein. Such fragments are typically produced by proteolytic cleavage, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')<sub>2</sub> fragments). ~~Alternatively, secreted CRCGCL protein-binding fragments can be~~ produced through the application of recombinant DNA technology or through  
20 synthetic chemistry.

For in vivo use of antibodies in humans, it may be preferable to use "humanized" chimeric monoclonal antibodies. Such antibodies can be produced using genetic constructs derived from hybridoma cells producing the monoclonal antibodies described above. Methods for producing chimeric antibodies are known in  
25 the art. (See, for review, Morrison, Science 229:1202 (1985); Oi et al., BioTechniques 4:214 (1986); Cabilly et al., U.S. Patent No. 4,816,567; Taniguchi et al., EP 171496; Morrison et al., EP 173494; Neuberger et al., WO 8601533; Robinson et al., WO 8702671; Boulianne et al., Nature 312:643 (1984); Neuberger et al., Nature 314:268 (1985).)

**b) Isolation of antibody fragments directed against CRCGCL from a library of scFvs.**

Naturally occurring V-genes isolated from human PBLs are constructed into a large library of antibody fragments which contain reactivities against CRCGCL to which the donor may or may not have been exposed (see e.g., U.S. Patent 5,885,793 incorporated herein in its entirety by reference).

**Rescue of the Library.** A library of scFvs is constructed from the RNA of human PBLs as described in WO92/01047. To rescue phage displaying antibody fragments, approximately  $10^9$  E. coli harbouring the phagemid are used to inoculate 50 ml of 2xTY containing 1% glucose and 100 ug/ml of ampicillin (2xTY-AMP-GLU) and grown to an O.D. of 0.8 with shaking. Five ml of this culture is used to inoculate 50 ml of 2xTY-AMP-GLU,  $2 \times 10^8$  TU of delta gene 3 helper (M13 delta gene III, see WO92/01047) are added and the culture incubated at 37 degree C for 45 minutes without shaking and then at 37 degree C for 45 minutes with shaking. The culture is centrifuged at 4000 r.p.m. for 10 min. and the pellet resuspended in 2 liters of 2xTY containing 100 ug/ml ampicillin and 50 ug/ml kanamycin and grown overnight. Phage are prepared as described in WO92/01047.

M13 delta gene III is prepared as follows: M13 delta gene III helper phage does not encode gene III protein, hence the phage(mid) displaying antibody fragments have a greater avidity of binding to antigen. Infectious M13 delta gene III particles are made by growing the helper phage in cells harbouring a pUC19 derivative supplying the wild type gene III protein during phage morphogenesis. The culture is incubated for 1 hour at 37 degree C without shaking and then for a further hour at 37 degree C with shaking. Cells are spun down (IEC-Centra 8, 4000 revs/min for 10 min), resuspended in 300 ml 2xTY broth containing 100 ug ampicillin/ml and 25 ug kanamycin/ml (2xTY-AMP-KAN) and grown overnight, shaking at 37° C. Phage particles are purified and concentrated from the culture medium by two PEG-precipitations (Sambrook et al., 1990), resuspended in 2 ml PBS and passed through a 0.45 um filter (Minisart NML; Sartorius) to give a final concentration of approximately  $10^{13}$  transducing units/ml (ampicillin-resistant clones).

**Panning of the Library.** Immuntubes (Nunc) are coated overnight in PBS with 4 ml of either 100 ug/ml or 10 ug/ml of a polypeptide of the present invention. Tubes are blocked with 2% Marvel-PBS for 2 hours at 37 degree C and then washed 3

times in PBS. Approximately  $10^{13}$  TU of phage is applied to the tube and incubated for 30 minutes at room temperature tumbling on an over and under turntable and then left to stand for another 1.5 hours. Tubes are washed 10 times with PBS 0.1% Tween-20 and 10 times with PBS. Phage are eluted by adding 1 ml of 100 mM triethylamine and rotating 15 minutes on an under and over turntable after which the solution is immediately neutralized with 0.5 ml of 1.0M Tris-HCl, pH 7.4. Phage are then used to infect 10 ml of mid-log E. coli TG1 by incubating eluted phage with bacteria for 30 minutes at 37 degree C. The E. coli are then plated on TYE plates containing 1% glucose and 100 ug/ml ampicillin. The resulting bacterial library is then rescued with delta gene 3 helper phage as described above to prepare phage for a subsequent round of selection. This process is then repeated for a total of 4 rounds of affinity purification with tube-washing increased to 20 times with PBS, 0.1% Tween-20 and 20 times with PBS for rounds 3 and 4.

**Characterization of Binders.** Eluted phage from the 3rd and 4th rounds of selection are used to infect E. coli HB 2151 and soluble scFv is produced (Marks, et al., 1991) from single colonies for assay. ELISAs are performed with microtitre plates coated with either 10 pg/ml of the polypeptide of the present invention in 50 mM bicarbonate pH 9.6. Clones positive in ELISA are further characterized by PCR fingerprinting (see e.g., WO92/01047) and then by sequencing.

### ***Example 12: Production Of CRCGCL Protein For High-Throughput Screening Assays***

The following protocol produces a supernatant containing CRCGCL polypeptide to be tested. This supernatant can then be used in the Screening Assays described in Examples 14-21.

First, dilute Poly-D-Lysine (644 587 Boehringer-Mannheim) stock solution (1mg/ml in PBS) 1:20 in PBS (w/o calcium or magnesium 17-516F Biowhittaker) for a working solution of 50ug/ml. Add 200 ul of this solution to each well (24 well plates) and incubate at RT for 20 minutes. Be sure to distribute the solution over each well (note: a 12-channel pipetter may be used with tips on every other channel). Aspirate off the Poly-D-Lysine solution and rinse with 1ml PBS (Phosphate Buffered

Saline). The PBS should remain in the well until just prior to plating the cells and plates may be poly-lysine coated in advance for up to two weeks.

Plate 293T cells (do not carry cells past P+20) at  $2 \times 10^5$  cells/well in .5ml DMEM(Dulbecco's Modified Eagle Medium)(with 4.5 G/L glucose and L-glutamine (12-604F Biowhittaker))/10% heat inactivated FBS(14-503F Biowhittaker)/1x Penstrep(17-602E Biowhittaker). Let the cells grow overnight.

The next day, mix together in a sterile solution basin: 300 ul Lipofectamine (18324-012 Gibco/BRL) and 5ml Optimem I (31985070 Gibco/BRL)/96-well plate. With a small volume multi-channel pipetter, aliquot approximately 2ug of an expression vector containing a polynucleotide insert, produced by the methods described in Examples 8-10, into an appropriately labeled 96-well round bottom plate. With a multi-channel pipetter, add 50ul of the Lipofectamine/Optimem I mixture to each well. Pipette up and down gently to mix. Incubate at RT 15-45 minutes. After about 20 minutes, use a multi-channel pipetter to add 150ul Optimem I to each well. As a control, one plate of vector DNA lacking an insert should be transfected with each set of transfections.

Preferably, the transfection should be performed by tag-teaming the following tasks. By tag-teaming, hands on time is cut in half, and the cells do not spend too much time on PBS. First, person A aspirates off the media from four 24-well plates of cells, and then person B rinses each well with .5-1ml PBS. Person A then aspirates off PBS rinse, and person B, using a 12-channel pipetter with tips on every other channel, adds the 200ul of DNA/Lipofectamine/Optimem I complex to the odd wells first, then to the even wells, to each row on the 24-well plates. Incubate at 37 degree C for 6 hours.

While cells are incubating, prepare appropriate media, either 1%BSA in DMEM with 1x penstrep, or HGS CHO-5 media (116.6 mg/L of  $\text{CaCl}_2$  (anhyd); 0.00130 mg/L  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ ; 0.050 mg/L of  $\text{Fe}(\text{NO}_3)_3 \cdot 9\text{H}_2\text{O}$ ; 0.417 mg/L of  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ ; 311.80 mg/L of Kcl; 28.64 mg/L of  $\text{MgCl}_2$ ; 48.84 mg/L of  $\text{MgSO}_4$ ; 6995.50 mg/L of NaCl; 2400.0 mg/L of  $\text{NaHCO}_3$ ; 62.50 mg/L of  $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$ ; 71.02 mg/L of  $\text{Na}_2\text{HPO}_4$ ; .4320 mg/L of  $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ ; .002 mg/L of Arachidonic Acid ; 1.022 mg/L of Cholesterol; .070 mg/L of DL-alpha-Tocopherol-Acetate; 0.0520 mg/L of Linoleic Acid; 0.010 mg/L of Linolenic Acid; 0.010 mg/L of Myristic

0927430 061300

- Acid; 0.010 mg/L of Oleic Acid; 0.010 mg/L of Palmitric Acid; 0.010 mg/L of Palmitic Acid; 100 mg/L of Pluronic F-68; 0.010 mg/L of Stearic Acid; 2.20 mg/L of Tween 80; 4551 mg/L of D-Glucose; 130.85 mg/ml of L- Alanine; 147.50 mg/ml of L-Arginine-HCL; 7.50 mg/ml of L-Asparagine-H<sub>2</sub>O; 6.65 mg/ml of L-Aspartic Acid;
- 5 29.56 mg/ml of L-Cystine-2HCL-H<sub>2</sub>O; 31.29 mg/ml of L-Cystine-2HCL; 7.35 mg/ml of L-Glutamic Acid; 365.0 mg/ml of L-Glutamine; 18.75 mg/ml of Glycine; 52.48 mg/ml of L-Histidine-HCL-H<sub>2</sub>O; 106.97 mg/ml of L-Isoleucine; 111.45 mg/ml of L-Leucine; 163.75 mg/ml of L-Lysine HCL; 32.34 mg/ml of L-Methionine; 68.48 mg/ml of L-Phenylalainine; 40.0 mg/ml of L-Proline; 26.25 mg/ml of L-Serine;
- 10 101.05 mg/ml of L-Threonine; 19.22 mg/ml of L-Tryptophan; 91.79 mg/ml of L-Tyrosine-2Na-2H<sub>2</sub>O; and 99.65 mg/ml of L-Valine; 0.0035 mg/L of Biotin; 3.24 mg/L of D-Ca Pantothenate; 11.78 mg/L of Choline Chloride; 4.65 mg/L of Folic Acid; 15.60 mg/L of i-Inositol; 3.02 mg/L of Niacinamide; 3.00 mg/L of Pyridoxal HCL; 0.031 mg/L of Pyridoxine HCL; 0.319 mg/L of Riboflavin; 3.17 mg/L of
- 15 Thiamine HCL; 0.365 mg/L of Thymidine; 0.680 mg/L of Vitamin B<sub>12</sub>; 25 mM of HEPES Buffer; 2.39 mg/L of Na Hypoxanthine; 0.105 mg/L of Lipoic Acid; 0.081 mg/L of Sodium Putrescine-2HCL; 55.0 mg/L of Sodium Pyruvate; 0.0067 mg/L of Sodium Selenite; 20uM of Ethanolamine; 0.122 mg/L of Ferric Citrate; 41.70 mg/L of Methyl-B-Cyclodextrin complexed with Linoleic Acid; 33.33 mg/L of Methyl-B-
- 20 Cyclodextrin complexed with Oleic Acid; 10 mg/L of Methyl-B-Cyclodextrin complexed with Retinal Acetate. Adjust osmolarity to 327 mOsm) with 2mm glutamine and 1x penstrep. (BSA (81-068-3 Bayer) 100gm dissolved in 1L DMEM for a 10% BSA stock solution). Filter the media and collect 50 ul for endotoxin assay in 15ml polystyrene conical.
- 25 The transfection reaction is terminated, preferably by tag-teaming, at the end of the incubation period. Person A aspirates off the transfection media, while person B adds 1.5ml appropriate media to each well. Incubate at 37 degree C for 45 or 72 hours depending on the media used: 1%BSA for 45 hours or CHO-5 for 72 hours.
- 30 On day four, using a 300ul multichannel pipetter, aliquot 600ul in one 1ml deep well plate and the remaining supernatant into a 2ml deep well. The supernatants from each well can then be used in the assays described in Examples 14-21.

It is specifically understood that when activity is obtained in any of the assays described below using a supernatant, the activity originates from either the CRCGCL polypeptide directly (e.g., as a secreted protein) or by CRCGCL inducing expression of other proteins, which are then secreted into the supernatant. Thus, the invention  
5 further provides a method of identifying the protein in the supernatant characterized by an activity in a particular assay.

### ***Example 13: Construction of GAS Reporter Construct***

10 One signal transduction pathway involved in the differentiation and proliferation of cells is called the Jaks-STATs pathway. Activated proteins in the Jaks-STATs pathway bind to gamma activation site "GAS" elements or interferon-sensitive responsive element ("ISRE"), located in the promoter of many genes. The binding of a protein to these elements alter the expression of the associated gene.

15 GAS and ISRE elements are recognized by a class of transcription factors called Signal Transducers and Activators of Transcription, or "STATs." There are six members of the STATs family. Stat1 and Stat3 are present in many cell types, as is Stat2 (as response to IFN-alpha is widespread). Stat4 is more restricted and is not in many cell types though it has been found in T helper class I, cells after treatment with  
20 IL-12. Stat5 was originally called mammary growth factor, but has been found at higher concentrations in other cells including myeloid cells. It can be activated in tissue culture cells by many cytokines.

The STATs are activated to translocate from the cytoplasm to the nucleus upon tyrosine phosphorylation by a set of kinases known as the Janus Kinase ("Jaks")  
25 family. Jaks represent a distinct family of soluble tyrosine kinases and include Tyk2, Jak1, Jak2, and Jak3. These kinases display significant sequence similarity and are generally catalytically inactive in resting cells.

The Jaks are activated by a wide range of receptors summarized in Table 2 below. (Adapted from review by Schidler and Darnell, Ann. Rev. Biochem. 64:621-  
30 51 (1995).) A cytokine receptor family, capable of activating Jaks, is divided into two groups: (a) Class 1 includes receptors for IL-2, IL-3, IL-4, IL-6, IL-7, IL-9, IL-11, IL-12, IL-15, Epo, PRL, GH, G-CSF, GM-CSF, LIF, CNTF, and thrombopoietin; and (b) Class 2 includes IFN-a, IFN-g, and IL-10. The Class 1 receptors share a conserved

cysteine motif (a set of four conserved cysteines and one tryptophan) and a WSXWS motif (a membrane proximal region encoding Trp-Ser-Xxx-Trp-Ser (SEQ ID NO:5)).

Thus, on binding of a ligand to a receptor, Jaks are activated, which in turn activate STATs, which then translocate and bind to GAS elements. This entire process is encompassed in the Jaks-STATs signal transduction pathway.

Therefore, activation of the Jaks-STATs pathway, reflected by the binding of the GAS or the ISRE element, can be used to indicate proteins involved in the proliferation and differentiation of cells. For example, growth factors and cytokines are known to activate the Jaks-STATs pathway. (See Table 2 below.) Thus, by using GAS elements linked to reporter molecules, activators of the Jaks-STATs pathway can be identified.

There is preliminary data that CRCGCL interacts with Jak1.

Table 2

|    | <u>ISRE</u><br><u>Ligand</u>          | <u>JAKs</u> |             |             |             | <u>STATS</u> | <u>GAS(elements) or</u> |
|----|---------------------------------------|-------------|-------------|-------------|-------------|--------------|-------------------------|
|    |                                       | <u>tyk2</u> | <u>Jak1</u> | <u>Jak2</u> | <u>Jak3</u> |              |                         |
|    | <u>IFN family</u>                     |             |             |             |             |              |                         |
| 20 | IFN- $\alpha$ /B                      | +           | +           | -           | -           | 1,2,3        | ISRE                    |
|    | IFN-g<br>(IRF1>Lys6>IFP)              |             | +           | +           | -           | 1            | GAS                     |
|    | IL-10                                 | +           | ?           | ?           | -           | 1,3          |                         |
| 25 | <u>gp130 family</u>                   |             |             |             |             |              |                         |
|    | IL-6 (Pleiotrohic)<br>(IRF1>Lys6>IFP) | +           | +           | +           | ?           | 1,3          | GAS                     |
|    | IL-11(Pleiotrohic)                    | ?           | +           | ?           | ?           | 1,3          |                         |
|    | OnM(Pleiotrohic)                      | ?           | +           | +           | ?           | 1,3          |                         |
| 30 | LIF(Pleiotrohic)                      | ?           | +           | +           | ?           | 1,3          |                         |
|    | CNTF(Pleiotrohic)                     | -/+         | +           | +           | ?           | 1,3          |                         |
|    | G-CSF(Pleiotrohic)                    | ?           | +           | ?           | ?           | 1,3          |                         |
|    | IL-12(Pleiotrohic)                    | +           | -           | +           | +           | 1,3          |                         |
| 35 | <u>g-C family</u>                     |             |             |             |             |              |                         |
|    | IL-2 (lymphocytes)                    | -           | +           | -           | +           | 1,3,5        | GAS                     |
|    | IL-4 (lymph/myeloid)-<br>>>Ly6(IgH)   | -           | +           | -           | +           | 6            | GAS (IRF1 = IFP)        |
|    | IL-7 (lymphocytes)                    | -           | +           | -           | +           | 5            | GAS                     |
| 40 | IL-9 (lymphocytes)                    | -           | +           | -           | +           | 5            | GAS                     |
|    | IL-13 (lymphocyte)                    | -           | +           | ?           | ?           | 6            | GAS                     |
|    | IL-15                                 | ?           | +           | ?           | +           | 5            | GAS                     |

|                                  |                    |   |     |   |   |                    |
|----------------------------------|--------------------|---|-----|---|---|--------------------|
| <u>gp140 family</u>              |                    |   |     |   |   |                    |
|                                  | IL-3 (myeloid)     | - | -   | + | - | 5 GAS              |
|                                  | (IRF1>IFP>>Ly6)    |   |     |   |   |                    |
| 5                                | IL-5 (myeloid)     | - | -   | + | - | 5 GAS              |
|                                  | GM-CSF (myeloid)   | - | -   | + | - | 5 GAS              |
| <u>Growth hormone family</u>     |                    |   |     |   |   |                    |
|                                  | GH                 | ? | -   | + | - | 5                  |
| 10                               | PRL                | ? | +/- | + | - | 1,3,5              |
|                                  | EPO                | ? | -   | + | - | 5 GAS(B-           |
|                                  | CAS>IRF1=IFP>>Ly6) |   |     |   |   |                    |
| <u>Receptor Tyrosine Kinases</u> |                    |   |     |   |   |                    |
| 15                               | EGF                | ? | +   | + | - | 1,3 GAS (IRF1)     |
|                                  | PDGF               | ? | +   | + | - | 1,3                |
|                                  | CSF-1              | ? | +   | + | - | 1,3 GAS (not IRF1) |

20

To construct a synthetic GAS containing promoter element, which is used in the Biological Assays described in Examples 14-15, a PCR based strategy is employed to generate a GAS-SV40 promoter sequence. The 5' primer contains four tandem copies of the GAS binding site found in the IRF1 promoter and previously

25 demonstrated to bind STATs upon induction with a range of cytokines (Rothman et al., Immunity 1:457-468 (1994).), although other GAS or ISRE elements can be used instead. The 5' primer also contains 18bp of sequence complementary to the SV40 early promoter sequence and is flanked with an XhoI site. The sequence of the 5' primer is:

30 5':GCGCCTCGAGATTTCCTCGAAATCTAGATTTCCTCGAAATGATTTCCTCGAAATGATTTCCTCGAAATATCTGCCATCTCAATTAG:3' (SEQ ID NO:6)

The downstream primer is complementary to the SV40 promoter and is flanked with a Hind III site: 5':GCGGCAAGCTTTTGTCAAAGCCTAGGC:3' (SEQ ID NO:7)

35 PCR amplification is performed using the SV40 promoter template present in the B-gal:promoter plasmid obtained from Clontech. The resulting PCR fragment is digested with XhoI/Hind III and subcloned into BLSK2-. (Stratagene.) Sequencing with forward and reverse primers confirms that the insert contains the following sequence:



5':CTCGAGATTTC~~CCCCGAAATCTAGATTTC~~CCCCGAAATGATTTC~~CCCCGAAA~~  
 TGATTTC~~CCCCGAAATATCTGCCATCTCAATTAGTCAGCAACCATAGTCCCG~~  
 CCCCTAACTCCGCCCATCCCGCCCCTAACTCCGCCCAGTTCCGCCCATTCT  
 CCGCCCCATGGCTGACTAATTTTTTTTATTTATGCAGAGGCCGAGGCCGCC  
 5 TCGGCCTCTGAGCTATTCCAGAAGTAGTGAGGAGGCTTTTTTTGGAGGCCT  
 AGGCTTTTGCAAAAAAGCTT:3' (SEQ ID NO:8)

With this GAS promoter element linked to the SV40 promoter, a GAS:SEAP2 reporter construct is next engineered. Here, the reporter molecule is a secreted alkaline phosphatase, or "SEAP." Clearly, however, any reporter molecule can be  
 10 instead of SEAP, in this or in any of the other Examples. Well known reporter molecules that can be used instead of SEAP include chloramphenicol acetyltransferase (CAT), luciferase, alkaline phosphatase, B-galactosidase, green fluorescent protein (GFP), or any protein detectable by an antibody.

The above sequence confirmed synthetic GAS-SV40 promoter element is  
 15 subcloned into the pSEAP-Promoter vector obtained from Clontech using HindIII and XhoI, effectively replacing the SV40 promoter with the amplified GAS:SV40 promoter element, to create the GAS-SEAP vector. However, this vector does not contain a neomycin resistance gene, and therefore, is not preferred for mammalian expression systems.

20 Thus, in order to generate mammalian stable cell lines expressing the GAS-SEAP reporter, the GAS-SEAP cassette is removed from the GAS-SEAP vector using SalI and NotI, and inserted into a backbone vector containing the neomycin resistance gene, such as pGFP-1 (Clontech), using these restriction sites in the multiple cloning site, to create the GAS-SEAP/Neo vector. Once this vector is transfected into  
 25 mammalian cells, this vector can then be used as a reporter molecule for GAS binding as described in Examples 14-15.

Other constructs can be made using the above description and replacing GAS with a different promoter sequence. For example, construction of reporter molecules containing NFK-B and EGR promoter sequences are described in Examples 16 and  
 30 17. However, many other promoters can be substituted using the protocols described in these Examples. For instance, SRE, IL-2, NFAT, or Osteocalcin promoters can be substituted, alone or in combination (e.g., GAS/NF-KB/EGR, GAS/NF-KB, IL-2/NFAT, or NF-KB/GAS). Similarly, other cell lines can be used to test reporter

09376430-08189

construct activity, such as HELA (epithelial), HUVEC (endothelial), Reh (B-cell), Saos-2 (osteoblast), HUVAC (aortic), or Cardiomyocyte.

***Example 14: High-Throughput Screening Assay  
for T-cell Activity.***

5

The following protocol is used to assess T-cell activity of CRCGCL by determining whether CRCGCL supernatant proliferates and/or differentiates T-cells. T-cell activity is assessed using the GAS/SEAP/Neo construct produced in Example  
10 13. Thus, factors that increase SEAP activity indicate the ability to activate the Jaks-STATS signal transduction pathway. The T-cell used in this assay is Jurkat T-cells (ATCC Accession No. TIB-152), although Molt-3 cells (ATCC Accession No. CRL-1552) and Molt-4 cells (ATCC Accession No. CRL-1582) cells can also be used.

Jurkat T-cells are lymphoblastic CD4+ Th1 helper cells. In order to generate  
15 stable cell lines, approximately 2 million Jurkat cells are transfected with the GAS-SEAP/neo vector using DMRIE-C (Life Technologies)(transfection procedure described below). The transfected cells are seeded to a density of approximately 20,000 cells per well and transfectants resistant to 1 mg/ml gentamicin selected.

Resistant colonies are expanded and then tested for their response to increasing  
20 concentrations of interferon gamma. The dose response of a selected clone is demonstrated.

Specifically, the following protocol will yield sufficient cells for 75 wells containing 200 ul of cells. Thus, it is either scaled up, or performed in multiple to generate sufficient cells for multiple 96 well plates. Jurkat cells are maintained in  
25 RPMI + 10% serum with 1%Pen-Strep. Combine 2.5 mls of OPTI-MEM (Life Technologies) with 10 ug of plasmid DNA in a T25 flask. Add 2.5 ml OPTI-MEM containing 50 ul of DMRIE-C and incubate at room temperature for 15-45 mins.

During the incubation period, count cell concentration, spin down the required number of cells ( $10^7$  per transfection), and resuspend in OPTI-MEM to a final  
30 concentration of  $10^7$  cells/ml. Then add 1ml of  $1 \times 10^7$  cells in OPTI-MEM to T25 flask and incubate at 37 degree C for 6 hrs. After the incubation, add 10 ml of RPMI + 15% serum.

668730 0619260

The Jurkat:GAS-SEAP stable reporter lines are maintained in RPMI + 10% serum, 1 mg/ml Genticin, and 1% Pen-Strep. These cells are treated with supernatants containing CRCGCL polypeptides or CRCGCL induced polypeptides as produced by the protocol described in Example 12.

5 On the day of treatment with the supernatant, the cells should be washed and resuspended in fresh RPMI + 10% serum to a density of 500,000 cells per ml. The exact number of cells required will depend on the number of supernatants being screened. For one 96 well plate, approximately 10 million cells (for 10 plates, 100 million cells) are required.

10 Transfer the cells to a triangular reservoir boat, in order to dispense the cells into a 96 well dish, using a 12 channel pipette. Using a 12 channel pipette, transfer 200 ul of cells into each well (therefore adding 100, 000 cells per well).

After all the plates have been seeded, 50 ul of the supernatants are transferred directly from the 96 well plate containing the supernatants into each well using a 12  
15 channel pipette. In addition, a dose of exogenous interferon gamma (0.1, 1.0, 10 ng) is added to wells H9, H10, and H11 to serve as additional positive controls for the assay.

The 96 well dishes containing Jurkat cells treated with supernatants are placed in an incubator for 48 hrs (note: this time is variable between 48-72 hrs). 35 ul  
20 samples from each well are then transferred to an opaque 96 well plate using a 12 channel pipette. The opaque plates should be covered (using sellophene covers) and stored at -20 degree C until SEAP assays are performed according to Example 18. The plates containing the remaining treated cells are placed at 4 degree C and serve as a source of material for repeating the assay on a specific well if desired.

25 As a positive control, 100 Unit/ml interferon gamma can be used which is known to activate Jurkat T cells. Over 30 fold induction is typically observed in the positive control wells.

### ***Example 15: High-Throughput Screening Assay Identifying Myeloid Activity***

30

The following protocol is used to assess myeloid activity of CRCGCL by determining whether CRCGCL proliferates and/or differentiates myeloid cells.

Myeloid cell activity is assessed using the GAS/SEAP/Neo construct produced in Example 13. Thus, factors that increase SEAP activity indicate the ability to activate the Jaks-STATS signal transduction pathway. The myeloid cell used in this assay is U937, a pre-monocyte cell line, although TF-1, HL60, or KG1 can be used.

- 5           To transiently transfect U937 cells with the GAS/SEAP/Neo construct produced in Example 13, a DEAE-Dextran method (Kharbanda et. al., 1994, Cell Growth & Differentiation, 5:259-265) is used. First, harvest  $2 \times 10^7$  U937 cells and wash with PBS. The U937 cells are usually grown in RPMI 1640 medium containing 10% heat-inactivated fetal bovine serum (FBS) supplemented with 100 units/ml penicillin and 100 mg/ml streptomycin.

Next, suspend the cells in 1 ml of 20 mM Tris-HCl (pH 7.4) buffer containing 0.5 mg/ml DEAE-Dextran, 8 ug GAS-SEAP2 plasmid DNA, 140 mM NaCl, 5 mM KCl, 375 uM  $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$ , 1 mM  $\text{MgCl}_2$ , and 675 uM  $\text{CaCl}_2$ . Incubate at 37 degree C for 45 min.

- 15           Wash the cells with RPMI 1640 medium containing 10% FBS and then resuspend in 10 ml complete medium and incubate at 37 degree C for 36 hr.

The GAS-SEAP/U937 stable cells are obtained by growing the cells in 400 ug/ml G418. ~~The G418-free medium is used for routine growth but every one to two~~ months, the cells should be re-grown in 400 ug/ml G418 for couple of passages.

- 20           These cells are tested by harvesting  $1 \times 10^8$  cells (this is enough for ten 96-well plates assay) and wash with PBS. Suspend the cells in 200 ml above described growth medium, with a final density of  $5 \times 10^5$  cells/ml. Plate 200 ul cells per well in the 96-well plate (or  $1 \times 10^5$  cells/well).

- 25           Add 50 ul of the supernatant prepared by the protocol described in Example 12. Incubate at 37 degree C for 48 to 72 hr. As a positive control, 100 Unit/ml interferon gamma can be used which is known to activate U937 cells. Over 30 fold induction is typically observed in the positive control wells. SEAP assay the supernatant according to the protocol described in Example 18.

***Example 16: High-Throughput Screening Assay  
Identifying Neuronal Activity.***

When cells undergo differentiation and proliferation, a group of genes are  
5 activated through many different signal transduction pathways. One of these genes,  
EGR1 (early growth response gene 1), is induced in various tissues and cell types  
upon activation. The promoter of EGR1 is responsible for such induction. Using the  
EGR1 promoter linked to reporter molecules, activation of cells can be assessed by  
CRCGCL.

10 Particularly, the following protocol is used to assess neuronal activity in PC12  
cell lines. PC12 cells (rat pheochromocytoma cells) are known to proliferate and/or  
differentiate by activation with a number of mitogens, such as TPA (tetradecanoyl  
phorbol acetate), NGF (nerve growth factor), and EGF (epidermal growth factor).  
The EGR1 gene expression is activated during this treatment. Thus, by stably  
15 transfecting PC12 cells with a construct containing an EGR promoter linked to SEAP  
reporter, activation of PC12 cells by CRCGCL can be assessed.

The EGR/SEAP reporter construct can be assembled by the following  
protocol. The EGR-1 promoter sequence (-633 to +1)(Sakamoto K et al., Oncogene  
6:867-871 (1991)) can be PCR amplified from human genomic DNA using the  
20 following primers:

5' GCGCTCGAGGGATGACAGCGATAGAACCCCGG -3' (SEQ ID NO:9)

5' GCGAAGCTTCGCGACTCCCCGGATCCGCCTC-3' (SEQ ID NO:10)

Using the GAS:SEAP/Neo vector produced in Example 13, EGR1 amplified  
product can then be inserted into this vector. Linearize the GAS:SEAP/Neo vector  
25 using restriction enzymes XhoI/HindIII, removing the GAS/SV40 stuffer. Restrict the  
EGR1 amplified product with these same enzymes. Ligate the vector and the EGR1  
promoter.

To prepare 96 well-plates for cell culture, two mls of a coating solution (1:30  
dilution of collagen type I (Upstate Biotech Inc. Cat#08-115) in 30% ethanol (filter  
30 sterilized)) is added per one 10 cm plate or 50 ml per well of the 96-well plate, and  
allowed to air dry for 2 hr.

PC12 cells are routinely grown in RPMI-1640 medium (Bio Whittaker)  
containing 10% horse serum (JRH BIOSCIENCES, Cat. # 12449-78P), 5% heat-

5 Transfect the EGR/SEAP/Neo construct into PC12 using the Lipofectamine protocol described in Example 12. EGR-SEAP/PC12 stable cells are obtained by growing the cells in 300 ug/ml G418. The G418-free medium is used for routine growth but every one to two months, the cells should be re-grown in 300 ug/ml G418 for couple of passages.

The next morning, remove the medium and wash the cells with PBS. Scrape  
15 off the cells from the plate, suspend the cells well in 2 ml low serum medium. Count  
the cell number and add more low serum medium to reach final cell density as  $5 \times 10^5$   
cells/ml.

25 **Example 17: High-Throughput Screening Assay for T-cell Activity**

NF-KB (Nuclear Factor KB) is a transcription factor activated by a wide variety of agents including the inflammatory cytokines IL-1 and TNF, CD30 and CD40, lymphotoxin-alpha and lymphotoxin-beta, by exposure to LPS or thrombin, and by expression of certain viral gene products. As a transcription factor, NF-KB regulates the expression of genes involved in immune cell activation, control of apoptosis (NF- KB appears to shield cells from apoptosis), B and T-cell development, anti-viral and antimicrobial responses, and multiple stress responses.

In non-stimulated conditions, NF- KB is retained in the cytoplasm with I-KB (Inhibitor KB). However, upon stimulation, I- KB is phosphorylated and degraded, causing NF- KB to shuttle to the nucleus, thereby activating transcription of target genes. Target genes activated by NF- KB include IL-2, IL-6, GM-CSF, ICAM-1 and class 1 MHC.

Due to its central role and ability to respond to a range of stimuli, reporter constructs utilizing the NF-KB promoter element are used to screen the supernatants produced in Example 12. Activators or inhibitors of NF-KB would be useful in treating diseases. For example, inhibitors of NF-KB could be used to treat those diseases related to the acute or chronic activation of NF-KB, such as rheumatoid arthritis.

To construct a vector containing the NF-KB promoter element, a PCR based strategy is employed. The upstream primer contains four tandem copies of the NF-KB binding site (GGGGACTTTCCC) (SEQ ID NO:11), 18 bp of sequence complementary to the 5' end of the SV40 early promoter sequence, and is flanked with an XhoI site:

5':GCGGCCTCGAGGGGACTTTCCCGGGGACTTTCCGGGGACTTTCCGGGAC  
TTTCCATCCTGCCATCTCAATTAG:3' (SEQ ID NO:12)

The downstream primer is complementary to the 3' end of the SV40 promoter and is flanked with a Hind III site:

5':GCGGCAAGCTTTTTGCAAAGCCTAGGC:3' (SEQ ID NO:7)

PCR amplification is performed using the SV40 promoter template present in the pB-gal:promoter plasmid obtained from Clontech. The resulting PCR fragment is digested with XhoI and Hind III and subcloned into BLSK2-. (Stratagene). Sequencing with the T7 and T3 primers confirms the insert contains the following sequence:

5':CTCGAGGGGACTTTCCCGGGGACTTTCCGGGGACTTTCCGGGACTTTCC  
ATCTGCCATCTCAATTAGTCAGCAACCATAGTCCCGCCCCTAACTCCGCCC  
ATCCCGCCCCTAACTCCGCCCAGTTCCGCCCATTCTCCGCCCCATGGCTGA  
CTAATTTTTTTTATTTATGCAGAGGCCGAGGCCGCCTCGGCCTCTGAGCTA  
TTCCAGAAGTAGTGAGGAGGCTTTTTTGGAGGCCTAGGCTTTTGCAAAAA  
GCTT:3' (SEQ ID NO:13)

Next, replace the SV40 minimal promoter element present in the pSEAP2-promoter plasmid (Clontech) with this NF-KB/SV40 fragment using XhoI and HindIII. However, this vector does not contain a neomycin resistance gene, and therefore, is not preferred for mammalian expression systems.

5 In order to generate stable mammalian cell lines, the NF-KB/SV40/SEAP cassette is removed from the above NF-KB/SEAP vector using restriction enzymes SalI and NotI, and inserted into a vector containing neomycin resistance. Particularly, the NF-KB/SV40/SEAP cassette was inserted into pGFP-1 (Clontech), replacing the GFP gene, after restricting pGFP-1 with SalI and NotI.

10 Once NF-KB/SV40/SEAP/Neo vector is created, stable Jurkat T-cells are created and maintained according to the protocol described in Example 14. Similarly, the method for assaying supernatants with these stable Jurkat T-cells is also described in Example 14. As a positive control, exogenous TNF alpha (0.1, 1, 10 ng) is added to wells H9, H10, and H11, with a 5-10 fold activation typically observed.

15

### ***Example 18: Assay for SEAP Activity***

As a reporter molecule for the assays described in Examples 14-17, SEAP activity is assayed using the Tropix Phospho-light Kit (Cat. BP-400) according to the  
20 following general procedure. The Tropix Phospho-light Kit supplies the Dilution, Assay, and Reaction Buffers used below.

Prime a dispenser with the 2.5x Dilution Buffer and dispense 15 ul of 2.5x dilution buffer into Optiplates containing 35 ul of a supernatant. Seal the plates with a plastic sealer and incubate at 65 degree C for 30 min. Separate the Optiplates to  
25 avoid uneven heating.

Cool the samples to room temperature for 15 minutes. Empty the dispenser and prime with the Assay Buffer. Add 50 ml Assay Buffer and incubate at room temperature 5 min. Empty the dispenser and prime with the Reaction Buffer (see Table 3 below). Add 50 ul Reaction Buffer and incubate at room temperature for 20  
30 minutes. Since the intensity of the chemiluminescent signal is time dependent, and it takes about 10 minutes to read 5 plates on luminometer, one should treat 5 plates at each time and start the second set 10 minutes later.



Read the relative light unit in the luminometer. Set H12 as blank, and print the results. An increase in chemiluminescence indicates reporter activity.

**Table 3**

5

**Reaction Buffer Formulation:**

| # of plates | Rxn buffer diluent (ml) | CSPD (ml) |
|-------------|-------------------------|-----------|
| 10          | 60                      | 3         |
| 11          | 65                      | 3.25      |
| 12          | 70                      | 3.5       |
| 13          | 75                      | 3.75      |
| 14          | 80                      | 4         |
| 15          | 85                      | 4.25      |
| 16          | 90                      | 4.5       |
| 17          | 95                      | 4.75      |
| 18          | 100                     | 5         |
| 19          | 105                     | 5.25      |
| 20          | 110                     | 5.5       |
| 21          | 115                     | 5.75      |
| 22          | 120                     | 6         |
| 23          | 125                     | 6.25      |
| 24          | 130                     | 6.5       |
| 25          | 135                     | 6.75      |
| 26          | 140                     | 7         |
| 27          | 145                     | 7.25      |
| 28          | 150                     | 7.5       |
| 29          | 155                     | 7.75      |
| 30          | 160                     | 8         |
| 31          | 165                     | 8.25      |
| 32          | 170                     | 8.5       |
| 33          | 175                     | 8.75      |
| 34          | 180                     | 9         |
| 35          | 185                     | 9.25      |
| 36          | 190                     | 9.5       |
| 37          | 195                     | 9.75      |
| 38          | 200                     | 10        |
| 39          | 205                     | 10.25     |
| 40          | 210                     | 10.5      |

003430 0E49E60

|    |     |       |
|----|-----|-------|
| 41 | 215 | 10.75 |
| 42 | 220 | 11    |
| 43 | 225 | 11.25 |
| 44 | 230 | 11.5  |
| 45 | 235 | 11.75 |
| 46 | 240 | 12    |
| 47 | 245 | 12.25 |
| 48 | 250 | 12.5  |
| 49 | 255 | 12.75 |
| 50 | 260 | 13    |

### ***Example 19: High-Throughput Screening Assay Identifying Changes in Small Molecule Concentration and Membrane Permeability***

5 Binding of a ligand to a receptor is known to alter intracellular levels of small molecules, such as calcium, potassium, sodium, and pH, as well as alter membrane potential. These alterations can be measured in an assay to identify supernatants which bind to receptors of a particular cell. Although the following protocol describes an assay for calcium, this protocol can easily be modified to detect changes  
10 ~~in potassium, sodium, pH, membrane potential, or any other small molecule which is~~  
detectable by a fluorescent probe.

The following assay uses Fluorometric Imaging Plate Reader (“FLIPR”) to measure changes in fluorescent molecules (Molecular Probes) that bind small molecules. Clearly, any fluorescent molecule detecting a small molecule can be used instead of the calcium fluorescent molecule, fluo-3, used here.

For adherent cells, seed the cells at 10,000 -20,000 cells/well in a Co-star black 96-well plate with clear bottom. The plate is incubated in a CO<sub>2</sub> incubator for 20 hours. The adherent cells are washed two times in Biotek washer with 200 ul of HBSS (Hank's Balanced Salt Solution) leaving 100 ul of buffer after the final wash.

20            A stock solution of 1 mg/ml fluo-3 is made in 10% pluronic acid DMSO. To load the cells with fluo-3, 50 ul of 12 ug/ml fluo-3 is added to each well. The plate is incubated at 37 degree C in a CO<sub>2</sub> incubator for 60 min. The plate is washed four times in the Biotek washer with HBSS leaving 100 ul of buffer.

For non-adherent cells, the cells are spun down from culture media. Cells are  
25 re-suspended to  $2-5 \times 10^6$  cells/ml with HBSS in a 50-ml conical tube. 4 ul of 1 mg/ml

5

10

15

### Identifying Tyrosine Kinase Activity

20

30

Because of the wide range of known factors capable of stimulating tyrosine kinase activity, identifying whether CRCGCL or a molecule induced by CRCGCL is capable of activating tyrosine kinase signal transduction pathways is of interest. Therefore, the following protocol is designed to identify such molecules capable of activating the tyrosine kinase signal transduction pathways.

Seed target cells (e.g., primary keratinocytes) at a density of approximately 25,000 cells per well in a 96 well Loprodyne Silent Screen Plates purchased from Nalge Nunc (Naperville, IL). The plates are sterilized with two 30 minute rinses with 100% ethanol, rinsed with water and dried overnight. Some plates are coated for 2 hr with 100 ml of cell culture grade type I collagen (50 mg/ml), gelatin (2%) or polylysine (50 mg/ml), all of which can be purchased from Sigma Chemicals (St. Louis, MO) or 10% Matrigel purchased from Becton Dickinson (Bedford, MA), or calf serum, rinsed with PBS and stored at 4 degree C. Cell growth on these plates is assayed by seeding 5,000 cells/well in growth medium and indirect quantitation of cell number through use of alamarBlue as described by the manufacturer Alamar Biosciences, Inc. (Sacramento, CA) after 48 hr. Falcon plate covers #3071 from Becton Dickinson (Bedford, MA) are used to cover the Loprodyne Silent Screen Plates. ~~Falcon Microtest III cell culture plates can also be used in some proliferation experiments.~~

To prepare extracts, A431 cells are seeded onto the nylon membranes of Loprodyne plates (20,000/200ml/well) and cultured overnight in complete medium. Cells are quiesced by incubation in serum-free basal medium for 24 hr. After 5-20 minutes treatment with EGF (60ng/ml) or 50 ul of the supernatant produced in Example 12, the medium was removed and 100 ml of extraction buffer ((20 mM HEPES pH 7.5, 0.15 M NaCl, 1% Triton X-100, 0.1% SDS, 2 mM Na<sub>3</sub>VO<sub>4</sub>, 2 mM Na<sub>4</sub>P<sub>2</sub>O<sub>7</sub> and a cocktail of protease inhibitors (# 1836170) obtained from Boehringer Mannheim (Indianapolis, IN) is added to each well and the plate is shaken on a rotating shaker for 5 minutes at 4°C. The plate is then placed in a vacuum transfer manifold and the extract filtered through the 0.45 mm membrane bottoms of each well using house vacuum. Extracts are collected in a 96-well catch/assay plate in the bottom of the vacuum manifold and immediately placed on ice. To obtain extracts clarified by centrifugation, the content of each well, after

detergent solubilization for 5 minutes, is removed and centrifuged for 15 minutes at 4 degree C at 16,000 x g.

Test the filtered extracts for levels of tyrosine kinase activity. Although many methods of detecting tyrosine kinase activity are known, one method is described here.

Generally, the tyrosine kinase activity of a supernatant is evaluated by determining its ability to phosphorylate a tyrosine residue on a specific substrate (a biotinylated peptide). Biotinylated peptides that can be used for this purpose include PSK1 (corresponding to amino acids 6-20 of the cell division kinase cdc2-p34) and PSK2 (corresponding to amino acids 1-17 of gastrin). Both peptides are substrates for a range of tyrosine kinases and are available from Boehringer Mannheim.

The tyrosine kinase reaction is set up by adding the following components in order. First, add 10ul of 5uM Biotinylated Peptide, then 10ul ATP/Mg<sub>2</sub><sup>+</sup> (5mM ATP/50mM MgCl<sub>2</sub>), then 10ul of 5x Assay Buffer (40mM imidazole hydrochloride, pH7.3, 40 mM beta-glycerophosphate, 1mM EGTA, 100mM MgCl<sub>2</sub>, 5 mM MnCl<sub>2</sub>, 0.5 mg/ml BSA), then 5ul of Sodium Vanadate(1mM), and then 5ul of water. Mix the components gently and preincubate the reaction mix at 30 degree C for 2 min. Initial the reaction by adding 10ul of the control enzyme or the filtered supernatant.

The tyrosine kinase assay reaction is then terminated by adding 10 ul of 120mM EDTA and place the reactions on ice.

Tyrosine kinase activity is determined by transferring 50 ul aliquot of reaction mixture to a microtiter plate (MTP) module and incubating at 37 degree C for 20 min. This allows the streptavidin coated 96 well plate to associate with the biotinylated peptide. Wash the MTP module with 300ul/well of PBS four times. Next add 75 ul of anti-phosphotyrosine antibody conjugated to horse radish peroxidase(anti-P-Tyr-POD(0.5u/ml)) to each well and incubate at 37 degree C for one hour. Wash the well as above.

Next add 100ul of peroxidase substrate solution (Boehringer Mannheim) and incubate at room temperature for at least 5 mins (up to 30 min). Measure the absorbance of the sample at 405 nm by using ELISA reader. The level of bound peroxidase activity is quantitated using an ELISA reader and reflects the level of tyrosine kinase activity.

***Example 21: High-Throughput Screening Assay  
Identifying Phosphorylation Activity***

5 As a potential alternative and/or compliment to the assay of protein tyrosine kinase activity described in Example 20, an assay which detects activation (phosphorylation) of major intracellular signal transduction intermediates can also be used. For example, as described below one particular assay can detect tyrosine phosphorylation of the Erk-1 and Erk-2 kinases. However, phosphorylation of other  
10 molecules, such as Raf, JNK, p38 MAP, Map kinase kinase (MEK), MEK kinase, Src, Muscle specific kinase (MuSK), IRAK, Tec, and Janus, as well as any other phosphoserine, phosphotyrosine, or phosphothreonine molecule, can be detected by substituting these molecules for Erk-1 or Erk-2 in the following assay.

Specifically, assay plates are made by coating the wells of a 96-well ELISA  
15 plate with 0.1ml of protein G (1ug/ml) for 2 hr at room temp, (RT). The plates are then rinsed with PBS and blocked with 3% BSA/PBS for 1 hr at RT. The protein G plates are then treated with 2 commercial monoclonal antibodies (100ng/well) against Erk-1 and Erk-2 (1 hr at RT) (Santa Cruz Biotechnology). (To detect other  
20 molecules, this step can easily be modified by substituting a monoclonal antibody detecting any of the above described molecules.) After 3-5 rinses with PBS, the plates are stored at 4 degree C until use.

A431 cells are seeded at 20,000/well in a 96-well Loprodyne filterplate and cultured overnight in growth medium. The cells are then starved for 48 hr in basal medium (DMEM) and then treated with EGF (6ng/well) or 50 ul of the supernatants  
25 obtained in Example 12 for 5-20 minutes. The cells are then solubilized and extracts filtered directly into the assay plate.

After incubation with the extract for 1 hr at RT, the wells are again rinsed. As a positive control, a commercial preparation of MAP kinase (10ng/well) is used in place of A431 extract. Plates are then treated with a commercial polyclonal (rabbit)  
30 antibody (1ug/ml) which specifically recognizes the phosphorylated epitope of the Erk-1 and Erk-2 kinases (1 hr at RT). This antibody is biotinylated by standard procedures. The bound polyclonal antibody is then quantitated by successive incubations with Europium-streptavidin and Europium fluorescence enhancing

00376430 081899

reagent in the Wallac DELFIA instrument (time-resolved fluorescence). An increased fluorescent signal over background indicates a phosphorylation by CRCGCL or a molecule induced by CRCGCL.

5                    ***Example 22: Method of Determining Alterations  
in the CRCGCL Gene***

10                    RNA isolated from entire families or individual patients presenting with a phenotype of interest (such as a disease) is be isolated. cDNA is then generated from these RNA samples using protocols known in the art. (See, Sambrook.) The cDNA is then used as a template for PCR, employing primers surrounding regions of interest in SEQ ID NO:1. Suggested PCR conditions consist of 35 cycles at 95 degree C for 30 seconds; 60-120 seconds at 52-58 degree C; and 60-120 seconds at 70 degree C, using buffer solutions described in Sidransky, D., et al., Science 252:706 (1991).

15                    PCR products are then sequenced using primers labeled at their 5' end with T4 polynucleotide kinase, employing SequiTherm Polymerase. (Epicentre Technologies). The intron-exon borders of selected exons of CRCGCL is also determined and genomic PCR products analyzed to confirm the results. PCR products harboring suspected mutations in CRCGCL is then cloned and sequenced to  
20                    validate the results of the direct sequencing.

                      PCR products of CRCGCL are cloned into T-tailed vectors as described in Holton, T.A. and Graham, M.W., Nucleic Acids Research, 19:1156 (1991) and sequenced with T7 polymerase (United States Biochemical). Affected individuals are identified by mutations in CRCGCL not present in unaffected individuals.

25                    Genomic rearrangements are also observed as a method of determining alterations in the CRCGCL gene. Genomic clones isolated according to Example 2 are nick-translated with digoxigenindeoxy-uridine 5'-triphosphate (Boehringer Mannheim), and FISH performed as described in Johnson, Cg. et al., Methods Cell Biol. 35:73-99 (1991). Hybridization with the labeled probe is carried out using  
30                    vast excess of human cot-1 DNA for specific hybridization to the CRCGCL genomic locus.

                      Chromosomes are counterstained with 4,6-diamino-2-phenylidole and propidium iodide, producing a combination of C- and R-bands. Aligned images for

precise mapping are obtained using a triple-band filter set (Chroma Technology, Brattleboro, VT) in combination with a cooled charge-coupled device camera (Photometrics, Tucson, AZ) and variable excitation wavelength filters. (Johnson, Cv. et al., Genet. Anal. Tech. Appl., 8:75 (1991).) Image collection, analysis and chromosomal fractional length measurements are performed using the ISee Graphical Program System. (Inovision Corporation, Durham, NC.) Chromosome alterations of the genomic region of CRCGCL (hybridized by the probe) are identified as insertions, deletions, and translocations. These CRCGCL alterations are used as a diagnostic marker for an associated disease.

***Example 23: Method of Detecting Abnormal Levels  
of CRCGCL in a Biological Sample***

CRCGCL polypeptides can be detected in a biological sample, and if an increased or decreased level of CRCGCL is detected, this polypeptide is a marker for a particular phenotype. Methods of detection are numerous, and thus, it is understood that one skilled in the art can modify the following assay to fit their particular needs.

For example, antibody-sandwich ELISAs are used to detect CRCGCL in a sample, preferably a biological sample. Wells of a microtiter plate are coated with specific antibodies to CRCGCL, at a final concentration of 0.2 to 10 ug/ml. The antibodies are either monoclonal or polyclonal and are produced by the method described in Example 11. The wells are blocked so that non-specific binding of CRCGCL to the well is reduced.

The coated wells are then incubated for > 2 hours at RT with a sample containing CRCGCL. Preferably, serial dilutions of the sample should be used to validate results. The plates are then washed three times with deionized or distilled water to remove unbounded CRCGCL.

Next, 50 ul of specific antibody-alkaline phosphatase conjugate, at a concentration of 25-400 ng, is added and incubated for 2 hours at room temperature. The plates are again washed three times with deionized or distilled water to remove unbounded conjugate.

Add 75 ul of 4-methylumbelliferyl phosphate (MUP) or p-nitrophenyl phosphate (NPP) substrate solution to each well and incubate 1 hour at room



temperature. Measure the reaction by a microtiter plate reader. Prepare a standard curve, using serial dilutions of a control sample, and plot CRCGCL polypeptide concentration on the X-axis (log scale) and fluorescence or absorbance of the Y-axis (linear scale). Interpolate the concentration of the CRCGCL in the sample using the standard curve.

#### ***Example 24: Formulating a Polypeptide***

The CRCGCL composition will be formulated and dosed in a fashion consistent with good medical practice, taking into account the clinical condition of the individual patient (especially the side effects of treatment with the CRCGCL polypeptide alone), the site of delivery, the method of administration, the scheduling of administration, and other factors known to practitioners. The "effective amount" for purposes herein is thus determined by such considerations.

As a general proposition, the total pharmaceutically effective amount of CRCGCL administered parenterally per dose will be in the range of about 1ug/kg/day to 10 mg/kg/day of patient body weight, although, as noted above, this will be subject to therapeutic discretion. More preferably, this dose is at least 0.01 mg/kg/day, and most preferably for humans between about 0.01 and 1 mg/kg/day for the hormone. If given continuously, CRCGCL is typically administered at a dose rate of about 1 ug/kg/hour to about 50 ug/kg/hour, either by 1-4 injections per day or by continuous subcutaneous infusions, for example, using a mini-pump. An intravenous bag solution may also be employed. The length of treatment needed to observe changes and the interval following treatment for responses to occur appears to vary depending on the desired effect.

Pharmaceutical compositions containing CRCGCL are administered orally, rectally, parenterally, intracistemally, intravaginally, intraperitoneally, topically (as by powders, ointments, gels, drops or transdermal patch), buccally, or as an oral or nasal spray. "Pharmaceutically acceptable carrier" refers to a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

CRCGCL is also suitably administered by sustained-release systems. Suitable examples of sustained-release compositions include semi-permeable polymer matrices in the form of shaped articles, e.g., films, or microcapsules. Sustained-release matrices include polylactides (U.S. Pat. No. 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma-ethyl-L-glutamate (Sidman, U. et al., Biopolymers 22:547-556 (1983)), poly (2- hydroxyethyl methacrylate) (R. Langer et al., J. Biomed. Mater. Res. 15:167-277 (1981), and R. Langer, Chem. Tech. 12:98-105 (1982)), ethylene vinyl acetate (R. Langer et al.) or poly-D- (-)-3-hydroxybutyric acid (EP 133,988). Sustained-release compositions also include liposomally entrapped CRCGCL polypeptides. Liposomes containing the CRCGCL are prepared by methods known per se: DE 3,218,121; Epstein et al., Proc. Natl. Acad. Sci. USA 82:3688-3692 (1985); Hwang et al., Proc. Natl. Acad. Sci. USA 77:4030-4034 (1980); EP 52,322; EP 36,676; EP 88,046; EP 143,949; EP 142,641; Japanese Pat. Appl. 83-118008; U.S. Pat. Nos. 4,485,045 and 4,544,545; and EP 102,324. Ordinarily, the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. percent cholesterol, the selected proportion being adjusted for the optimal secreted polypeptide therapy.

For parenteral administration, in one embodiment, CRCGCL is formulated generally by mixing it at the desired degree of purity, in a unit dosage injectable form (solution, suspension, or emulsion), with a pharmaceutically acceptable carrier, i.e., one that is non-toxic to recipients at the dosages and concentrations employed and is compatible with other ingredients of the formulation. For example, the formulation preferably does not include oxidizing agents and other compounds that are known to be deleterious to polypeptides.

Generally, the formulations are prepared by contacting CRCGCL uniformly and intimately with liquid carriers or finely divided solid carriers or both. Then, if necessary, the product is shaped into the desired formulation. Preferably the carrier is a parenteral carrier, more preferably a solution that is isotonic with the blood of the recipient. Examples of such carrier vehicles include water, saline, Ringer's solution, and dextrose solution. Non-aqueous vehicles such as fixed oils and ethyl oleate are also useful herein, as well as liposomes.

The carrier suitably contains minor amounts of additives such as substances that enhance isotonicity and chemical stability. Such materials are non-toxic to

recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, succinate, acetic acid, and other organic acids or their salts; antioxidants such as ascorbic acid; low molecular weight (less than about ten residues) polypeptides, e.g., polyarginine or tripeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids, such as glycine, glutamic acid, aspartic acid, or arginine; monosaccharides, disaccharides, and other carbohydrates including cellulose or its derivatives, glucose, manose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; counterions such as sodium; and/or nonionic surfactants such as polysorbates, poloxamers, or PEG.

CRCGCL is typically formulated in such vehicles at a concentration of about 0.1 mg/ml to 100 mg/ml, preferably 1-10 mg/ml, at a pH of about 3 to 8. It will be understood that the use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of polypeptide salts.

CRCGCL used for therapeutic administration can be sterile. Sterility is readily accomplished by filtration through sterile filtration membranes (e.g., 0.2 micron membranes). Therapeutic polypeptide compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

CRCGCL polypeptides ordinarily will be stored in unit or multi-dose containers, for example, sealed ampoules or vials, as an aqueous solution or as a lyophilized formulation for reconstitution. As an example of a lyophilized formulation, 10-ml vials are filled with 5 ml of sterile-filtered 1% (w/v) aqueous CRCGCL polypeptide solution, and the resulting mixture is lyophilized. The infusion solution is prepared by reconstituting the lyophilized CRCGCL polypeptide using bacteriostatic Water-for-Injection.

The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration. In addition, CRCGCL may be employed in conjunction with other therapeutic compounds.

5  
10

15  
20  
25  
30

30

not limited to, steroids, cyclosporine, cyclosporine analogs, cyclophosphamide methylprednisone, prednisone, azathioprine, FK-506, 15-deoxyspergualin, and other immunosuppressive agents that act by suppressing the function of responding T cells.

5 In a further embodiment, the compositions of the invention are administered in combination with an antibiotic agent. Antibiotic agents that may be administered with the compositions of the invention include, but are not limited to, tetracycline, metronidazole, amoxicillin, beta-lactamases, aminoglycosides, macrolides, quinolones, fluoroquinolones, cephalosporins, erythromycin, ciprofloxacin, and streptomycin.

10 In an additional embodiment, the compositions of the invention are administered alone or in combination with an anti-inflammatory agent. Anti-inflammatory agents that may be administered with the compositions of the invention include, but are not limited to, glucocorticoids and the nonsteroidal anti-inflammatories, aminoarylcarboxylic acid derivatives, arylacetic acid derivatives,  
15 arylbutyric acid derivatives, arylcarboxylic acids, arylpropionic acid derivatives, pyrazoles, pyrazolones, salicylic acid derivatives, thiazinecarboxamides, e-acetamidocaproic acid, S-adenosylmethionine, 3-amino-4-hydroxybutyric acid, amixetrine, bendazac, benzydamine, bucolome, difenpiramide, ditazol, emorfazone,  
20 guaiazulene, nabumetone, nimesulide, orgotein, oxaceprol, paranyline, perisoxal, pifoxime, proquazone, proxazole, and tenidap.

In another embodiment, compositions of the invention are administered in combination with a chemotherapeutic agent. Chemotherapeutic agents that may be administered with the compositions of the invention include, but are not limited to, antibiotic derivatives (e.g., doxorubicin, bleomycin, daunorubicin, and  
25 dactinomycin); antiestrogens (e.g., tamoxifen); antimetabolites (e.g., fluorouracil, 5-FU, methotrexate, floxuridine, interferon alpha-2b, glutamic acid, plicamycin, mercaptopurine, and 6-thioguanine); cytotoxic agents (e.g., carmustine, BCNU, lomustine, CCNU, cytosine arabinoside, cyclophosphamide, estramustine, hydroxyurea, procarbazine, mitomycin, busulfan, cis-platin, and vincristine sulfate);  
30 hormones (e.g., medroxyprogesterone, estramustine phosphate sodium, ethinyl estradiol, estradiol, megestrol acetate, methyltestosterone, diethylstilbestrol diphosphate, chlorotrianisene, and testolactone); nitrogen mustard derivatives (e.g., mephallen, chorambucil, mechlorethamine (nitrogen mustard) and thiotepa); steroids

and combinations (e.g., bethamethasone sodium phosphate); and others (e.g., dicarbazine, asparaginase, mitotane, vincristine sulfate, vinblastine sulfate, and etoposide).

5 In an additional embodiment, the compositions of the invention are administered in combination with cytokines. Cytokines that may be administered with the compositions of the invention include, but are not limited to, IL2, IL3, IL4, IL5, IL6, IL7, IL10, IL12, IL13, IL15, anti-CD40, CD40L, IFN-gamma and TNF-alpha.

10 In an additional embodiment, the compositions of the invention are administered in combination with angiogenic proteins. Angiogenic proteins that may be administered with the compositions of the invention include, but are not limited to, Glioma Derived Growth Factor (GDGF), as disclosed in European Patent Number EP-399816; Platelet Derived Growth Factor-A (PDGF-A), as disclosed in European Patent Number EP-682110; Platelet Derived Growth Factor-B (PDGF-B), as  
15 disclosed in European Patent Number EP-282317; Placental Growth Factor (PIGF), as disclosed in International Publication Number WO 92/06194; Placental Growth Factor-2 (PIGF-2), as disclosed in Hauser et al., Growth Factors, 4:259-268 (1993); Vascular Endothelial Growth Factor (VEGF), as disclosed in International Publication  
Number WO 90/13649; Vascular Endothelial Growth Factor-A (VEGF-A), as  
20 disclosed in European Patent Number EP-506477; Vascular Endothelial Growth Factor-2 (VEGF-2), as disclosed in International Publication Number WO 96/39515; Vascular Endothelial Growth Factor B-186 (VEGF-B186), as disclosed in International Publication Number WO 96/26736; Vascular Endothelial Growth Factor-D (VEGF-D), as disclosed in International Publication Number WO 98/02543;  
25 Vascular Endothelial Growth Factor-D (VEGF-D), as disclosed in International Publication Number WO 98/07832; and Vascular Endothelial Growth Factor-E (VEGF-E), as disclosed in German Patent Number DE19639601. The above mentioned references are incorporated herein by reference herein.

30 In an additional embodiment, the compositions of the invention are administered in combination with Fibroblast Growth Factors. Fibroblast Growth Factors that may be administered with the compositions of the invention include, but are not limited to, FGF-1, FGF-2, FGF-3, FGF-4, FGF-5, FGF-6, FGF-7, FGF-8, FGF-9, FGF-10, FGF-11, FGF-12, FGF-13, FGF-14, and FGF-15.

65280\*0009260

In additional embodiments, the compositions of the invention are administered in combination with other therapeutic or prophylactic regimens, such as, for example, radiation therapy.

5

***Example 25: Method of Treating Decreased Levels of CRCGCL***

The present invention relates to a method for treating an individual in need of a decreased level of CRCGCL activity in the body comprising, administering to such  
10 an individual a composition comprising a therapeutically effective amount of CRCGCL antagonist. Preferred antagonists for use in the present invention are CRCGCL-specific antibodies.

Moreover, it will be appreciated that conditions caused by a decrease in the standard or normal expression level of CRCGCL in an individual can be treated by  
15 administering CRCGCL, preferably in the secreted form. Thus, the invention also provides a method of treatment of an individual in need of an increased level of CRCGCL polypeptide comprising administering to such an individual a pharmaceutical composition comprising an amount of CRCGCL to increase the activity level of CRCGCL in such an individual.

20 For example, a patient with decreased levels of CRCGCL polypeptide receives a daily dose 0.1-100 ug/kg of the polypeptide for six consecutive days. Preferably, the polypeptide is in the secreted form. The exact details of the dosing scheme, based on administration and formulation, are provided in Example 24.

25 ***Example 26: Method of Treating Increased Levels of CRCGCL***

The present invention also relates to a method for treating an individual in need of an increased level of CRCGCL activity in the body comprising administering to such an individual a composition comprising a therapeutically effective amount of  
30 CRCGCL or an agonist thereof.

Antisense technology is used to inhibit production of CRCGCL. This technology is one example of a method of decreasing levels of CRCGCL polypeptide, preferably a secreted form, due to a variety of etiologies, such as cancer.

09376430-081899

For example, a patient diagnosed with abnormally increased levels of CRCGCL is administered intravenously antisense polynucleotides at 0.5, 1.0, 1.5, 2.0 and 3.0 mg/kg day for 21 days. This treatment is repeated after a 7-day rest period if the treatment was well tolerated. The formulation of the antisense polynucleotide is provided in Example 24.

### ***Example 27: Method of Treatment Using Gene Therapy - Ex Vivo***

One method of gene therapy transplants fibroblasts, which are capable of expressing CRCGCL polypeptides, onto a patient. Generally, fibroblasts are obtained from a subject by skin biopsy. The resulting tissue is placed in tissue-culture medium and separated into small pieces. Small chunks of the tissue are placed on a wet surface of a tissue culture flask, approximately ten pieces are placed in each flask. The flask is turned upside down, closed tight and left at room temperature over night. After 24 hours at room temperature, the flask is inverted and the chunks of tissue remain fixed to the bottom of the flask and fresh media (e.g., Ham's F12 media, with 10% FBS, penicillin and streptomycin) is added. The flasks are then incubated at 37 degree C for approximately one week.

At this time, fresh media is added and subsequently changed every several days. After an additional two weeks in culture, a monolayer of fibroblasts emerge. The monolayer is trypsinized and scaled into larger flasks.

pMV-7 (Kirschmeier, P.T. et al., DNA, 7:219-25 (1988)), flanked by the long terminal repeats of the Moloney murine sarcoma virus, is digested with EcoRI and HindIII and subsequently treated with calf intestinal phosphatase. The linear vector is fractionated on agarose gel and purified, using glass beads.

The cDNA encoding CRCGCL can be amplified using PCR primers which correspond to the 5' and 3' end sequences respectively as set forth in Example 1. Preferably, the 5' primer contains an EcoRI site and the 3' primer includes a HindIII site. Equal quantities of the Moloney murine sarcoma virus linear backbone and the amplified EcoRI and HindIII fragment are added together, in the presence of T4 DNA ligase. The resulting mixture is maintained under conditions appropriate for ligation of the two fragments. The ligation mixture is then used to transform bacteria HB101,



which are then plated onto agar containing kanamycin for the purpose of confirming that the vector contains properly inserted CRCGCL.

5 The amphotropic pA317 or GP+am12 packaging cells are grown in tissue culture to confluent density in Dulbecco's Modified Eagles Medium (DMEM) with 10% calf serum (CS), penicillin and streptomycin. The MSV vector containing the CRCGCL gene is then added to the media and the packaging cells transduced with the vector. The packaging cells now produce infectious viral particles containing the CRCGCL gene(the packaging cells are now referred to as producer cells).

10 Fresh media is added to the transduced producer cells, and subsequently, the media is harvested from a 10 cm plate of confluent producer cells. The spent media, containing the infectious viral particles, is filtered through a millipore filter to remove detached producer cells and this media is then used to infect fibroblast cells. Media is removed from a sub-confluent plate of fibroblasts and quickly replaced with the media from the producer cells. This media is removed and replaced with fresh media.  
15 If the titer of virus is high, then virtually all fibroblasts will be infected and no selection is required. If the titer is very low, then it is necessary to use a retroviral vector that has a selectable marker, such as neo or his. Once the fibroblasts have been efficiently infected, the fibroblasts are analyzed to determine whether CRCGCL protein is produced.

20 The engineered fibroblasts are then transplanted onto the host, either alone or after having been grown to confluence on cytodex 3 microcarrier beads.

### ***Example 28: Gene Therapy Using Endogenous CRCGCL Gene***

25 Another method of gene therapy according to the present invention involves operably associating the endogenous CRCGCL sequence with a promoter via homologous recombination as described, for example, in U.S. Patent No. 5,641,670, issued June 24, 1997; International Publication No. WO 96/29411, published September 26, 1996; International Publication No. WO 94/12650, published August  
30 4, 1994; Koller et al., Proc. Natl. Acad. Sci. USA 86:8932-8935 (1989); and Zijlstra et al., Nature 342:435-438 (1989). This method involves the activation of a gene which is present in the target cells, but which is not expressed in the cells, or is expressed at a lower level than desired.

The amplified promoter and the amplified targeting sequences are digested with the appropriate restriction enzymes and subsequently treated with calf intestinal phosphatase. The digested promoter and digested targeting sequences are added together in the presence of T4 DNA ligase. The resulting mixture is maintained under conditions appropriate for ligation of the two fragments. The construct is size fractionated on an agarose gel then purified by phenol extraction and ethanol precipitation.

Once the cells are transfected, homologous recombination will take place which results in the promoter being operably linked to the endogenous CRCGCL sequence. This results in the expression of CRCGCL in the cell. Expression may be detected by immunological staining, or any other method known in the art.

Fibroblasts are obtained from a subject by skin biopsy. The resulting tissue is placed in DMEM + 10% fetal calf serum. Exponentially growing or early stationary phase fibroblasts are trypsinized and rinsed from the plastic surface with nutrient medium. An aliquot of the cell suspension is removed for counting, and the remaining cells are subjected to centrifugation. The supernatant is aspirated and the pellet is resuspended in 5 ml of electroporation buffer (20 mM HEPES pH 7.3, 137 mM NaCl, 5 mM KCl, 0.7 mM Na<sub>2</sub> HPO<sub>4</sub>, 6 mM dextrose). The cells are recentrifuged, the

supernatant aspirated, and the cells resuspended in electroporation buffer containing 1 mg/ml acetylated bovine serum albumin. The final cell suspension contains approximately  $3 \times 10^6$  cells/ml. Electroporation should be performed immediately following resuspension.

5 Plasmid DNA is prepared according to standard techniques. For example, to construct a plasmid for targeting to the CRCGCL locus, plasmid pUC18 (MBI Fermentas, Amherst, NY) is digested with HindIII. The CMV promoter is amplified by PCR with an XbaI site on the 5' end and a BamHI site on the 3' end. Two CRCGCL non-coding sequences are amplified via PCR: one CRCGCL non-coding  
10 sequence (CRCGCL fragment 1) is amplified with a HindIII site at the 5' end and an Xba site at the 3' end; the other CRCGCL non-coding sequence (CRCGCL fragment 2) is amplified with a BamHI site at the 5' end and a HindIII site at the 3' end. The CMV promoter and CRCGCL fragments are digested with the appropriate enzymes (CMV promoter - XbaI and BamHI; CRCGCL fragment 1 - XbaI; CRCGCL  
15 fragment 2 - BamHI) and ligated together. The resulting ligation product is digested with HindIII, and ligated with the HindIII-digested pUC18 plasmid.

Plasmid DNA is added to a sterile cuvette with a 0.4 cm electrode gap (Bio-Rad). The final DNA concentration is generally at least 120  $\mu$ g/ml. 0.5 ml of the cell suspension (containing approximately  $1.5 \times 10^6$  cells) is then added to the cuvette,  
20 and the cell suspension and DNA solutions are gently mixed. Electroporation is performed with a Gene-Pulser apparatus (Bio-Rad). Capacitance and voltage are set at 960  $\mu$ F and 250-300 V, respectively. As voltage increases, cell survival decreases, but the percentage of surviving cells that stably incorporate the introduced DNA into their genome increases dramatically. Given these parameters, a pulse time of  
25 approximately 14-20 mSec should be observed.

Electroporated cells are maintained at room temperature for approximately 5 min, and the contents of the cuvette are then gently removed with a sterile transfer pipette. The cells are added directly to 10 ml of prewarmed nutrient media (DMEM with 15% calf serum) in a 10 cm dish and incubated at 37 degree C. The following  
30 day, the media is aspirated and replaced with 10 ml of fresh media and incubated for a further 16-24 hours.

The engineered fibroblasts are then injected into the host, either alone or after having been grown to confluence on cytodex 3 microcarrier beads. The fibroblasts

668730-06192660

5

## 10

20

25

30

The CRCGCL polynucleotide vector constructs used in the gene therapy method are preferably constructs that will not integrate into the host genome nor will

they contain sequences that allow for replication. Any strong promoter known to those skilled in the art can be used for driving the expression of DNA. Unlike other gene therapies techniques, one major advantage of introducing naked nucleic acid sequences into target cells is the transitory nature of the polynucleotide synthesis in the cells. Studies have shown that non-replicating DNA sequences can be introduced into cells to provide production of the desired polypeptide for periods of up to six months.

The CRCGCL polynucleotide construct can be delivered to the interstitial space of tissues within the an animal, including of muscle, skin, brain, lung, liver, spleen, bone marrow, thymus, heart, lymph, blood, bone, cartilage, pancreas, kidney, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland, and connective tissue. Interstitial space of the tissues comprises the intercellular fluid, mucopolysaccharide matrix among the reticular fibers of organ tissues, elastic fibers in the walls of vessels or chambers, collagen fibers of fibrous tissues, or that same matrix within connective tissue ensheathing muscle cells or in the lacunae of bone. It is similarly the space occupied by the plasma of the circulation and the lymph fluid of the lymphatic channels. Delivery to the interstitial space of muscle tissue is preferred for the reasons discussed below. They may be conveniently

delivered by injection into the tissues comprising these cells. They are preferably delivered to and expressed in persistent, non-dividing cells which are differentiated, although delivery and expression may be achieved in non-differentiated or less completely differentiated cells, such as, for example, stem cells of blood or skin fibroblasts. In vivo muscle cells are particularly competent in their ability to take up and express polynucleotides.

For the naked CRCGCL polynucleotide injection, an effective dosage amount of DNA or RNA will be in the range of from about 0.05 g/kg body weight to about 50 mg/kg body weight. Preferably the dosage will be from about 0.005 mg/kg to about 20 mg/kg and more preferably from about 0.05 mg/kg to about 5 mg/kg. Of course, as the artisan of ordinary skill will appreciate, this dosage will vary according to the tissue site of injection. The appropriate and effective dosage of nucleic acid sequence can readily be determined by those of ordinary skill in the art and may depend on the condition being treated and the route of administration. The preferred route of administration is by the parenteral route of injection into the interstitial space of

tissues. However, other parenteral routes may also be used, such as, inhalation of an aerosol formulation particularly for delivery to lungs or bronchial tissues, throat or mucous membranes of the nose. In addition, naked CRCGCL polynucleotide constructs can be delivered to arteries during angioplasty by the catheter used in the procedure.

The dose response effects of injected CRCGCL polynucleotide in muscle in vivo is determined as follows. Suitable CRCGCL template DNA for production of mRNA coding for CRCGCL polypeptide is prepared in accordance with a standard recombinant DNA methodology. The template DNA, which may be either circular or linear, is either used as naked DNA or complexed with liposomes. The quadriceps muscles of mice are then injected with various amounts of the template DNA.

Five to six week old female and male Balb/C mice are anesthetized by intraperitoneal injection with 0.3 ml of 2.5% Avertin. A 1.5 cm incision is made on the anterior thigh, and the quadriceps muscle is directly visualized. The CRCGCL template DNA is injected in 0.1 ml of carrier in a 1 cc syringe through a 27 gauge needle over one minute, approximately 0.5 cm from the distal insertion site of the muscle into the knee and about 0.2 cm deep. A suture is placed over the injection site for future localization, and the skin is closed with stainless steel clips.

After an appropriate incubation time (e.g., 7 days) muscle extracts are prepared by excising the entire quadriceps. Every fifth 15 um cross-section of the individual quadriceps muscles is histochemically stained for CRCGCL protein expression. A time course for CRCGCL protein expression may be done in a similar fashion except that quadriceps from different mice are harvested at different times. Persistence of CRCGCL DNA in muscle following injection may be determined by Southern blot analysis after preparing total cellular DNA and HIRT supernatants from injected and control mice. The results of the above experimentation in mice can be use to extrapolate proper dosages and other treatment parameters in humans and other animals using CRCGCL naked DNA.

### ***Example 30: CRCGCL Transgenic Animals.***

The CRCGCL polypeptides can also be expressed in transgenic animals. Animals of any species, including, but not limited to, mice, rats, rabbits, hamsters,

09376430-081899

guinea pigs, pigs, micro-pigs, goats, sheep, cows and non-human primates, e.g., baboons, monkeys, and chimpanzees may be used to generate transgenic animals. In a specific embodiment, techniques described herein or otherwise known in the art, are used to express polypeptides of the invention in humans, as part of a gene therapy protocol.

Any technique known in the art may be used to introduce the transgene (i.e., polynucleotides of the invention) into animals to produce the founder lines of transgenic animals. Such techniques include, but are not limited to, pronuclear microinjection (Paterson et al., Appl. Microbiol. Biotechnol. 40:691-698 (1994); Carver et al., Biotechnology (NY) 11:1263-1270 (1993); Wright et al., Biotechnology (NY) 9:830-834 (1991); and Hoppe et al., U.S. Pat. No. 4,873,191 (1989)); retrovirus mediated gene transfer into germ lines (Van der Putten et al., Proc. Natl. Acad. Sci., USA 82:6148-6152 (1985)), blastocysts or embryos; gene targeting in embryonic stem cells (Thompson et al., Cell 56:313-321 (1989)); electroporation of cells or embryos (Lo, 1983, Mol Cell. Biol. 3:1803-1814 (1983)); introduction of the polynucleotides of the invention using a gene gun (see, e.g., Ulmer et al., Science 259:1745 (1993); introducing nucleic acid constructs into embryonic pluripotent stem cells and transferring the stem cells back into the blastocyst; and sperm-mediated gene transfer (Lavitrano et al., Cell 57:717-723 (1989); etc. For a review of such techniques, see Gordon, "Transgenic Animals," Intl. Rev. Cytol. 115:171-229 (1989), which is incorporated by reference herein in its entirety.

Any technique known in the art may be used to produce transgenic clones containing polynucleotides of the invention, for example, nuclear transfer into enucleated oocytes of nuclei from cultured embryonic, fetal, or adult cells induced to quiescence (Campell et al., Nature 380:64-66 (1996); Wilmut et al., Nature 385:810-813 (1997)).

The present invention provides for transgenic animals that carry the transgene in all their cells, as well as animals which carry the transgene in some, but not all their cells, i.e., mosaic animals or chimeric. The transgene may be integrated as a single transgene or as multiple copies such as in concatamers, e.g., head-to-head tandems or head-to-tail tandems. The transgene may also be selectively introduced into and activated in a particular cell type by following, for example, the teaching of Lasko et al. (Lasko et al., Proc. Natl. Acad. Sci. USA 89:6232-6236 (1992)). The regulatory

sequences required for such a cell-type specific activation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art. When it is desired that the polynucleotide transgene be integrated into the chromosomal site of the endogenous gene, gene targeting is preferred.

5 Briefly, when such a technique is to be utilized, vectors containing some nucleotide sequences homologous to the endogenous gene are designed for the purpose of integrating, via homologous recombination with chromosomal sequences, into and disrupting the function of the nucleotide sequence of the endogenous gene. The transgene may also be selectively introduced into a particular cell type, thus  
10 inactivating the endogenous gene in only that cell type, by following, for example, the teaching of Gu et al. (Gu et al., Science 265:103-106 (1994)). The regulatory sequences required for such a cell-type specific inactivation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art. The contents of each of the documents recited in this paragraph is herein incorporated by  
15 reference in its entirety.

Any of the CRCGCL polypeptides disclose throughout this application can be used to generate transgenic animals. For example, DNA encoding amino acids M1-K231 of SEQ ID NO:2 can be inserted into a vector containing a promoter, such as the actin promoter, which will ubiquitously express the inserted fragment. Primers  
20 that can be used to generate such fragments include a 5' primer containing a BglII restriction site shown in bold:

CCGGTTAGATCTGCCATCATGGGGCGGCTGGTTCTGCCGGTTAGATCTGC  
CATCATGGGGCGGCTGGTTCTGCCGGTTAGATCTGCCATCATGGGGCGGC  
TGGTTCTGCCGGTTAGATCTGCCATCATGGGGCGGCTGGTTCTG (SEQ ID  
25 NO: 32) and a 3' primer, containing a Xba restriction site shown in bold:

GGCCGGTCTAGATTATTTGGACAGCTTTGGTTTG (SEQ ID NO: 31). This construct will express the predicted soluble domain of CRCGCL under the control of the actin promoter for ubiquitous expression. The region of CRCGCL included in this construct extends from M1-K231 of SEQ ID NO:2.

30 Similarly, the DNA encoding the full length CRCGCL protein can also be inserted into a vector.

Alternatively, polynucleotides of the invention can be inserted in a vector which controls tissue specific expression through a tissue specific promoter. For

658130-0619260



example, a construct having a transferrin promoter would express the CRCGCL polypeptide in the liver of transgenic animals. Therefore, DNA encoding amino acids M1-K231 of SEQ ID NO:2 can be amplified using a 5' primer, having a BglII restriction site shown in bold:

- 5 CCGGTTAGATCTGCCATCATGGGGCGGCTGGTTCTG (SEQ ID NO: 28), and a 3' primer, containing a Xba restriction site shown in bold: GGCCGGTCTAGATTATTTGGACAGCTTTGGTTTG (SEQ ID NO: 31).

Similarly, the DNA encoding the full length CRCGCL protein can also be inserted into a vector for tissue specific expression.

- 10 Once transgenic animals have been generated, the expression of the recombinant gene may be assayed utilizing standard techniques. Initial screening may be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to verify that integration of the transgene has taken place. The level of mRNA expression of the transgene in the tissues of the transgenic animals may also be  
15 assessed using techniques which include, but are not limited to, Northern blot analysis of tissue samples obtained from the animal, in situ hybridization analysis, and reverse transcriptase-PCR (rt-PCR). Samples of transgenic gene-expressing tissue may also be evaluated immunocytochemically or immunohistochemically using antibodies specific for the transgene product.

- 20 Once the founder animals are produced, they may be bred, inbred, outbred, or crossbred to produce colonies of the particular animal. Examples of such breeding strategies include, but are not limited to: outbreeding of founder animals with more than one integration site in order to establish separate lines; inbreeding of separate lines in order to produce compound transgenics that express the transgene at higher  
25 levels because of the effects of additive expression of each transgene; crossing of heterozygous transgenic animals to produce animals homozygous for a given integration site in order to both augment expression and eliminate the need for screening of animals by DNA analysis; crossing of separate homozygous lines to produce compound heterozygous or homozygous lines; and breeding to place the  
30 transgene on a distinct background that is appropriate for an experimental model of interest.

Transgenic animals of the invention have uses which include, but are not limited to, animal model systems useful in elaborating the biological function of

0936430-049260

CRCGCL polypeptides, studying conditions and/or disorders associated with aberrant CRCGCL expression, and in screening for compounds effective in ameliorating such conditions and/or disorders.

5

***Example 31: CRCGCL Knock-Out Animals.***

Endogenous CRCGCL gene expression can also be reduced by inactivating or "knocking out" the CRCGCL gene and/or its promoter using targeted homologous recombination. (E.g., see Smithies et al., Nature 317:230-234 (1985); Thomas & Capecchi, Cell 51:503-512 (1987); Thompson et al., Cell 5:313-321 (1989); each of which is incorporated by reference herein in its entirety). For example, a mutant, non-functional polynucleotide of the invention (or a completely unrelated DNA sequence) flanked by DNA homologous to the endogenous polynucleotide sequence (either the coding regions or regulatory regions of the gene) can be used, with or without a selectable marker and/or a negative selectable marker, to transfect cells that express polypeptides of the invention in vivo. In another embodiment, techniques known in the art are used to generate knockouts in cells that contain, but do not express the gene of interest. Insertion of the DNA construct, via targeted homologous recombination, results in inactivation of the targeted gene. Such approaches are particularly suited in research and agricultural fields where modifications to embryonic stem cells can be used to generate animal offspring with an inactive targeted gene (e.g., see Thomas & Capecchi 1987 and Thompson 1989, supra). However this approach can be routinely adapted for use in humans provided the recombinant DNA constructs are directly administered or targeted to the required site in vivo using appropriate viral vectors that will be apparent to those of skill in the art.

In further embodiments of the invention, cells that are genetically engineered to express the polypeptides of the invention, or alternatively, that are genetically engineered not to express the polypeptides of the invention (e.g., knockouts) are administered to a patient in vivo. Such cells may be obtained from the patient (i.e., animal, including human) or an MHC compatible donor and can include, but are not limited to fibroblasts, bone marrow cells, blood cells (e.g., lymphocytes), adipocytes, muscle cells, endothelial cells etc. The cells are genetically engineered in vitro using recombinant DNA techniques to introduce the coding sequence of polypeptides of the

09376430-081899

invention into the cells, or alternatively, to disrupt the coding sequence and/or endogenous regulatory sequence associated with the polypeptides of the invention, e.g., by transduction (using viral vectors, and preferably vectors that integrate the transgene into the cell genome) or transfection procedures, including, but not limited to, the use of plasmids, cosmids, YACs, naked DNA, electroporation, liposomes, etc. The coding sequence of the polypeptides of the invention can be placed under the control of a strong constitutive or inducible promoter or promoter/enhancer to achieve expression, and preferably secretion, of the CRCGCL polypeptides. The engineered cells which express and preferably secrete the polypeptides of the invention can be introduced into the patient systemically, e.g., in the circulation, or intraperitoneally.

Alternatively, the cells can be incorporated into a matrix and implanted in the body, e.g., genetically engineered fibroblasts can be implanted as part of a skin graft; genetically engineered endothelial cells can be implanted as part of a lymphatic or vascular graft. (See, for example, Anderson et al. U.S. Patent No. 5,399,349; and Mulligan & Wilson, U.S. Patent No. 5,460,959 each of which is incorporated by reference herein in its entirety).

When the cells to be administered are non-autologous or non-MHC compatible cells, they can be administered using well known techniques which prevent the development of a host immune response against the introduced cells. For example, the cells may be introduced in an encapsulated form which, while allowing for an exchange of components with the immediate extracellular environment, does not allow the introduced cells to be recognized by the host immune system.

Knock-out animals of the invention have uses which include, but are not limited to, animal model systems useful in elaborating the biological function of CRCGCL polypeptides, studying conditions and/or disorders associated with aberrant CRCGCL expression, and in screening for compounds effective in ameliorating such conditions and/or disorders. For example, a knock-out mouse can be made using the sequences disclosed as AA008694 and W98372, herein incorporated by reference in their entirety.

### ***Example 32: Assays Detecting Stimulation or Inhibition of B cell Proliferation and Differentiation***

Generation of functional humoral immune responses requires both soluble and  
 5 cognate signaling between B-lineage cells and their microenvironment. Signals may  
 impart a positive stimulus that allows a B-lineage cell to continue its programmed  
 development, or a negative stimulus that instructs the cell to arrest its current  
 developmental pathway. To date, numerous stimulatory and inhibitory signals have been  
 found to influence B cell responsiveness including IL-2, IL-4, IL-5, IL-6, IL-7, IL10, IL-  
 10 13, IL-14 and IL-15. Interestingly, these signals are by themselves weak effectors but can,  
 in combination with various co-stimulatory proteins, induce activation, proliferation,  
 differentiation, homing, tolerance and death among B cell populations.

One of the best studied classes of B-cell co-stimulatory proteins is the TNF-  
 superfamily. Within this family CD40, CD27, and CD30 along with their respective  
 15 ligands CD154, CD70, and CD153 have been found to regulate a variety of immune  
 responses. Assays which allow for the detection and/or observation of the proliferation  
 and differentiation of these B-cell populations and their precursors are valuable tools in  
 determining the effects various proteins may have on these B-cell populations in terms of  
 proliferation and differentiation. Listed below are two assays designed to allow for the  
 20 detection of the differentiation, proliferation, or inhibition of B-cell populations and their  
 precursors.

**In Vitro Assay-** Purified CRCGCL protein, or truncated forms thereof, is  
 assessed for its ability to induce activation, proliferation, differentiation or inhibition  
 and/or death in B-cell populations and their precursors. The activity of CRCGCL protein  
 25 on purified human tonsillar B cells, measured qualitatively over the dose range from 0.1  
 to 10,000 ng/mL, is assessed in a standard B-lymphocyte co-stimulation assay in which  
 purified tonsillar B cells are cultured in the presence of either formalin-fixed  
 Staphylococcus aureus Cowan I (SAC) or immobilized anti-human IgM antibody as the  
 priming agent. Second signals such as IL-2 and IL-15 synergize with SAC and IgM  
 30 crosslinking to elicit B cell proliferation as measured by tritiated-thymidine  
 incorporation. Novel synergizing agents can be readily identified using this assay. The  
 assay involves isolating human tonsillar B cells by magnetic bead (MACS) depletion of

0536430-001899

CD3-positive cells. The resulting cell population is greater than 95% B cells as assessed by expression of CD45R(B220).

Various dilutions of each sample are placed into individual wells of a 96-well plate to which are added  $10^5$  B-cells suspended in culture medium (RPMI 1640 containing 10% FBS,  $5 \times 10^{-5}$  M 2ME, 100U/ml penicillin, 10ug/ml streptomycin, and  $10^{-5}$  dilution of SAC) in a total volume of 150ul. Proliferation or inhibition is quantitated by a 20h pulse (1uCi/well) with 3H-thymidine (6.7 Ci/mM) beginning 72h post factor addition. The positive and negative controls are IL2 and medium respectively.

**In Vivo Assay-** BALB/c mice are injected (i.p.) twice per day with buffer only, or 2 mg/Kg of CRCGCL protein, or truncated forms thereof. Mice receive this treatment for 4 consecutive days, at which time they are sacrificed and various tissues and serum collected for analyses. Comparison of H&E sections from normal and CRCGCL protein-treated spleens identify the results of the activity of CRCGCL protein on spleen cells, such as the diffusion of peri-arterial lymphatic sheaths, and/or significant increases in the nucleated cellularity of the red pulp regions, which may indicate the activation of the differentiation and proliferation of B-cell populations. Immunohistochemical studies using a B cell marker, anti-CD45R(B220), are used to determine whether any physiological changes to splenic cells, such as splenic disorganization, are due to increased B-cell representation within loosely defined B-cell zones that infiltrate established T-cell regions.

Flow cytometric analyses of the spleens from CRCGCL protein-treated mice is used to indicate whether CRCGCL protein specifically increases the proportion of ThB+, CD45R(B220)dull B cells over that which is observed in control mice.

Likewise, a predicted consequence of increased mature B-cell representation in vivo is a relative increase in serum Ig titers. Accordingly, serum IgM and IgA levels are compared between buffer and CRCGCL protein-treated mice.

The studies described in this example tested activity in CRCGCL protein. However, one skilled in the art could easily modify the exemplified studies to test the activity of CRCGCL polynucleotides (e.g., gene therapy), agonists, and/or antagonists of CRCGCL.

### ***Example 33: T Cell Proliferation Assay***

A CD3-induced proliferation assay is performed on PBMCs and is measured by the uptake of <sup>3</sup>H-thymidine. The assay is performed as follows. Ninety-six well plates are coated with 100 µl/well of mAb to CD3 (HIT3a, Pharmingen) or isotype-matched control mAb (B33.1) overnight at 4°C (1 µg/ml in .05M bicarbonate buffer, pH 9.5), then washed three times with PBS. PBMC are isolated by F/H gradient centrifugation from human peripheral blood and added to quadruplicate wells (5 x 10<sup>4</sup>/well) of mAb coated plates in RPMI containing 10% FCS and P/S in the presence of varying concentrations of CRGCL protein (total volume 200 µl). Relevant protein buffer and medium alone are controls. After 48 hr. culture at 37°C, plates are spun for 2 min. at 1000 rpm and 100 µl of supernatant is removed and stored -20°C for measurement of IL-2 (or other cytokines) if effect on proliferation is observed. Wells are supplemented with 100 µl of medium containing 0.5 µCi of <sup>3</sup>H-thymidine and cultured at 37°C for 18-24 hr. Wells are harvested and incorporation of <sup>3</sup>H-thymidine used as a measure of proliferation. Anti-CD3 alone is the positive control for proliferation. IL-2 (100 U/ml) is also used as a control which enhances proliferation. Control antibody which does not induce proliferation of T cells is used as the negative controls for the effects of CRGCL proteins.

The studies described in this example tested activity in CRGCL protein. However, one skilled in the art could easily modify the exemplified studies to test the activity of CRGCL polynucleotides (e.g., gene therapy), agonists, and/or antagonists of CRGCL.

### ***Example 34: Effect of CRGCL on the Expression of MHC Class II, Costimulatory and Adhesion Molecules and Cell Differentiation of Monocytes and Monocyte-Derived Human Dendritic Cells***

Dendritic cells are generated by the expansion of proliferating precursors found in the peripheral blood: adherent PBMC or elutriated monocytic fractions are cultured for 7-10 days with GM-CSF (50 ng/ml) and IL-4 (20 ng/ml). These dendritic cells have the characteristic phenotype of immature cells (expression of CD1, CD80, CD86, CD40 and

MHC class II antigens). Treatment with activating factors, such as TNF- $\alpha$ , causes a rapid change in surface phenotype (increased expression of MHC class I and II, costimulatory and adhesion molecules, downregulation of FC $\gamma$ RII, upregulation of CD83). These changes correlate with increased antigen-presenting capacity and with functional maturation of the dendritic cells.

FACS analysis of surface antigens is performed as follows. Cells are treated 1-3 days with increasing concentrations of CRCGCL or LPS (positive control), washed with PBS containing 1% BSA and 0.02 mM sodium azide, and then incubated with 1:20 dilution of appropriate FITC- or PE-labeled monoclonal antibodies for 30 minutes at 4°C. After an additional wash, the labeled cells are analyzed by flow cytometry on a FACScan (Becton Dickinson).

Effect on the production of cytokines. Cytokines generated by dendritic cells, in particular IL-12, are important in the initiation of T-cell dependent immune responses. IL-12 strongly influences the development of Th1 helper T-cell immune response, and induces cytotoxic T and NK cell function. An ELISA is used to measure the IL-12 release as follows. Dendritic cells ( $10^6$ /ml) are treated with increasing concentrations of CRCGCL for 24 hours. LPS (100 ng/ml) is added to the cell culture as positive control. Supernatants from the cell cultures are then collected and analyzed for IL-12 content using commercial ELISA kit (e.g., R & D Systems (Minneapolis, MN)). The standard protocols provided with the kits are used.

Effect on the expression of MHC Class II, costimulatory and adhesion molecules. Three major families of cell surface antigens can be identified on monocytes: adhesion molecules, molecules involved in antigen presentation, and Fc receptor. Modulation of the expression of MHC class II antigens and other costimulatory molecules, such as B7 and ICAM-1, may result in changes in the antigen presenting capacity of monocytes and ability to induce T cell activation. Increase expression of Fc receptors may correlate with improved monocyte cytotoxic activity, cytokine release and phagocytosis.

FACS analysis is used to examine the surface antigens as follows. Monocytes are treated 1-5 days with increasing concentrations of CRCGCL or LPS (positive control), washed with PBS containing 1% BSA and 0.02 mM sodium azide, and then incubated with 1:20 dilution of appropriate FITC- or PE-labeled monoclonal antibodies for 30

09376430-031399

minutes at 4°C. After an additional wash, the labeled cells are analyzed by flow cytometry on a FACScan (Becton Dickinson).

Monocyte activation and/or increased survival. Assays for molecules that activate (or alternatively, inactivate) monocytes and/or increase monocyte survival (or alternatively, decrease monocyte survival) are known in the art and may routinely be applied to determine whether a molecule of the invention functions as an inhibitor or activator of monocytes. CRCGCL, agonists, or antagonists of CRCGCL can be screened using the three assays described below. For each of these assays, Peripheral blood mononuclear cells (PBMC) are purified from single donor leukopacks (American Red Cross, Baltimore, MD) by centrifugation through a Histopaque gradient (Sigma). Monocytes are isolated from PBMC by counterflow centrifugal elutriation.

Monocyte Survival Assay. Human peripheral blood monocytes progressively lose viability when cultured in absence of serum or other stimuli. Their death results from internally regulated process (apoptosis). Addition to the culture of activating factors, such as TNF-alpha dramatically improves cell survival and prevents DNA fragmentation. ~~Propidium iodide (PI) staining is used to measure apoptosis as follows.~~ Monocytes are cultured for 48 hours in polypropylene tubes in serum-free medium (positive control), in the presence of 100 ng/ml TNF-alpha (negative control), and in the presence of varying concentrations of the compound to be tested. Cells are suspended at a concentration of  $2 \times 10^6$ /ml in PBS containing PI at a final concentration of 5 µg/ml, and then incubated at room temperature for 5 minutes before FACScan analysis. PI uptake has been demonstrated to correlate with DNA fragmentation in this experimental paradigm.

25

Effect on cytokine release. An important function of monocytes/macrophages is their regulatory activity on other cellular populations of the immune system through the release of cytokines after stimulation. An ELISA to measure cytokine release is performed as follows. Human monocytes are incubated at a density of  $5 \times 10^5$  cells/ml with increasing concentrations of CRCGCL and under the same conditions, but in the absence of CRCGCL. For IL-12 production, the cells are primed overnight with IFN (100 U/ml) in presence of CRCGCL. LPS (10 ng/ml) is then added. Conditioned media are collected after 24h and kept frozen until use. Measurement of TNF-alpha, IL-10, MCP-1 and IL-8 is

09376430-081309



then performed using a commercially available ELISA kit (e.g., R & D Systems (Minneapolis, MN)) and applying the standard protocols provided with the kit.

Oxidative burst. Purified monocytes are plated in 96-w plate at  $2 \times 10^5$  cell/well.

5 Increasing concentrations of CRCGCL are added to the wells in a total volume of 0.2 ml culture medium (RPMI 1640 + 10% FCS, glutamine and antibiotics). After 3 days incubation, the plates are centrifuged and the medium is removed from the wells. To the macrophage monolayers, 0.2 ml per well of phenol red solution (140 mM NaCl, 10 mM potassium phosphate buffer pH 7.0, 5.5 mM dextrose, 0.56 mM phenol red and 19 U/ml of  
10 HRPO) is added, together with the stimulant (200 nM PMA). The plates are incubated at 37°C for 2 hours and the reaction is stopped by adding 20 µl 1N NaOH per well. The absorbance is read at 610 nm. To calculate the amount of H<sub>2</sub>O<sub>2</sub> produced by the macrophages, a standard curve of a H<sub>2</sub>O<sub>2</sub> solution of known molarity is performed for each experiment.

15 The studies described in this example tested activity in CRCGCL protein. However, one skilled in the art could easily modify the exemplified studies to test the activity of CRCGCL polynucleotides (e.g., gene therapy), agonists, and/or antagonists of CRCGCL.

### 20 ***Example 35: CRCGCL Biological Effects***

#### **Astrocyte and Neuronal Assays\***

Recombinant CRCGCL, expressed in *Escherichia coli* and purified as described above, can be tested for activity in promoting the survival, neurite outgrowth, or  
25 phenotypic differentiation of cortical neuronal cells and for inducing the proliferation of glial fibrillary acidic protein immunopositive cells, astrocytes. The selection of cortical cells for the bioassay is based on the prevalent expression of FGF-1 and FGF-2 in cortical structures and on the previously reported enhancement of cortical neuronal survival resulting from FGF-2 treatment. A thymidine incorporation assay, for example, can be  
30 used to elucidate CRCGCL's activity on these cells.

Moreover, previous reports describing the biological effects of FGF-2 (basic FGF) on cortical or hippocampal neurons *in vitro* have demonstrated increases in both neuron

survival and neurite outgrowth (Walicke, P. et al., "Fibroblast growth factor promotes survival of dissociated hippocampal neurons and enhances neurite extension." *Proc. Natl. Acad. Sci. USA* 83:3012-3016. (1986), assay herein incorporated by reference in its entirety). However, reports from experiments done on PC-12 cells suggest that these two responses are not necessarily synonymous and may depend on not only which FGF is being tested but also on which receptor(s) are expressed on the target cells. Using the primary cortical neuronal culture paradigm, the ability of CRCGCL to induce neurite outgrowth can be compared to the response achieved with FGF-2 using, for example, a thymidine incorporation assay.

10

**Fibroblast and endothelial cell assays**

Human lung fibroblasts are obtained from Clonetics (San Diego, CA) and maintained in growth media from Clonetics. Dermal microvascular endothelial cells are obtained from Cell Applications (San Diego, CA). For proliferation assays, the human lung fibroblasts and dermal microvascular endothelial cells can be cultured at 5,000 cells/well in a 96-well plate for one day in growth medium. The cells are then incubated for one day in 0.1% BSA basal medium. After replacing the medium with fresh 0.1% BSA medium, the cells are incubated with the test proteins for 3 days. Alamar Blue (Alamar Biosciences, Sacramento, CA) is added to each well to a final concentration of 10%. The cells are incubated for 4 hr. Cell viability is measured by reading in a CytoFluor fluorescence reader. For the PGE<sub>2</sub> assays, the human lung fibroblasts are cultured at 5,000 cells/well in a 96-well plate for one day. After a medium change to 0.1% BSA basal medium, the cells are incubated with FGF-2 or CRCGCL with or without IL-1 $\alpha$  for 24 hours. The supernatants are collected and assayed for PGE<sub>2</sub> by EIA kit (Cayman, Ann Arbor, MI). For the IL-6 assays, the human lung fibroblasts are cultured at 5,000 cells/well in a 96-well plate for one day. After a medium change to 0.1% BSA basal medium, the cells are incubated with FGF-2 or CRCGCL with or without IL-1 $\alpha$  for 24 hours. The supernatants are collected and assayed for IL-6 by ELISA kit (Endogen, Cambridge, MA).

30

Human lung fibroblasts are cultured with FGF-2 or CRCGCL for 3 days in basal medium before the addition of Alamar Blue to assess effects on growth of the fibroblasts.

558130-0649260

FGF-2 should show a stimulation at 10 - 2500 ng/ml which can be used to compare stimulation with CRGCL.

### Parkinson Models.

5        The loss of motor function in Parkinson's disease is attributed to a deficiency of striatal dopamine resulting from the degeneration of the nigrostriatal dopaminergic projection neurons. An animal model for Parkinson's that has been extensively characterized involves the systemic administration of 1-methyl-4 phenyl 1,2,3,6-tetrahydropyridine (MPTP). In the CNS, MPTP is taken-up by astrocytes and catabolized  
10 by monoamine oxidase B to 1-methyl-4-phenyl pyridine (MPP<sup>+</sup>) and released. Subsequently, MPP<sup>+</sup> is actively accumulated in dopaminergic neurons by the high-affinity reuptake transporter for dopamine. MPP<sup>+</sup> is then concentrated in mitochondria by the electrochemical gradient and selectively inhibits nicotinamide adenine disphosphate: ubiquinone oxidoreductionase (complex I), thereby interfering with electron transport and  
15 eventually generating oxygen radicals.

It has been demonstrated in tissue culture paradigms that FGF-2 (basic FGF) has trophic activity towards nigral dopaminergic neurons (Ferrari et al., Dev. Biol. 1989).  
~~Recently, Dr. Unsicker's group has demonstrated that administering FGF-2 in gel foam~~  
implants in the striatum results in the near complete protection of nigral dopaminergic  
20 neurons from the toxicity associated with MPTP exposure (Otto and Unsicker, J. Neuroscience, 1990).

Based on the data with FGF-2, CRGCL can be evaluated to determine whether it has an action similar to that of FGF-2 in enhancing dopaminergic neuronal survival *in vitro* and it can also be tested *in vivo* for protection of dopaminergic neurons in the  
25 striatum from the damage associated with MPTP treatment. The potential effect of CRGCL is first examined *in vitro* in a dopaminergic neuronal cell culture paradigm. The cultures are prepared by dissecting the midbrain floor plate from gestation day 14 Wistar rat embryos. The tissue is dissociated with trypsin and seeded at a density of 200,000 cells/cm<sup>2</sup> on polyorthinine-laminin coated glass coverslips. The cells are maintained in  
30 Dulbecco's Modified Eagle's medium and F12 medium containing hormonal supplements (N1). The cultures are fixed with paraformaldehyde after 8 days *in vitro* and are processed for tyrosine hydroxylase, a specific marker for dopaminergic neurons, immunohistochemical staining. Dissociated cell cultures are prepared from embryonic

668180-0649260

rats. The culture medium is changed every third day and the factors are also added at that time.

Since the dopaminergic neurons are isolated from animals at gestation day 14, a developmental time which is past the stage when the dopaminergic precursor cells are proliferating, an increase in the number of tyrosine hydroxylase immunopositive neurons would represent an increase in the number of dopaminergic neurons surviving *in vitro*. Therefore, if CRCGCL acts to prolong the survival of dopaminergic neurons, it would suggest that CRCGCL may be involved in Parkinson's Disease.

The studies described in this example tested activity in CRCGCL protein. However, one skilled in the art could easily modify the exemplified studies to test the activity of CRCGCL polynucleotides (e.g., gene therapy), agonists, and/or antagonists of CRCGCL.

### ***Example 36: The Effect of CRCGCL on the Growth of Vascular Endothelial Cells***

On day 1, human umbilical vein endothelial cells (HUVEC) are seeded at  $2-5 \times 10^4$  cells/35 mm dish density in M199 medium containing 4% fetal bovine serum (FBS), 16 units/ml heparin, and 50 units/ml endothelial cell growth supplements (ECGS, Biotechnology, Inc.). On day 2, the medium is replaced with M199 containing 10% FBS, 8 units/ml heparin. CRCGCL protein of SEQ ID NO. 2, and positive controls, such as VEGF and basic FGF (bFGF) are added, at varying concentrations. On days 4 and 6, the medium is replaced. On day 8, cell number is determined with a Coulter Counter.

An increase in the number of HUVEC cells indicates that CRCGCL may proliferate vascular endothelial cells.

The studies described in this example tested activity in CRCGCL protein. However, one skilled in the art could easily modify the exemplified studies to test the activity of CRCGCL polynucleotides (e.g., gene therapy), agonists, and/or antagonists of CRCGCL.

668780\*0219260

***Example 37: Stimulatory Effect of CRCGCL  
on the Proliferation of Vascular Endothelial Cells***

For evaluation of mitogenic activity of growth factors, the colorimetric MTS  
5 (3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)2H-  
tetrazolium) assay with the electron coupling reagent PMS (phenazine methosulfate) was  
performed (CellTiter 96 AQ, Promega). Cells are seeded in a 96-well plate (5,000  
cells/well) in 0.1 mL serum-supplemented medium and are allowed to attach overnight.  
After serum-starvation for 12 hours in 0.5% FBS, conditions (bFGF, VEGF<sub>165</sub> or  
10 CRCGCL in 0.5% FBS) with or without Heparin (8 U/ml) are added to wells for 48 hours.  
20 mg of MTS/PMS mixture (1:0.05) are added per well and allowed to incubate for 1  
hour at 37°C before measuring the absorbance at 490 nm in an ELISA plate reader.  
Background absorbance from control wells (some media, no cells) is subtracted, and seven  
wells are performed in parallel for each condition. See, Leak *et al. In Vitro Cell. Dev.*  
15 *Biol.* 30A:512-518 (1994).

The studies described in this example tested activity in CRCGCL protein.  
However, one skilled in the art could easily modify the exemplified studies to test the  
activity of CRCGCL polynucleotides (e.g., gene therapy), agonists, and/or antagonists of  
CRCGCL.

20

***Example 38: Inhibition of PDGF-induced Vascular Smooth Muscle  
Cell Proliferation Stimulatory Effect***

HAoSMC proliferation can be measured, for example, by BrdUrd incorporation.  
25 Briefly, subconfluent, quiescent cells grown on the 4-chamber slides are transfected with  
CRP or FITC-labeled AT2-3LP. Then, the cells are pulsed with 10% calf serum and 6  
mg/ml BrdUrd. After 24 h, immunocytochemistry is performed by using BrdUrd Staining  
Kit (Zymed Laboratories). In brief, the cells are incubated with the biotinylated mouse  
anti-BrdUrd antibody at 4 °C for 2 h after being exposed to denaturing solution and then  
30 incubated with the streptavidin-peroxidase and diaminobenzidine. After counterstaining  
with hematoxylin, the cells are mounted for microscopic examination, and the BrdUrd-  
positive cells are counted. The BrdUrd index is calculated as a percent of the BrdUrd-  
positive cells to the total cell number. In addition, the simultaneous detection of the  
BrdUrd staining (nucleus) and the FITC uptake (cytoplasm) is performed for individual

The studies described in this example tested activity in CRCGCL protein. However, one skilled in the art could easily modify the exemplified studies to test the activity of CRCGCL polynucleotides (e.g., gene therapy), agonists, and/or antagonists of CRCGCL.

10            This example will be used to explore the possibility that CRCGCL may stimulate lymphatic endothelial cell migration.

Endothelial cell migration assays are performed using a 48 well microchemotaxis chamber (Neuroprobe Inc., Cabin John, MD; Falk, W., et al., J. Immunological Methods 1980;33:239-247). Polyvinylpyrrolidone-free polycarbonate filters with a pore size of 8  
15 um (Nucleopore Corp. Cambridge, MA) are coated with 0.1% gelatin for at least 6 hours at room temperature and dried under sterile air. Test substances are diluted to appropriate concentrations in M199 supplemented with 0.25% bovine serum albumin (BSA), and 25 ul of the final dilution is placed in the lower chamber of the modified Boyden apparatus.

Subconfluent, early passage (2-6) HUVEC or BMEC cultures are washed and trypsinized for the minimum time required to achieve cell detachment. After placing the filter between lower and upper chamber,  $2.5 \times 10^5$  cells suspended in 50  $\mu$ l M199 containing 1% FBS are seeded in the upper compartment. The apparatus is then incubated for 5 hours at 37°C in a humidified chamber with 5% CO<sub>2</sub> to allow cell migration. After the incubation period, the filter is removed and the upper side of the filter with the non-migrated cells is scraped with a rubber policeman. The filters are fixed with methanol and stained with a Giemsa solution (Diff-Quick, Baxter, McGraw Park, IL). Migration is quantified by counting cells of three random high-power fields (40x) in each well, and all groups are performed in quadruplicate.


The studies described in this example tested activity in CRCGCL protein. However, one skilled in the art could easily modify the exemplified studies to test the activity of CRCGCL polynucleotides (e.g., gene therapy), agonists, and/or antagonists of CRCGCL.

### ***Example 40: Stimulation of Nitric Oxide Production by Endothelial Cells***

Nitric oxide released by the vascular endothelium is believed to be a mediator of  
5 vascular endothelium relaxation. Thus, CRCGCL activity can be assayed by determining  
nitric oxide production by endothelial cells in response to CRCGCL.

Nitric oxide is measured in 96-well plates of confluent microvascular endothelial  
cells after 24 hours starvation and a subsequent 4 hr exposure to various levels of a  
positive control (such as VEGF-1) and CRCGCL. Nitric oxide in the medium is  
10 determined by use of the Griess reagent to measure total nitrite after reduction of nitric  
oxide-derived nitrate by nitrate reductase. The effect of CRCGCL on nitric oxide release  
is examined on HUVEC.

Briefly, NO release from cultured HUVEC monolayer is measured with a NO-  
specific polarographic electrode connected to a NO meter (Iso-NO, World Precision  
15 Instruments Inc.) (1049). Calibration of the NO elements is performed according to the  
following equation:



The standard calibration curve is obtained by adding graded concentrations of  
KNO<sub>2</sub> (0, 5, 10, 25, 50, 100, 250, and 500 nmol/L) into the calibration solution containing  
20 KI and H<sub>2</sub>SO<sub>4</sub>. The specificity of the Iso-NO electrode to NO is previously determined by  
measurement of NO from authentic NO gas (1050). The culture medium is removed and  
HUVECs are washed twice with Dulbecco's phosphate buffered saline. The cells are then  
bathed in 5 ml of filtered Krebs-Henseleit solution in 6-well plates, and the cell plates are  
kept on a slide warmer (Lab Line Instruments Inc.) To maintain the temperature at 37°C.  
25 The NO sensor probe is inserted vertically into the wells, keeping the tip of the electrode 2  
mm under the surface of the solution, before addition of the different conditions.  
S-nitroso acetyl penicillamin (SNAP) is used as a positive control. The amount of  
released NO is expressed as picomoles per 1x10<sup>6</sup> endothelial cells. All values reported are  
means of four to six measurements in each group (number of cell culture wells). See,  
30 Leak *et al. Biochem. and Biophys. Res. Comm.* 217:96-105 (1995).

The studies described in this example tested activity in CRCGCL protein.  
However, one skilled in the art could easily modify the exemplified studies to test the

09375430-061899

activity of CRCGCL polynucleotides (e.g., gene therapy), agonists, and/or antagonists of CRCGCL.

***Example 41: Effect of CRCGCL on  
Cord Formation in Angiogenesis***

Another step in angiogenesis is cord formation, marked by differentiation of endothelial cells. This bioassay measures the ability of microvascular endothelial cells to form capillary-like structures (hollow structures) when cultured *in vitro*.

CADMEC (microvascular endothelial cells) are purchased from Cell Applications, Inc. as proliferating (passage 2) cells and are cultured in Cell Applications' CADMEC Growth Medium and used at passage 5. For the *in vitro* angiogenesis assay, the wells of a 48-well cell culture plate are coated with Cell Applications' Attachment Factor Medium (200 ml/well) for 30 min. at 37°C. CADMEC are seeded onto the coated wells at 7,500 cells/well and cultured overnight in Growth Medium. The Growth Medium is then replaced with 300 mg Cell Applications' Chord Formation Medium containing control buffer or CRCGCL (0.1 to 100 ng/ml) and the cells are cultured for an additional 48 hr. The numbers and lengths of the capillary-like chords are quantitated through use of the Boeckeler VIA-170 video image analyzer. All assays are done in triplicate.

Commercial (R&D) VEGF (50 ng/ml) is used as a positive control. b-esteradiol (1 ng/ml) is used as a negative control. The appropriate buffer (without protein) is also utilized as a control.

The studies described in this example tested activity in CRCGCL protein. However, one skilled in the art could easily modify the exemplified studies to test the activity of CRCGCL polynucleotides (e.g., gene therapy), agonists, and/or antagonists of CRCGCL.

***Example 42: Angiogenic Effect on  
Chick Chorioallantoic Membrane***

Chick chorioallantoic membrane (CAM) is a well-established system to examine angiogenesis. Blood vessel formation on CAM is easily visible and quantifiable. The ability of CRCGCL to stimulate angiogenesis in CAM can be examined.



Fertilized eggs of the White Leghorn chick (*Gallus gallus*) and the Japanese quail (*Coturnix coturnix*) are incubated at 37.8°C and 80% humidity. Differentiated CAM of 16-day-old chick and 13-day-old quail embryos is studied with the following methods.

On Day 4 of development, a window is made into the egg shell of chick eggs. The embryos are checked for normal development and the eggs sealed with cello tape. They are further incubated until Day 13. Thermanox coverslips (Nunc, Naperville, IL) are cut into disks of about 5 mm in diameter. Sterile and salt-free growth factors are dissolved in distilled water and about 3.3 mg/ 5 ml are pipetted on the disks. After air-drying, the inverted disks are applied on CAM. After 3 days, the specimens are fixed in 3% glutaraldehyde and 2% formaldehyde and rinsed in 0.12 M sodium cacodylate buffer. They are photographed with a stereo microscope [Wild M8] and embedded for semi- and ultrathin sectioning as described above. Controls are performed with carrier disks alone.

The studies described in this example tested activity in CRCGCL protein. However, one skilled in the art could easily modify the exemplified studies to test the activity of CRCGCL polynucleotides (e.g., gene therapy), agonists, and/or antagonists of CRCGCL.

#### ***Example 43: Angiogenesis Assay Using a Matrigel Implant in Mouse***

*In vivo* angiogenesis assay of CRCGCL measures the ability of an existing capillary network to form new vessels in an implanted capsule of murine extracellular matrix material (Matrigel). The protein is mixed with the liquid Matrigel at 4 degree C and the mixture is then injected subcutaneously in mice where it solidifies. After 7 days, the solid "plug" of Matrigel is removed and examined for the presence of new blood vessels. Matrigel is purchased from Becton Dickinson Labware/Collaborative Biomedical Products.

When thawed at 4 degree C the Matrigel material is a liquid. The Matrigel is mixed with CRCGCL at 150 ng/ml at 4 degree C and drawn into cold 3 ml syringes. Female C57Bl/6 mice approximately 8 weeks old are injected with the mixture of Matrigel and experimental protein at 2 sites at the midventral aspect of the abdomen (0.5 ml/site). After 7 days, the mice are sacrificed by cervical dislocation, the Matrigel plugs are

5

10

### Example 44: Rescue of Ischemia in Rabbit Lower Limb Model

15

20

25

30

reflected by the ratio of max FL: resting FL; (c) Angiographic Score - This is measured by the angiogram of collateral vessels. A score is determined by the percentage of circles in an overlaying grid that with crossing opacified arteries divided by the total number in the rabbit thigh; (d) Capillary density - The number of collateral capillaries determined in light microscopic sections taken from hindlimbs.

The studies described in this example tested activity in CRCGCL protein. However, one skilled in the art could easily modify the exemplified studies to test the activity of CRCGCL polynucleotides (e.g., gene therapy), agonists, and/or antagonists of CRCGCL.

#### ***Example 45: Effect of CRCGCL on Vasodilation***

Since dilation of vascular endothelium is important in reducing blood pressure, the ability of CRCGCL to affect the blood pressure in spontaneously hypertensive rats (SHR) is examined. Increasing doses (0, 10, 30, 100, 300, and 900 mg/kg) of the CRCGCL are administered to 13-14 week old spontaneously hypertensive rats (SHR). Data are expressed as the mean +/- SEM. Statistical analysis are performed with a paired t-test and statistical significance is defined as  $p < 0.05$  vs. the response to buffer alone.

The studies described in this example tested activity in CRCGCL protein. However, one skilled in the art could easily modify the exemplified studies to test the activity of CRCGCL polynucleotides (e.g., gene therapy), agonists, and/or antagonists of CRCGCL.

#### ***Example 46: Rat Ischemic Skin Flap Model***

The evaluation parameters include skin blood flow, skin temperature, and factor VIII immunohistochemistry or endothelial alkaline phosphatase reaction. CRCGCL expression, during the skin ischemia, is studied using in situ hybridization.

The study in this model is divided into three parts as follows:

- a) Ischemic skin
- b) Ischemic skin wounds
- c) Normal wounds

The experimental protocol includes:

a) Raising a 3x4 cm, single pedicle full-thickness random skin flap (myocutaneous flap over the lower back of the animal).

b) An excisional wounding (4-6 mm in diameter) in the ischemic skin (skin-flap).

c) Topical treatment with CRCGCL of the excisional wounds (day 0, 1, 2, 3, 4 post-wounding) at the following various dosage ranges: 1mg to 100 mg.

d) Harvesting the wound tissues at day 3, 5, 7, 10, 14 and 21 post-wounding for histological, immunohistochemical, and in situ studies.

The studies described in this example tested activity in CRCGCL protein. However, one skilled in the art could easily modify the exemplified studies to test the activity of CRCGCL polynucleotides (e.g., gene therapy), agonists, and/or antagonists of CRCGCL.

#### ***Example 47: Peripheral Arterial Disease Model***

Angiogenic therapy using CRCGCL is a novel therapeutic strategy to obtain restoration of blood flow around the ischemia in case of peripheral arterial diseases. The experimental protocol includes:

a) One side of the femoral artery is ligated to create ischemic muscle of the hindlimb, the other side of hindlimb serves as a control.

b) CRCGCL protein, in a dosage range of 20 mg - 500 mg, is delivered intravenously and/or intramuscularly 3 times (perhaps more) per week for 2-3 weeks.

c) The ischemic muscle tissue is collected after ligation of the femoral artery at 1, 2, and 3 weeks for the analysis of CRCGCL expression and histology. Biopsy is also performed on the other side of normal muscle of the contralateral hindlimb.

The studies described in this example tested activity in CRCGCL protein. However, one skilled in the art could easily modify the exemplified studies to test the activity of CRCGCL polynucleotides (e.g., gene therapy), agonists, and/or antagonists of CRCGCL.

#### ***Example 48: Ischemic Myocardial Disease Model***

CRCGCL is evaluated as a potent mitogen capable of stimulating the development of collateral vessels, and restructuring new vessels after coronary artery occlusion.

Alteration of CRCGCL expression is investigated in situ. The experimental protocol includes:

a) The heart is exposed through a left-side thoracotomy in the rat. Immediately, the left coronary artery is occluded with a thin suture (6-0) and the thorax is closed.

5        b) CRCGCL protein, in a dosage range of 20 mg - 500 mg, is delivered intravenously and/or intramuscularly 3 times (perhaps more) per week for 2-4 weeks.

c) Thirty days after the surgery, the heart is removed and cross-sectioned for morphometric and in situ analyzes.

10       The studies described in this example tested activity in CRCGCL protein. However, one skilled in the art could easily modify the exemplified studies to test the activity of CRCGCL polynucleotides (e.g., gene therapy), agonists, and/or antagonists of CRCGCL.

#### ***Example 49: Rat Corneal Wound Healing Model***

15

This animal model shows the effect of CRCGCL on neovascularization. The experimental protocol includes:

a)       Making a 1-1.5 mm long incision from the center of cornea into the stromal layer.

20       b)       Inserting a spatula below the lip of the incision facing the outer corner of the eye.

c)       Making a pocket (its base is 1-1.5 mm from the edge of the eye).

d) Positioning a pellet, containing 50ng- 5ug of CRCGCL, within the pocket.

e)       CRCGCL treatment can also be applied topically to the corneal wounds in  
25       a dosage range of 20mg - 500mg (daily treatment for five days).

The studies described in this example tested activity in CRCGCL protein. However, one skilled in the art could easily modify the exemplified studies to test the activity of CRCGCL polynucleotides (e.g., gene therapy), agonists, and/or antagonists of CRCGCL.

30

### A. Diabetic *db+ / db+* Mouse Model.

The diabetic animals have many of the characteristic features observed in Type II diabetes mellitus. Homozygous (db+/db+) mice are obese in comparison to their normal heterozygous (db+/+m) littermates. Mutant diabetic (db+/db+) mice have a single autosomal recessive mutation on chromosome 4 (db+) (Coleman *et al. Proc. Natl. Acad. Sci. USA* 77:283-293 (1982)). Animals show polyphagia, polydipsia and polyuria. Mutant diabetic mice (db+/db+) have elevated blood glucose, increased or normal insulin levels, and suppressed cell-mediated immunity (Mandel *et al., J. Immunol.* 120:1375 (1978); Debray-Sachs, M. *et al., Clin. Exp. Immunol.* 51(1):1-7 (1983); Leiter *et al., Am. J. of Pathol.* 114:46-55 (1985)). Peripheral neuropathy, myocardial complications, and microvascular lesions, basement membrane thickening and glomerular filtration abnormalities have been described in these animals (Norido, F. *et al., Exp. Neurol.* 83(2):221-232 (1984); Robertson *et al., Diabetes* 29(1):60-67 (1980); Giacomelli *et al., Lab Invest.* 40(4):460-473 (1979); Coleman, D.L., *Diabetes* 31 (Suppl):1-6 (1982)). These homozygous diabetic mice develop hyperglycemia that is resistant to insulin analogous to human type II diabetes (Mandel *et al., J. Immunol.* 120:1375-1377 (1978)).

The characteristics observed in these animals suggests that healing in this model may be similar to the healing observed in human diabetes (Greenhalgh, *et al.*, *Am. J. of Pathol.* 136:1235-1246 (1990)).

Genetically diabetic female C57BL/KsJ (db+/db+) mice and their non-diabetic (db+/+m) heterozygous littermates are used in this study (Jackson Laboratories). The animals are purchased at 6 weeks of age and are 8 weeks old at the beginning of the study. Animals are individually housed and received food and water ad libitum. All manipulations are performed using aseptic techniques. The experiments are conducted

according to the rules and guidelines of Human Genome Sciences, Inc. Institutional Animal Care and Use Committee and the Guidelines for the Care and Use of Laboratory Animals.

Wounding protocol is performed according to previously reported methods (Tsuboi, R. and Rifkin, D.B., *J. Exp. Med.* 172:245-251 (1990)). Briefly, on the day of wounding, animals are anesthetized with an intraperitoneal injection of Avertin (0.01 mg/mL), 2,2,2-tribromoethanol and 2-methyl-2-butanol dissolved in deionized water. The dorsal region of the animal is shaved and the skin washed with 70% ethanol solution and iodine. The surgical area is dried with sterile gauze prior to wounding. An 8 mm full-thickness wound is then created using a Keyes tissue punch. Immediately following wounding, the surrounding skin is gently stretched to eliminate wound expansion. The wounds are left open for the duration of the experiment. Application of the treatment is given topically for 5 consecutive days commencing on the day of wounding. Prior to treatment, wounds are gently cleansed with sterile saline and gauze sponges.

Wounds are visually examined and photographed at a fixed distance at the day of surgery and at two day intervals thereafter. Wound closure is determined by daily measurement on days 1-5 and on day 8. Wounds are measured horizontally and vertically using a calibrated Jameson caliper. Wounds are considered healed if granulation tissue is no longer visible and the wound is covered by a continuous epithelium.

CRCGCL is administered using at a range different doses of CRCGCL, from 4mg to 500mg per wound per day for 8 days in vehicle. Vehicle control groups received 50mL of vehicle solution.

Animals are euthanized on day 8 with an intraperitoneal injection of sodium pentobarbital (300mg/kg). The wounds and surrounding skin are then harvested for histology and immunohistochemistry. Tissue specimens are placed in 10% neutral buffered formalin in tissue cassettes between biopsy sponges for further processing.

Three groups of 10 animals each (5 diabetic and 5 non-diabetic controls) are evaluated: 1) Vehicle placebo control, 2) CRCGCL.

Wound closure is analyzed by measuring the area in the vertical and horizontal axis and obtaining the total square area of the wound. Contraction is then estimated by establishing the differences between the initial wound area (day 0) and that of post treatment (day 8). The wound area on day 1 is 64mm<sup>2</sup>, the corresponding size of the dermal punch. Calculations are made using the following formula:

[Open area on day 8] - [Open area on day 1] / [Open area on day 1]

Specimens are fixed in 10% buffered formalin and paraffin embedded blocks are sectioned perpendicular to the wound surface (5mm) and cut using a Reichert-Jung microtome. Routine hematoxylin-eosin (H&E) staining is performed on cross-sections of bisected wounds. Histologic examination of the wounds are used to assess whether the healing process and the morphologic appearance of the repaired skin is altered by treatment with CRCGCL. This assessment included verification of the presence of cell accumulation, inflammatory cells, capillaries, fibroblasts, re-epithelialization and epidermal maturity (Greenhalgh, D.G. *et al.*, *Am. J. Pathol.* 136:1235 (1990)). A calibrated lens micrometer is used by a blinded observer.

Tissue sections are also stained immunohistochemically with a polyclonal rabbit anti-human keratin antibody using ABC Elite detection system. Human skin is used as a positive tissue control while non-immune IgG is used as a negative control. Keratinocyte growth is determined by evaluating the extent of reepithelialization of the wound using a calibrated lens micrometer.

Proliferating cell nuclear antigen/cyclin (PCNA) in skin specimens is demonstrated by using anti-PCNA antibody (1:50) with an ABC Elite detection system. Human colon cancer served as a positive tissue control and human brain tissue is used as a negative tissue control. Each specimen included a section with omission of the primary antibody and substitution with non-immune mouse IgG. Ranking of these sections is based on the extent of proliferation on a scale of 0-8, the lower side of the scale reflecting slight proliferation to the higher side reflecting intense proliferation.

Experimental data are analyzed using an unpaired t test. A p value of < 0.05 is considered significant.

### ***B. Steroid Impaired Rat Model***

The inhibition of wound healing by steroids has been well documented in various *in vitro* and *in vivo* systems (Wahl, S.M. Glucocorticoids and Wound healing. In: Anti-Inflammatory Steroid Action: Basic and Clinical Aspects. 280-302 (1989); Wahl, S.M.*et al.*, *J. Immunol.* 115: 476-481 (1975); Werb, Z. *et al.*, *J. Exp. Med.* 147:1684-1694 (1978)). Glucocorticoids retard wound healing by inhibiting angiogenesis, decreasing



vascular permeability ( Ebert, R.H., *et al.*, *An. Intern. Med.* 37:701-705 (1952)), fibroblast proliferation, and collagen synthesis (Beck, L.S. *et al.*, *Growth Factors.* 5: 295-304 (1991); Haynes, B.F. *et al.*, *J. Clin. Invest.* 61: 703-797 (1978)) and producing a transient reduction of circulating monocytes (Haynes, B.F., *et al.*, *J. Clin. Invest.* 61: 703-797 (1978); Wahl, S. M., "Glucocorticoids and wound healing", *In: Antiinflammatory Steroid Action: Basic and Clinical Aspects*, Academic Press, New York, pp. 280-302 (1989)). The systemic administration of steroids to impaired wound healing is a well establish phenomenon in rats (Beck, L.S. *et al.*, *Growth Factors.* 5: 295-304 (1991); Haynes, B.F., *et al.*, *J. Clin. Invest.* 61: 703-797 (1978); Wahl, S. M., "Glucocorticoids and wound healing", *In: Antiinflammatory Steroid Action: Basic and Clinical Aspects*, Academic Press, New York, pp. 280-302 (1989); Pierce, G.F. *et al.*, *Proc. Natl. Acad. Sci. USA* 86: 2229-2233 (1989)).

To demonstrate that CRCGCL can accelerate the healing process, the effects of multiple topical applications of CRCGCL on full thickness excisional skin wounds in rats in which healing has been impaired by the systemic administration of methylprednisolone is assessed.

Young adult male Sprague Dawley rats weighing 250-300 g (Charles River Laboratories) are used in this example. The animals are purchased at 8 weeks of age and are 9 weeks old at the beginning of the study. The healing response of rats is impaired by the systemic administration of methylprednisolone (17mg/kg/rat intramuscularly) at the time of wounding. Animals are individually housed and received food and water *ad libitum*. All manipulations are performed using aseptic techniques. This study is conducted according to the rules and guidelines of Human Genome Sciences, Inc. Institutional Animal Care and Use Committee and the Guidelines for the Care and Use of Laboratory Animals.

The wounding protocol is followed according to section A, above. On the day of wounding, animals are anesthetized with an intramuscular injection of ketamine (50 mg/kg) and xylazine (5 mg/kg). The dorsal region of the animal is shaved and the skin washed with 70% ethanol and iodine solutions. The surgical area is dried with sterile gauze prior to wounding. An 8 mm full-thickness wound is created using a Keyes tissue punch. The wounds are left open for the duration of the experiment. Applications of the testing materials are given topically once a day for 7 consecutive days commencing on the

day of wounding and subsequent to methylprednisolone administration. Prior to treatment, wounds are gently cleansed with sterile saline and gauze sponges.

Wounds are visually examined and photographed at a fixed distance at the day of wounding and at the end of treatment. Wound closure is determined by daily measurement on days 1-5 and on day 8. Wounds are measured horizontally and vertically using a calibrated Jameson caliper. Wounds are considered healed if granulation tissue is no longer visible and the wound is covered by a continuous epithelium.

CRCGCL is administered using at a range different doses of CRCGCL, from 4mg to 500mg per wound per day for 8 days in vehicle. Vehicle control groups received 50mL of vehicle solution.

Animals are euthanized on day 8 with an intraperitoneal injection of sodium pentobarbital (300mg/kg). The wounds and surrounding skin are then harvested for histology. Tissue specimens are placed in 10% neutral buffered formalin in tissue cassettes between biopsy sponges for further processing.

Four groups of 10 animals each (5 with methylprednisolone and 5 without glucocorticoid) are evaluated: 1) Untreated group 2) Vehicle placebo control 3) CRCGCL treated groups.

Wound closure is analyzed by measuring the area in the vertical and horizontal axis and obtaining the total area of the wound. Closure is then estimated by establishing the differences between the initial wound area (day 0) and that of post treatment (day 8). The wound area on day 1 is 64mm<sup>2</sup>, the corresponding size of the dermal punch. Calculations are made using the following formula:

$$[\text{Open area on day 8}] - [\text{Open area on day 1}] / [\text{Open area on day 1}]$$

Specimens are fixed in 10% buffered formalin and paraffin embedded blocks are sectioned perpendicular to the wound surface (5mm) and cut using an Olympus microtome. Routine hematoxylin-eosin (H&E) staining is performed on cross-sections of bisected wounds. Histologic examination of the wounds allows assessment of whether the healing process and the morphologic appearance of the repaired skin is improved by treatment with CRCGCL. A calibrated lens micrometer is used by a blinded observer to determine the distance of the wound gap.

Experimental data are analyzed using an unpaired t test. A p value of  $< 0.05$  is considered significant.

The studies described in this example tested activity in CRCGCL protein. However, one skilled in the art could easily modify the exemplified studies to test the activity of CRCGCL polynucleotides (e.g., gene therapy), agonists, and/or antagonists of CRCGCL.

### ***Example 51: Lymphadema Animal Model***

10 The purpose of this experimental approach is to create an appropriate and consistent lymphedema model for testing the therapeutic effects of CRCGCL in lymphangiogenesis and re-establishment of the lymphatic circulatory system in the rat hind limb. Effectiveness is measured by swelling volume of the affected limb, quantification of the amount of lymphatic vasculature, total blood plasma protein, and  
15 histopathology. Acute lymphedema is observed for 7-10 days. Perhaps more importantly, the chronic progress of the edema is followed for up to 3-4 weeks.

Prior to beginning surgery, blood sample is drawn for protein concentration analysis. Male rats weighing approximately ~350g are dosed with Pentobarbital. Subsequently, the right legs are shaved from knee to hip. The shaved area is swabbed  
20 with gauze soaked in 70% EtOH. Blood is drawn for serum total protein testing. Circumference and volumetric measurements are made prior to injecting dye into paws after marking 2 measurement levels (0.5 cm above heel, at mid-pt of dorsal paw). The intradermal dorsum of both right and left paws are injected with 0.05 ml of 1% Evan's Blue. Circumference and volumetric measurements are then made following injection of  
25 dye into paws.

Using the knee joint as a landmark, a mid-leg inguinal incision is made circumferentially allowing the femoral vessels to be located. Forceps and hemostats are used to dissect and separate the skin flaps. After locating the femoral vessels, the lymphatic vessel that runs along side and underneath the vessel(s) is located. The main  
30 lymphatic vessels in this area are then electrically coagulated or suture ligated.

Using a microscope, muscles in back of the leg (near the semitendinosus and adductors) are bluntly dissected. The popliteal lymph node is then located. The 2 proximal and 2 distal lymphatic vessels and distal blood supply of the popliteal node are

0037430-08189

then and ligated by suturing. The popliteal lymph node, and any accompanying adipose tissue, is then removed by cutting connective tissues.

Care is taken to control any mild bleeding resulting from this procedure. After lymphatics are occluded, the skin flaps are sealed by using liquid skin (Vetbond) (AJ Buck). The separated skin edges are sealed to the underlying muscle tissue while leaving a gap of ~0.5 cm around the leg. Skin also may be anchored by suturing to underlying muscle when necessary.

To avoid infection, animals are housed individually with mesh (no bedding). Recovering animals are checked daily through the optimal edematous peak, which typically occurred by day 5-7. The plateau edematous peak are then observed. To evaluate the intensity of the lymphedema, the circumference and volumes of 2 designated places on each paw before operation and daily for 7 days are measured. The effect plasma proteins on lymphedema is determined and whether protein analysis is a useful testing perimeter is also investigated. The weights of both control and edematous limbs are evaluated at 2 places. Analysis is performed in a blind manner.

**Circumference Measurements:** Under brief gas anesthetic to prevent limb movement, a cloth tape is used to measure limb circumference. Measurements are done at the ankle bone and dorsal paw by 2 different people then those 2 readings are averaged. Readings are taken from both control and edematous limbs.

**Volumetric Measurements:** On the day of surgery, animals are anesthetized with Pentobarbital and are tested prior to surgery. For daily volumetrics animals are under brief halothane anesthetic (rapid immobilization and quick recovery), both legs are shaved and equally marked using waterproof marker on legs. Legs are first dipped in water, then dipped into instrument to each marked level then measured by Buxco edema software(Chen/Victor). Data is recorded by one person, while the other is dipping the limb to marked area.

**Blood-plasma protein measurements:** Blood is drawn, spun, and serum separated prior to surgery and then at conclusion for total protein and Ca<sup>2+</sup> comparison.

**Limb Weight Comparison:** After drawing blood, the animal is prepared for tissue collection. The limbs are amputated using a quillitine, then both experimental and control legs are cut at the ligature and weighed. A second weighing is done as the tibio-cacaneal joint is disarticulated and the foot is weighed.

**Histological Preparations:** The transverse muscle located behind the knee (popliteal) area is dissected and arranged in a metal mold, filled with freezeGel, dipped into cold methylbutane, placed into labeled sample bags at - 80EC until sectioning. Upon sectioning, the muscle is observed under fluorescent microscopy for lymphatics..

5        The studies described in this example tested activity in CRCGCL protein. However, one skilled in the art could easily modify the exemplified studies to test the activity of CRCGCL polynucleotides (e.g., gene therapy), agonists, and/or antagonists of CRCGCL.

10        ***Example 52: Suppression of TNF alpha-induced adhesion molecule expression by CRCGCL***

15        The recruitment of lymphocytes to areas of inflammation and angiogenesis involves specific receptor-ligand interactions between cell surface adhesion molecules (CAMs) on lymphocytes and the vascular endothelium. The adhesion process, in both normal and pathological settings, follows a multi-step cascade that involves intercellular adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule-1 (VCAM-1), and endothelial leukocyte adhesion molecule-1 (E-selectin) expression on endothelial cells (EC). The expression of these molecules and others on the vascular endothelium  
20        determines the efficiency with which leukocytes may adhere to the local vasculature and extravasate into the local tissue during the development of an inflammatory response. The local concentration of cytokines and growth factor participate in the modulation of the expression of these CAMs.

25        Tumor necrosis factor alpha (TNF-a), a potent proinflammatory cytokine, is a stimulator of all three CAMs on endothelial cells and may be involved in a wide variety of inflammatory responses, often resulting in a pathological outcome.

30        The potential of CRCGCL to mediate a suppression of TNF-a induced CAM expression can be examined. A modified ELISA assay which uses ECs as a solid phase absorbent is employed to measure the amount of CAM expression on TNF-a treated ECs when co-stimulated with a member of the FGF family of proteins.

To perform the experiment, human umbilical vein endothelial cell (HUVEC) cultures are obtained from pooled cord harvests and maintained in growth medium (EGM-2; Clonetics, San Diego, CA) supplemented with 10% FCS and 1%

penicillin/streptomycin in a 37 degree C humidified incubator containing 5% CO<sub>2</sub>. HUVECs are seeded in 96-well plates at concentrations of  $1 \times 10^4$  cells/well in EGM medium at 37 degree C for 18-24 hrs or until confluent. The monolayers are subsequently washed 3 times with a serum-free solution of RPMI-1640 supplemented with 100 U/ml penicillin and 100 mg/ml streptomycin, and treated with a given cytokine and/or growth factor(s) for 24 h at 37 degree C. Following incubation, the cells are then evaluated for CAM expression.

Human Umbilical Vein Endothelial cells (HUVECs) are grown in a standard 96 well plate to confluence. Growth medium is removed from the cells and replaced with 90 ul of 199 Medium (10% FBS). Samples for testing and positive or negative controls are added to the plate in triplicate (in 10 ul volumes). Plates are incubated at 37 degree C for either 5 h (selectin and integrin expression) or 24 h (integrin expression only). Plates are aspirated to remove medium and 100  $\mu$ l of 0.1% paraformaldehyde-PBS(with Ca++ and Mg++) is added to each well. Plates are held at 4°C for 30 min.

Fixative is then removed from the wells and wells are washed 1X with PBS(+Ca,Mg)+0.5% BSA and drained. Do not allow the wells to dry. Add 10  $\mu$ l of diluted primary antibody to the test and control wells. Anti-ICAM-1-Biotin, Anti-VCAM-1-Biotin and Anti-E-selectin-Biotin are used at a concentration of 10  $\mu$ g/ml (1:10 dilution of 0.1 mg/ml stock antibody). Cells are incubated at 37°C for 30 min. in a humidified environment. Wells are washed X3 with PBS(+Ca,Mg)+0.5% BSA.

Then add 20  $\mu$ l of diluted ExtrAvidin-Alkaline Phosphatase (1:5,000 dilution) to each well and incubated at 37°C for 30 min. Wells are washed X3 with PBS(+Ca,Mg)+0.5% BSA. 1 tablet of p-Nitrophenol Phosphate pNPP is dissolved in 5 ml of glycine buffer (pH 10.4). 100  $\mu$ l of pNPP substrate in glycine buffer is added to each test well. Standard wells in triplicate are prepared from the working dilution of the ExtrAvidin-Alkaline Phosphatase in glycine buffer:  $1:5,000$  ( $10^0$ )  $> 10^{-0.5} > 10^{-1} > 10^{-1.5}$ . 5  $\mu$ l of each dilution is added to triplicate wells and the resulting AP content in each well is 5.50 ng, 1.74 ng, 0.55 ng, 0.18 ng. 100  $\mu$ l of pNNP reagent must then be added to each of the standard wells. The plate must be incubated at 37°C for 4h. A volume of 50  $\mu$ l of 3M NaOH is added to all wells. The results are quantified on a plate reader at 405 nm. The background subtraction option is used on blank wells filled with glycine buffer only. The template is set up to indicate the concentration of AP-conjugate in each standard well [

5.50 ng; 1.74 ng; 0.55 ng; 0.18 ng]. Results are indicated as amount of bound AP-conjugate in each sample.

5 The studies described in this example tested activity in CRCGCL protein. However, one skilled in the art could easily modify the exemplified studies to test the activity of CRCGCL polynucleotides (e.g., gene therapy), agonists, and/or antagonists of CRCGCL.

10 It will be clear that the invention may be practiced otherwise than as particularly described in the foregoing description and examples. Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, are within the scope of the appended claims.

15 The entire disclosure of each document cited (including patents, patent applications, journal articles, abstracts, laboratory manuals, books, or other disclosures) in the Background of the Invention, Detailed Description, and Examples is hereby incorporated herein by reference. Moreover, the sequence listing from U.S application Serial Nos. 60/078,563, filed March 19, 1998, 60/086,505, filed May 22, 1998, and 09/263,626, filed March 5, 1999, are herein incorporated by reference.

09376430-06492660